



Isolation and Identification of Bacteria from Wound Sepsis among Patients at Obigwe Orthopedic Clinic, Amorji Ubaha, Okigwe

**Chizurum P. Christian^a, George C. Njoku^{b*}, Chibuzo V. Alisigwe^c,
David U. Iloanusi^b, Emmanuel K. Amanze^c, Prisca C. Aririguzo^d
and Isabel C. Nwagu^b**

^a Department of Microbiology, Faculty of Biological Science, Abia State University, Uturu, PMB-2000
Abia State, Nigeria.

^b Department of Biochemistry, College of Natural Sciences, Michael Okpara University of Agriculture,
Umudike, PMB-7267, Umuahia, Abia State, Nigeria.

^c Department of Microbiology, College of Natural Sciences, Michael Okpara University of Agriculture,
Umudike, PMB-7267, Umuahia, Abia State, Nigeria.

^d Department of Public Health, School of Health Technology, Federal University of Technology Owerri
PMB-1526, Owerri, Imo State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/SAJRM/2022/v12i230267

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85290>

Original Research Article

Received 12 December 2021

Accepted 14 February 2022

Published 15 February 2022

ABSTRACT

Infection in wound delays healing and may cause wound breakdown, herniation, and complete wound dehiscence. This study investigated the isolation and identification of bacteria associated with wound sepsis. A total number of ten (10) swab samples were collected at random from the wound surface of patients with infected wounds. The swabs with the samples were analyzed using standard microbiological procedures. Bacterial isolates were characterized using their colonial, microscopic, and biochemical properties. Identification was with reference to Bergey's Manual. It was observed that *Escherichia coli* is the most frequently occurring isolates with a percentage occurrence of (25%) followed by *Staphylococcus aureus* with a percentage of (20%), then *Pseudomonas aeruginosa* (20%), *Klebsiella pneumonia* (20%), and *Streptococcus pyogenes*

*Corresponding author: Email: njokugeorgechigozie@gmail.com;

(15%). The overall distribution of bacterial isolates from the wound samples indicated that the wound samples from the male patients have a higher percentage distribution of (80%) than the samples obtained from the female patients with (20%). This study revealed that various bacteria, including opportunistic bacteria, are found in different wounds. Some of the wounds were infected with more than one bacteria species at a time. Although complete eradication of wound infections is not possible, however, by adopting prompt, clean surgical procedures, proper care of wounds, and antibiotics, the incidence of wound infection may be limited to a minimum.

Keywords: *Antibiotics; Escherichia coli; Klebsiella pneumonia; Pseudomonas aeruginosa; Staphylococcus aureus; Wound sepsis.*

1. INTRODUCTION

Wounds refer to injuries that break the skin, either breaking the cell membranes or a laceration, usually resulting in damage to underlying tissues [1]. "They are usually a result of violence, accidents, and surgery leading to the loss of the integrity of the first line of defense (skin) and exposure of the subcutaneous layer. Wound provides a moist, warm, and nutritive environment conducive to microbial and colonization, which results in infection" [1]. "Wounds make sepsis more expected following the entry of microorganisms into the wound if not treated well. Sepsis is a toxic condition resulting from the spread of bacteria or their toxic products from a focus of infection" [2]. "Wound sepsis occurs when virulence factors expressed by one or more microorganisms in a wound outcompete the host's natural immune system. The subsequent invasion and dissemination of microorganisms in viable tissue provoke a series of local and systemic host responses. Characteristic local responses are a purulent discharge or painful spreading erythema indicative of cellulitis around a wound" [3].

Wound infection poses a global health challenge leading to delay in wound healing and surgical complications like dehiscence or wound breakdown [4]. "Despite the latest advances in technology, wound infection has been identified as the most common nosocomial infection, especially in patients undergoing surgery [4], resulting in a prolonged hospital stay, increased trauma care, treatment costs, and more demanding general wound management practices" [1].

"Evidence from bacteriological studies has shown that many bacteria species are universally involved in wound infection. This variation is due to geographical locations, bacteria resident on the skin, clothing at the wound site, the time between the wound, and examination" [3].

Several bacterial genera cause wound infections [5]. "Isolates found in cases of wound infections include *Staphylococcus aureus*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Klebsiella aerogenes*, *Escherichia coli*, *Staphylococcus epidermidis*, *Streptococcus pyogenes*, and *Streptococcus faecalis*, *Candida albicans* and *Candida tropicalis* have also been implicated as etiological agents" [6]. These microorganisms are transmitted into the wound through various ways, including direct contact with the organisms, which are either transferred from surgical equipment airborne dispersal from the environment contaminated with microorganisms deposited onto the wounds. Contamination could also occur through one-self by physical migration of the patient's endogenous flora on the skin, mucous membrane, or gastrointestinal tract to the surgical site [7].

"In treating wound infections, several antibiotics are used. Ideally, proper antibiotics are given after the culture and sensitivity of isolates from the wound swab, pus, or infected tissue. Unfortunately, this practice is uncommon among physicians, especially in developing countries. Consequently, improper use of broad and narrow-spectrum antibiotics leads to drug resistance" [8]. "The control of wound infections has become more challenging due to pervasive bacterial resistance to antibiotics and increased infections caused by methicillin-resistant *Staphylococcus aureus* (MRSA), extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria, and polymicrobial flora" [3]. Therefore, the effects of antimicrobial resistance cannot be over-emphasized, and it can increase complications and costs associated with procedures and treatment [9].

Knowledge of the causative agents of wound infection has proven helpful in selecting empiric antimicrobial therapy and on infection control measures taken in health institutions. Therefore, this study aims to identify and characterize

isolates from wound sepsis to select the proper antimicrobial agent needed for their eradication.

2. MATERIALS AND METHODS

2.1 Sample Collection

A total of 10 wound swab samples were collected from patients at Obigwe Orthopaedic Clinic, Amorji Ubaha, Okigwe, using labeled sterile swab sticks. The wound was first cleansed with normal saline (irrigation) to remove superficial debris and a swab of purulent exudates; wound secretion was aseptically obtained using sterile cotton swab sticks for each patient. The wound specimen was collected on moistened cotton swab sticks to swab across the wound surface in a zigzag motion, without contaminating the skin commensals. The swabs taken were immersed in a small tube and labeled. Soon after collection, each sample was taken to the laboratory of microbiology, Abia State University, Uturu.

2.2 Preparation of Samples

On getting to the laboratory, the samples were stored immediately in a refrigerator at a temperature of 4°C to prevent the microbes' proliferation.

2.3 Culture of Samples

The collected swab specimens were streaked on freshly prepared nutrient agar and placed in the incubator. The swab specimens were incubated for 24 hours at 37°C; the plates were examined for growth. Isolates from the labeled nutrient agar plates were further sub-cultured into MacConkey agar (selective for only gram-negative enteric organisms) and blood agar using a sterile wire loop, and pure cultures were obtained. According to manufacturers' instructions, all growth media used in his process were prepared and autoclaved appropriately.

2.4 Morphological Characterization, Gram Reaction, and Microscopic Examination of Isolates

Before isolating the pathogens involved in wound sepsis, macroscopic examinations (for cultural characteristics such as colony size, pigmentation, elevation, and margin) were studied after incubation. Different morphological features of the yielded colonies, including color, size, shape, margin elevation, and texture, were recorded.

Following the macroscopic examination, the Gram staining technique was used in this experiment to help identify pathogens by their Gram reaction. According to Cheesbrough [10], the procedure was carried out as follows; the smear was prepared from an overnight culture on a clean, grease-free glass slide. The smear was left to air dry. After that, the slide was rapidly passed three times through the flame of a Bunsen burner and then allowed to cool before staining. The fixed smear was covered with Crystal violet stain for 60 seconds and then washed in slow-running clean tap water. The smear was tipped off all the water before being covered with Lugol's iodine (mordant) for 60 seconds. The stain was washed off in slow-running clean tap water. Acetone was used to decolorize the smear rapidly and immediately washed in slow-running clean tap water. Finally, the smear was covered with Safranin stain for 2 minutes and washed in slow-running clean tap water. The back of the slide was wiped clean and placed in a draining rack for the smear to air dry. A drop of oil was added to the dried smear got examined under the microscope with x100 oil immersion objective.

2.5 Biochemical Tests

After microscopic examination, the isolates were characterized using the following biochemical tests: catalase test, coagulase test, citrate test, oxidase test, and indole test using the standard operating procedure in Cheesbrough [10]. After biochemical tests, the ninth edition of Bergey's Manual of Determinative Bacteriology was used to identify the pathogens from wound sepsis as *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, and *Streptococcus pyogenes*.

2.6 Percentage Occurrence of Isolates

The percentage occurrence of isolates was determined by dividing the number of times a particular isolate appeared by the total number of isolates and then multiplied by hundred (%).

3. RESULTS

3.1 Morphological Identification, Biochemical Identification, Gram Reaction of Bacterial Isolates from the Wound Sepsis

Table 1 shows the Morphological identification, Biochemical Identification, and Gram Reaction of

bacterial isolates from the wound sepsis. The bacterial isolates obtained from this study include; *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, and *Streptococcus pyogenes*.

3.2 Percentage Occurrence of the Isolates from the Wound Sepsis

Table 2 shows the percentage occurrence of the isolates from the wound sepsis. It was observed that *E. coli* is the most frequently occurring isolate with a percentage occurrence of 5(25%), followed by *Staphylococcus aureus* 4(20%), *Pseudomonas aeruginosa* 4(20%), *Klebsiella pneumoniae* 4(20%), and *Streptococcus pyogenes* 3(15%).

3.3 Distribution of Bacteria Isolates from the Wound Swab Samples

Table 3 shows the distribution of bacteria isolates from the wound swab samples. The overall distribution of bacterial isolates from the wound samples indicated that the wound samples from the male patients have a higher percentage distribution of (80%) than the samples obtained from the female patients with (20%).

4. DISCUSSION

“Wound sepsis is one of the most common infections in which bacterial flora proliferates and delays wound healing. It is a significant complication of wounds with a significant increase in costs, morbidity, and potential mortality” [2]. “Wound sepsis occurs when virulence factors expressed by one or more microorganisms in a wound outcompete the host’s natural immune system. The subsequent invasion and dissemination of microorganisms in viable tissue provoke a series of local and systemic host responses” [11]. “Wound sepsis is one of the most common hospital-acquired infections, a significant cause of morbidity, and accounts for 70-80% mortality” [12]. “The development of such infections represents delayed healing, causes anxiety and discomfort for patients, longer stays at hospitals, and significantly adds to the cost of healthcare services” [13]. This study evaluated the bacteriology of patients’ infected wounds. In this study, the predominant bacteria isolated were *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, and *Streptococcus pyogenes* (Table 1). This conforms with earlier reports by Mahat et

al. [14], which indicated that “*S. aureus*, *Streptococcus spp*, *Enterococcus spp*, *Pseudomonas spp*, *E. coli*, *Acinetobacter Baumannii*, *E. aerogens*, *K. pneumoniae*, *Streptococcus pyogenes*, and *C. freundii* were among the most commonly isolated bacteria from the wound.” The report of Adeyemi (2012) in the same vein agrees with the findings of this work that the predominant bacteria isolated were *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, *Streptococcus species*, *Bacillus species*, *Klebsiella aerogenes*, *Klebsiella oxytoca*, *Enterobacter species*, *Proteus mirabilis*, *Proteus Vulgaris*, *Pseudomonas aeruginosa*, and *Streptococcus pyogenes*. This is also similar to what was earlier reported by Aizza et al. [15] and Bessa et al. [16], in their studies to explore the common bacterial pathogens responsible for wound infection and determine the antibiotic susceptibility pattern of the bacterial isolates.

From the findings in this study, it was observed that *Escherichia coli* is the most frequently occurring isolates with a percentage occurrence of (25%) followed by *Staphylococcus aureus* with a percentage of (20%) then *Pseudomonas aeruginosa* (20%), *Klebsiella pneumoniae* (20%), and *Streptococcus pyogenes* (15%). “In all, Gram-negative bacilli are more prevalent than Gram-positive bacteria. The prevalence of 70% and 30%, respectively, is similar to the previously reported prevalence of 67% and 32%” by [11]. Our report shows that *Escherichia coli* is the most common isolated Gram-negative bacilli, while *S. aureus* is Gram-positive bacteria. This is consistent with earlier reports [17]. The trend in the rate of isolation of Gram-negative bacilli over the study period is that of an increase, while that of Gram-positive bacteria is one of decrease. Although these changes are not statistically significant, they are clinically meaningful. The two most critical bacterial pathogens are *S. aureus* and *E. coli*. This report shows an increase in their rates of isolation. This also is of clinical importance. The overall distribution of bacterial isolates from the wound samples indicated that the wound samples from the male patients have a higher percentage distribution of (80%) as compared to the samples obtained from the female patients (20%) (Table 3). This is similar to the findings of Aizza et al. [15], who reported that “male patients with wound infection were 20% higher than female patients.” “The higher growth positive cases in male patients (58.8%) than in females (42.2%) was observed in this study and was supported by the similar

Table 1. Characterization and identification of isolates from Wound Sepsis

Specimen	Shape	Size	Margin	Color	Elevation	Texture	Gram reaction	CA	CI	CO	IN	OX	Probable isolates
Male 1A	Circular	Small	Entire	Dark pink	Flat	Moist	Negative bacilli (pairs)	+	-	-	+	-	<i>E. coli</i>
Male 1B	Circular	Small	Entire	Golden-yellow	Convex	Smooth	Positive cocci (cluster)	+	+	+	-	-	<i>Staphylococcus aureus</i>
Male 2A	Circular	medium	Entire	off-white	Flat	Smooth	Negative bacilli (chains)	+	+	-	-	+	<i>Pseudomonas aeruginosa</i>
Male 2B	Ovoid	Small	Entire	Grayish white	Raised	Smooth	Positive cocci (chains)	-	-	-	-	-	<i>Streptococcus pyogenes</i>
Male 3A	Circular	medium	Entire	Pink	Convex	Mucoid	Negative bacilli (pairs)	+	+	-	-	-	<i>Klebsiella pneumoniae</i>
Male 3B	Circular	Small	Entire	Golden-yellow	Convex	Smooth	Positive cocci (clusters)	+	+	+	-	-	<i>Staphylococcus aureus</i>
Male 4A	Circular	Small	Entire	Dark pink	Flat	Moist	Negative bacilli (pairs)	+	-	-	+	-	<i>E. coli</i>
Male 4B	Circular	Large	Entire	Creamy	Convex	Mucoid	Negative bacilli (chains)	+	+	-	-	-	<i>Klebsiella pneumoniae</i>
Male 5A	Circular	Small	Entire	Off white	Flat	Smooth	Negative bacilli (pairs)	+	+	-	-	+	<i>Pseudomonas aeruginosa</i>
Male 5B	Ovoid	Small	Entire	Grayish white	Raised	Smooth	Positive cocci (chains)	-	-	-	-	-	<i>Streptococcus pyogenes</i>
Male 6A	Circular	Medium	Entire	Pink	Convex	Mucoid	Negative bacilli (short chains)	+	+	-	-	-	<i>Klebsiella pneumoniae</i>
Male 6B	Ovoid	Small	Entire	Grayish white	Raised	Smooth	Positive cocci (chains)	-	-	-	-	-	<i>Streptococcus pyogenes</i>
Male 7A	Circular	Small	Entire	Dark pink	Flat	Moist	Negative bacilli (clusters)	+	-	-	+	-	<i>E. coli</i>
Male 7B	Circular	Large	Entire	Green	Flat	Mucoid	Negative bacilli (pairs)	+	+	-	-	+	<i>Pseudomonas aeruginosa</i>
Male 8A	Circular	Small	Entire	Dark pink	Flat	Moist	Negative bacilli (pairs)	+	-	-	+	-	<i>E. coli</i>
Male 8B	Circular	Small	Entire	Golden yellow	Convex	Smooth	Positive cocci (clusters)	+	+	+	-	-	<i>Staphylococcus aureus</i>
Female 1A	Circular	Medium	Entire	Pink	Convex	Mucoid	Negative bacilli (clusters)	+	+	-	-	-	<i>Klebsiella pneumoniae</i>
Female 1B	Circular	Small	Entire	Golden yellow	Convex	Smooth	Positive cocci (clusters)	+	+	+	-	-	<i>Staphylococcus aureus</i>
Female 2A	Circular	Small	Entire	Dark pink	Flat	Moist	Negative bacilli (pairs)	+	-	-	-	+	<i>E. coli</i>
Female 2B	Circular	Large	Entire	Green	Flat	Mucoid	Negative bacilli (pairs)	+	+	-	-	+	<i>Pseudomonas aeruginosa</i>

KEY: a = MacConkey agar, b = Blood agar, + = Positive, - = Negative, CA. = Catalase, CI. = Citrate, CO. = Coagulase, IN. = Indole, OX=Oxidase,MR.=MethylRed

Table 2. Percentage occurrence of bacterial isolates from wound sepsis

Isolates	No of isolates	Percentage occurrence (%)
<i>Escherichia coli</i>	5	25
<i>Staphylococcus aureus</i>	4	20
<i>Pseudomonas aeruginosa</i>	4	20
<i>Klebsiella pneumoniae</i>	4	20
<i>Streptococcus pyogenes</i>	3	15
Total	20	100

Table 3. Distribution of Bacteria Isolates from the wound swab samples

Isolates	No of isolates	Percentage distribution of isolates (%)	
		Male	Female
<i>Escherichia coli</i>	5	4(80%)	1(20%)
<i>Staphylococcus aureus</i>	4	3(75%)	1(25%)
<i>Pseudomonas aeruginosa</i>	4	3(75%)	1(25%)
<i>Klebsiella pneumoniae</i>	4	3(75%)	1(25%)
<i>Streptococcus pyogenes</i>	3	3(100%)	0(0%)
Total	20	16(80%)	4(20%)

studies" carried out" by Mahat et al. [14]. The slight difference noted is due to our social behavior where males are given superiority to the female and, if they get diseased, are brought immediately to hospitals compared to females for treatment.

5. CONCLUSION

This study concluded that contracting wound infection remains an ongoing problem. The main culprit for the wound infection is trivial organisms like *Staphylococcus aureus*, *Escherichia coli*, *Streptococcus pyogenes*, *Pseudomonas aeruginosa*, *Klebsiella*, etc. Although complete eradication of wound infections is not possible, the incidence of wound infection may be limited to a minimum by taking preventive measures and adopting prompt, clean surgical procedures and proper care of wounds. Otherwise, wound infections may lead to the morbidity and mortality of a high count, especially in children.

6. RECOMMENDATIONS

The result obtained from this study will be helpful for policymakers in evaluating the infection control measures in hospitals. Routine antimicrobial susceptibility testing before antibiotic administration is highly recommended since it can play a pivotal role in limiting wound infections to a minimum. As the emergence of drug-resistant bacteria is less likely when there is empirical drug therapy, it fastens the process of wound healing; hence the appropriate use of drugs decreases the cost of wound infection treatment.

CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical permission has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Miron RJ. Platelet-rich fibrin and soft tissue wound healing: A systematic review. *Tissue Eng. part. B: Rev.* 2017;23:83–99.
2. Russell J A. Shock Syndrome related to sepsis. Goldman, I., Schafer A. I, eds *Medicine*, 24th ed. Philadelphia, Pa: Saunders Elsevier. 2011; Chap 108.
3. Nasser A. *Staphylococcus aureus* versus neutrophil: Scrutiny of ancient combat. *Microbes' pathogen.* 2019;131:259–269.
4. Kashfi M. The prevalence of aminoglycoside-modifying enzyme genes among *Pseudomonas aeruginosa* strains isolated from burn patients. *Achieved Clin. Infect. and Disease.* 2017;12:15-18.
5. Copeland-Halperin LR, Kaminsky AJ, Bluefield N, Miraliakbari R. Sample procurement for cultures of infected

- wounds: a systematic review. *J. of Wound Care*. 2016;25(4):4–10.
6. Bahramian A. Molecular characterization of the pilS2 gene and its association with the frequency of *Pseudomonas aeruginosa* plasmid pKlc102 and PaPI-1 pathogenicity island *Infect and drug resist.* 2019; 12:221.
 7. Talebi G. Survey of ermA, ermB, ermC and mecA genes among *Staphylococcus aureus* isolates isolated from patients admitted to hospitals in Tehran, Iran by PCR. *Biomed. Res.* 2019;30:259–263.
 8. Edwards-Jones V, Flanagan M, Wolcott R. Technological advancements in the fight against antimicrobial resistance. *Wounds*. 2015;6(2):47–51.
 9. Fabbro MD, Bortolin M, Taschieri S, Ceci C, Weinstein RL. Antimicrobial properties of platelet-rich preparations. A systematic review of the current pre-clinical evidence. *Platelets*. 2016;27:276–285.
 10. Cheesbrough M. *District laboratory practices in tropical countries*. Cambridge Press; 2010.
 11. Bowler PG, Duerden BJ, Armstrong DG. *Wound Microbiology and Associated Approaches to wound management*, Clin. Microbiology. 2001;14:244-269.
 12. Wilson APR, Gibbons C, Reeves BC, Hodgson B, Liu M, Plummer D. Surgical wound infections as a performance indicator: agreement of common definitions of wound infections in 4773 Patients. *British Mol. J.* 2004;329:720-722.
 13. Mohantay S, Kapil A, Dhawan B, Das BK. Bacteriological and antimicrobial susceptibility profile of soft tissue infections from Northern India. *Indian Journal of Model Science*. 2004;58:10-15.
 14. Mahat P, Manandhar S, Baidya R. Bacteriological Profile of Wound Infection and Antibiotic Susceptibility Pattern of the Isolates. *J. Microb and Experim.* 2017;4 (5):12-18
 15. Aizza Z, Naeem A, Hasan E. Bacteriology of Infected Wounds - A Study Conducted at Children Hospital Lahore. *Biomed. J.* 2007;23:3-9.
 16. Bessa LJ, Fazii P, Di Giulio M, Cellini L. Bacterial isolates from infected wounds and their antibiotic susceptibility pattern: some remarks about wound infection. *Intl Wound J.* 2015;12:47–52.
 17. Basu S, Ramchuran PT, Bali ST, Gulat I A, Shukla V. A prospective, descriptive study to identify the microbiological profile of chronic wounds in outpatients. *Ostomy Wound Manage.* 2009;55(1):14–20.

© 2022 Christian et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/85290>