



Pregnancy and its Outcomes in a Tertiary Care Hospital of Northern India: A Retrospective Study

Preeti Chowdhry^{a++}, Neeti Shree^{a#} and Sugandha Bansal^{a†*}

^a *SSB Heart and Multispecialty Hospital, Faridabad, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95763>

Original Research Article

Received: 25/10/2022

Accepted: 29/12/2022

Published: 29/12/2022

ABSTRACT

Indian women's maternal health status was found to be worse than that of women in other developed nations. One of the most crucial aspects of managing and caring for mothers and children has been the promotion of maternal and child health. Safe motherhood by providing good antenatal care (ANC) is very important to reduce maternal mortality ratio and infant mortality rate and achieve millennium development goals. Therefore, the present study focused on the effect of ANC services and the current status of pregnancies in a tertiary care hospital. The present retrospective study was conducted on pregnant women of OPD and IPD in the Obstetrics and Gynecology Department of a tertiary care hospital in North India between April to October 2022. According to the current study, the number of complicated deliveries was less as compared to the number of delivered women with positive neonatal outcomes. These findings can be used to plan a health intervention programme aiming to improve maternal and health practices which can improve the status of health in pregnant women. This also highlights the importance of proper ANC care and institutional deliveries.

⁺⁺ *Senior consultant;*

[#] *Associate Consultant;*

[†] *Consultant;*

^{*} *Corresponding author: E-mail: trialclini@gmail.com;*

Keywords: Antenatal care; pregnant women; newborns; Tertiary Care Hospital.

1. INTRODUCTION

The World Health Organization's Safe Motherhood Initiatives, a global initiative that was started in 1987, sought to lower the number of deaths related to pregnancy and childbirth [1,2]. The primary healthcare delivery system of a nation that strives for a healthy society is thought to be anchored by antenatal care (ANC) services. Improving maternal health is one of the eight-millennium development goals (MDGs). According to the MDGs, nations agreed to cut maternal mortality in half between 1990 and 2015. The number of maternal deaths worldwide have decreased by 47% since 1990 [3]. In India, despite numerous changes in a rapidly changing socioeconomic environment over the past 60 years, the maternal health situation has been appalling. For a long time, India had a very high maternal mortality ratio (MMR). According to a report, the national average MMR for Indian women is 212 per 100,000 live births (SRS - 2007-2009), which is extremely high when compared to international standards like Sweden's (5), the USA's (24), Brazil's (58), and even neighbouring nations like Sri Lanka's (39) and Thailand's (48) [1]. In addition to maternal mortality, 50 million women experience acute pregnancy-related complications that could be avoided by encouraging women to give birth in a medical facility or with the help of trained birth attendants [4]. In this context, primary care physicians' duties and responsibilities have also undergone ongoing revision. To address the issue, various health worker cadres have been appointed under their direction [5]. In order to ensure the best possible outcomes for the mother and her unborn child, ANC offers care to pregnant women that monitors the course of the pregnancy and promotes birth readiness and complication readiness [6]. According to the National Family Health Survey (NFHS)-4, only 21% of pregnant women in India used full ANC services, with state-by-state usage ranging from 2.3 to 65.9%. Numerous pregnancy-related deaths can be avoided with proper medical care during pregnancy and childbirth [7,8]. Therefore, the present study focused on ANC services and the current status of pregnancy in a tertiary care hospital in a tier 2 city in India.

2. METHODOLOGY

The present retrospective study was conducted in a tertiary care hospital on pregnant women of both OPD and IPD patients in the Obstetrics and

Gynaecology Department of SSB Heart and Multispecialty Hospital, Faridabad, Haryana, India, between April to October 2022 and followed up till its neonatal outcomes. The study population comprises of all pregnant women in their 3rd trimester attending the antenatal clinic.

3. RESULTS

A total of 111 deliveries were conducted during the study period, and the study revealed that the maximum number of deliveries were recorded in the month of October i.e. 23, followed by 19, 16, 17, 13, 12, 11 in the month of September'22, May'22, July'22, April'22, June'22 and August'22. Fig. 1 shows that out of the total 111 patients who delivered, 95(85.58%) were discharged within 48 hours of delivery. 100% patients were discharged within 48 hours of delivery in the month of June'22. 26 high risk and complicated deliveries were conducted in the hospital during this time with the highest number of patients with obstetric complications like Antepartum Haemorrhage (APH), Postpartum Haemorrhage (PPH), Sepsis and Eclampsia were seen in the month of September'22 i.e. 8 patients, out of which 5 patients received blood transfusions due to the complications within the month of September'22 (Fig. 2).

The highest numbers of deliveries (23) were recorded in the month of October which include 11 vaginal deliveries and 12 C-section deliveries, out of which 5 caesarean deliveries were carried out in the night shift (8 pm to 8 am). Out of the total 70 caesarean sections conducted during the study period, only 20 were conducted in the night shift (8 pm to 8 am).

112 babies were born in this period with one being a twin birth and total pregnancy outcomes were a total of 50 female and 62 male live childbirths. Out of all the babies born in the hospital, 37(33%) babies were low birth weight i.e. less than 2.5kg and 45(40%) babies had a pre-term delivery i.e. born before 37 weeks of gestation (Figs. 4 and 5). A total of 5 childbirths with less than 2.5 kg were recorded in April, June, July and October month and 6 children in the month of September'22. Among the 112 babies born, only 24(21.4%) babies were breastfed within 1 hour of delivery (Fig. 4). A total of 37 Medical termination of pregnancies (MTP) were conducted in the hospital, out of which 35 were done within 12 weeks period of gestation.

Only 2 MTP's were done in patients with more than 12 weeks of pregnancy, 1 each in month of May'22 and August'22 (Fig. 6).

Apart from MTPs, contraception is a very important part of healthcare of reproductive age women in India. 148 pregnancy test kits were used in the IPD and OPD of the hospital in the

period of study. 92 patients who came to the hospital, opted for combined oral contraceptive pills followed by 5 women who got postpartum sterilization within 7 days of delivery or concurrently with C-section, 2 patients opted for interval IUCD insertion and only 1 woman underwent interval laparoscopic sterilization method (Fig. 7).




Fig. 1. Shows the delivery and discharge pattern of pregnant women from April to October'22




Fig. 2. Shows the number of patients with complications during delivery and those which needed transfusion from April to October'22




Fig. 3. Shows the status of delivery during between April –October-22




Fig. 4. Shows the status of newborn in the tertiary care centre between April –October-22




Fig. 5. Shows the pregnancy outcomes and details of newborns between April –October-22




Fig. 6. Shows the status of medical termination of pregnancy (MTP) between April –October’22




Fig. 7. Shows the family planning method between April to October-22

4. DISCUSSION

Various Maternal healthcare indicators were evaluated to see the services in a tertiary care hospital in a tier 2 city. Most patients are discharged within 48 hours of delivery, while some cases are required to prolong hospital stay which can be due to various obstetric and non-obstetric reasons. Being a tertiary care hospital in a tier 2 city gave us insights that out of all the deliveries conducted, 23% of patients were those with obstetric complications and there was a high caesarean rate which can also be attributed to various factors like a patient preference for the specific date of delivery due to certain customs and beliefs, the low pain threshold of patients for vaginal delivery, fear of litigation in doctors, a high number of complicated patients (obstetric and non-obstetric causes) being referred for delivery from nearby areas and villages [9]. The current study also showed that despite higher pre-term deliveries, less number of babies had low birth weight status. This might be because people are becoming more aware of factors affecting mother and child health. However, only 21% of the babies born could be breastfed within 1 hour of delivery as most patients coming to a private hospital are resting in the first 1 hour and are not willing to start breastfeeding despite encouragement and counselling. The study also

revealed the most common choice of contraception being Oral contraceptives.

This study has a few limitations, but not many. The study's conclusions can only be generalized to pregnant women undergoing delivery in a private institute. Due to various cultural practices, norms, and beliefs, the results of the study may vary depending on where it is conducted. The sample size was also small and thus larger-scale studies are required to analyse the patterns in different areas and institutions of India.

5. CONCLUSION

In the end, this study came to the conclusion that the results will be useful in the planning and implementation of new health initiatives that enhance maternal health procedures. Specific intervention programs need to be planned and conducted to improve their maternal health practices and eventually improve their health status.

CONSENT

It is not applicable.

ETHICAL APPROVAL

Ethical approval for conducting the study was obtained from the institutional ethics committee

of the hospital and data was collected from patients' records.

ACKNOWLEDGEMENT

We would like to thank and acknowledge the co-operative faculty and staff at the Department of Obstetrics and Gynecology and Department of Pediatrics at the SSB heart and multispecialty Hospital, a unit of SSB Central Hospital and Research Centre limited, Faridabad for their support throughout the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Patel BB et al. A study on knowledge and practices of antenatal care among pregnant women attending antenatal clinic at a Tertiary Care Hospital of Pune, Maharashtra. Medical Journal of Dr. DY Patil University. 2016;9(3):354.
2. Muhsen K, El-Hai RA, Amit-Aharon A, Nehama H, Gondia M, Davidovitch N, Goren S, Cohen D. Risk factors of underutilization of childhood immunizations in ultraorthodox Jewish communities in Israel despite high access to health care services. Vaccine. 2012;30(12):2109-2115.
3. Chiabi A, Nguéfack FD, Njapndounke F, Kobela M, Kenfack K, Nguéfack S, Mah E, Nguéfack-Tsague G, Angwafo F. Vaccination of infants aged 0 to 11 months at the Yaounde Gynaeco-obstetric and pediatric hospital in Cameroon: how complete and how timely?. BMC Pediatrics. 2017;17(1):1-7.
4. Gupta RK, Shora TN, Verma AK, Jan R. Knowledge regarding antenatal care services, its utilization, and delivery practices in mothers (aged 15-49 years) in a rural area of North India. Trop J Med Res. 2015;18(2):89-94.
5. Roy MP, Mohan U, Singh SK, Singh VK, Srivastava AK. Determinants of utilization of antenatal care services in rural Lucknow, India. Journal of Family Medicine and Primary Care. 2013;2(1):55.
6. Rustagi R, Basu S, Garg S, Singh MM, Mala YM. Utilization of antenatal care services and its sociodemographic correlates in urban and rural areas in Delhi, India. European Journal of Midwifery. 2021;5.
7. Kumar H, Chacko IV, Mane S, Govindan NN, Prasanth S. Study of utilization of antenatal care services and its determinants among pregnant women admitted in a tertiary care hospital in Mangaluru, Karnataka, India. Int J Community Med Public Health. 2020; 7:1960-7.
8. Garovic VD, Kattah A, Rose CH, Arendt KW. September. Medical and surgical illnesses during pregnancy: perspectives on immediate and long-term outcomes. In Mayo Clinic Proceedings Elsevier. 2016; 91(9):1151-1154.
9. Muula AS. Ethical and practical consideration of women choosing cesarean section deliveries without "medical indication" in developing countries. Croatian Medical Journal. 2007; 48(1):94.

© 2022 Chowdhry et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95763>