



Relative Abundance of Insect Pests on French Bean (*Phaseolus vulgaris* L.) in Relation to Abiotic Factors

Bijnesh Kumar ^{a++}, Sameer Kumar Singh ^{a#*},
Pankaj Kumar Yadav ^{a++} and Rajneesh Kumar Verma ^{a++}

^a Department of Entomology, College of Agriculture, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya - 224 229, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i214035

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107897>

Original Research Article

Received: 20/08/2023

Accepted: 25/10/2023

Published: 31/10/2023

ABSTRACT

The present investigation was carried out during *Rabi* 2021-22. The incidence of insect pests like aphid, whitefly, Jassid, leaf miner, pod borer was initiated 25 days after sowing till harvesting. The highest incidence of Jassid was observed on French bean crop at 9th SMW (7.20 per cage) and lowest 1st SMW (0.30 per cage), Aphid 8thSMW (45.20 aphids per 10cm terminal shoot) and 52thSMW (2.60 aphids per 10cm terminal shoot). The highest incidence of whitefly observed in 8th SMW (20.70 per cage) and lowest 2nd SMW (0.60 per cage). The highest incidence of Leaf miner was recorded in 11th SMW (40.50% leaf damage) and lowest in 51th SMW (1.50% leaf damage). The highest incidence of pod borer was noticed in 9th SMW (8.70 larvae per plant) and lowest in 1st SMW (0.30 larvae per plant). The Jassid population showed highly significant positive correlation with sunshine and positive significant correlation with maximum temperature and positive correlation with minimum temperature whereas, significant negative correlation with rainfall and

⁺⁺ PG Students;

[#] Assistant Professor;

^{*}Corresponding author: E-mail: drsameerent@nduat.org;

negative non-significant correlation with relative humidity. The aphid population showed highly significant positive correlation with sunshine and positive non-significant correlation with minimum and maximum temperature. The whitefly population showed highly significant positive correlation with sunshine and positive non-significant correlation with minimum and maximum temperature. The leaf miner population showed positive highly significant correlation with minimum and maximum temperature and significant positive correlation with relative humidity. The pod borer population showed highly significant correlation with maximum temperature and sunshine. The information generated in the present research can be successfully utilized to formulated effective insect pest management strategies in French bean.

Keywords: French bean; Jassid; aphid; whitefly; leaf miner; pod borer.

1. INTRODUCTION

The Leguminosae family includes the French bean (*Phaseolus vulgaris* L.), a significant and extremely lucrative vegetable and pulse crop in India. Mesoamerica was the origin place of the French bean [1]. Rajmah, common bean, kidney bean, haricot bean, salad bean, snap bean, and string bean are some other names for *P. vulgaris*. There are about 80 cultivated and wild species in the huge genus *Phaseolus*, but *P. vulgaris* is the one that is most frequently cultivated. French bean is widely grown in India due to the short duration and nutritional benefits. The crop has been introduced in northern plains during *Rabi* season also. Considering the nutritive value, 100g of green pod contains 1.70g protein, 0.10g fat, 4.5g carbohydrate, 1.80g fiber and is also rich in minerals and vitamins. It contains calcium (50 mg), phosphorus (28.00 mg), iron (1.70 mg), carotene (132.00 mg), thiamine (0.08 mg), riboflavin (0.06 mg) and vitamin C (24.0 mg) per 100 g of edible part. Additionally, it has some medical qualities that are helpful in managing diabetes and some heart issues, and it is a wonderful all-natural treatment for bladder burn. It is both carminative and reparative in its effects on diarrhea and constipation [2].

Jharkhand, Karnataka, Andhra Pradesh, Maharashtra, Tamil Nadu, Odisha, Bihar, Nagaland, Meghalaya, Sikkim, Mizoram, Jammu and Kashmir, Himachal Pradesh, and West Bengal are the principal states where French beans are grown. In Karnataka French bean is being grown in Chikkamagaluru, Hassan and Shivamogga. In hilly region, it is grown during *Kharif* and in lower hill region, sown as spring crop. In north-east plains and hilly tracts of Maharashtra, it is cultivated during *Rabi*. It is highly sensitive to frost and water logging. The average French bean production and productivity in India was 6.75 lakh tonnes and 27.94 t/ha,

respectively, on an area of 2.42 lakh ha. On a surface of 15.80 lakh ha, the global production and productivity were 242.21 tonnes and 153.30 t/ha, respectively [3].

Leguminous plant species are susceptible to a wide range of biotic stressors, pest insects, and diseases. The French bean's agricultural output is severely hampered by pest and disease issues, especially in the tropics [4]. In India, disease and insect infestation are the main causes of low productivity in the French bean crop. The French bean crop has been affected by reports of over 37 different insect pest species [5]. Among them several serious insect pests attack the French bean of which aphids (*Aphis craccivora*), leafhopper (*Amrasca biguttula biguttula*), whitefly (*Bemisia tabaci*), and thrips (*Caliothrips indicus*) [6,7] and foliage feeders viz., serpentine leaf miner (*Liriomyza trifolii*) and bean stem fly (*Ophiomyia* spp.) are the major insect pests. The spotted pod borer (*Maruca vitrata*) and red spider mite (*Tetranychus* spp.) are other pests that attack French beans [8]. In addition to causing direct damage by feeding the sap suckers are the vectors of viral diseases in French bean. Whitefly act as a vector for more than 100 plant viruses, which cause disease to many commercial crops in different parts of the world [9]. Which adversely influence the quality and quantity of yield [10,11]. Considering above facts, the present study was conducted to find out Relative abundance of Insect Pests on French bean in relation to abiotic factors.

2. MATERIALS AND METHODS

The Monitoring Seasonal incidence of Insect pest of French bean was conducted at Student's Instructional Farm (SIF), Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya, Uttar Pradesh, India during *Rabi*, 2021-22 on French bean Variety *Chitra*

with a plot size of 6m x 6m and spacing of 30cm x 10cm.

The population build-up of insect pest was recorded from 10 randomly selected plants and the numbers of Insect pests present was the counted at weekly intervals on a Standard Meteorological weekly basis from the appearance of insects to till the harvesting. "Direct visual counting method" was adopted for the estimation of the population of whitefly and jassids by using rectangular split cage. The incidence of leaf miner was recorded by counting total number of leaves and damaged leaves by leaf miner Total number of larvae present on per plant was counted to estimate the population of pod borer. Total of 10 plants were randomly selected to observe the incidence insect pests. The incidence of aphid was recorded from top 10cm terminal shoot. The observations were noted at weekly intervals during morning hours between 6.00 AM to 7.30 AM.

3. RESULTS AND DISCUSSION

3.1 Relative Abundance of Insect Pests in French Bean

3.1.1 Jassid (*A. biguttula biguttula*)

The incidence of Jassid was observed on 51th SMW (0.80 Jassid per cage). The peak population was observed at 9th SMW (7.20 Jassid per cage). Whereas, the relative humidity (morning) was also maximum (79.60%) and no rainfall, and sunshine (9.0 hrs). Though, the minimum population was recorded at 1st SMW (0.30 jassid per cage), during *Rabi* season, 2021-22. The present findings are in accordance to Sahoo and Giraddi [12] who found that the peak population leafhopper was observed from vegetative to flowering stage. The present findings are similar to Singh and Yadav [13] who found the mean population of jassids 0.4-5.80/cage and 0.20-7.40/cage, respectively. Incidence of the jassids and whiteflies started in 50th standard week with 0.40/cage and 0.20/cage, respectively.

3.1.2 Aphid (*Aphis craccivora*)

The mean population of aphid (*A. craccivora*) was observed at first time on 51th SMW during *Rabi*, 2021-22. The mean population of *A. craccivora* showed its peak (45.20 per 10 cm terminal shoot) during 8th SMW. The results are in conformity with Kishor et al. [14] who found that the incidence of aphid (*Aphis craccivora*)

started from 53rd Meteorological Standard Week (MSW) (23.80 aphid/10 cm apical twigs). The aphid population gradually increased and reached its peak (35.40 aphids/10cm apical twigs) on 7th MSW (22nd February) and thereafter its population gradually decreased from 8th MSW on 4th week of February (31.30 aphids/10 cm apical twigs) to 12th MSW (4th week of March) (9.0 aphids/10cm apical twigs).

3.1.3 Whitefly (*Bemisia tabaci*)

The Population of whitefly (*B. tabaci*) was recorded at first time on 51th SMW during *Rabi*, 2021-22. The highest incidence (20.70 per cage) of whitefly was recorded at 8th Standard Meteorological Week. These findings are similar to Rani et al. [15] who found that the incidence of whitefly, *Bemisia tabaci* (Gennadius) population was noticed from the second week of November to second week of December during first season with peak incidence during the third week of November and a mean population of 0.12 per leaf. Whereas during second season, its population was noticed from second week of February to second week of March with peak incidence during the last week of February and a mean population of 0.28 per leaf.

3.1.4 Leaf miner (*Liriomyza trifolii*)

The mean population of serpentine leaf miner (*L. trifolii*) was observed at first time at 45th SMW. The highest attack (40.50% leaf miner damage) of leaf miner damage was observed at Standard Meteorological Week number 11th during *Rabi* season 2021-22. The present findings are in accordance to Sahoo and Giraddi [12] who found that the peak of leaf miner was observed from vegetative to maturity stage. The present findings are in conformity with Khaliq et al. [16] who found that first appearance being during the 49th standard meteorological week (SMW) 0.95 miners/leaf which reached its peak (11.00 miners/leaf) during the 4th SMW reached to minimum (1.26 miners/leaf) during the 12th SMW. These findings are similar to Mahesh et al. [17] reported that the population of leaf miner appeared on the crop from seedling stage till pod maturation (15-70 days after sowing) and the maximum number of insect pests infested the crop during vegetative and flowering stage.

3.1.5 Pod borer (*Helicoverpa armigera*)

The Incidence of pod borer (*H. armigera*) was recorded at first time on 1st SMW during *Rabi* season 2021-22. The highest population (8.70

larvae per plant) of pod borer was recorded at 9th Standard Meteorological Week. The results are similar to Jakkaray et al. [18] who found that the pod borer incidence started from the first week of January (1stMSW) and continued till January last week (5thMSW). The lowest and highest population of bean pod borer was 0.35 larvae per plant noticed in January first week and 1.89 larvae per plant in the third week (3rdMSW), respectively.

3.2 Effect of Abiotic Factors on Relative Abundance of Insect Pests in French Bean

The correlation between incidence of major insect pest and abiotic factors viz., maximum temperature (°C), minimum temperature (°C), relative humidity (%) and rainfall (mm), and sunshine (hrs.) have been described under this section.

3.2.1 Jassid (*A. biguttula biguttula*)

The incidence of Jassid had positive and positive significant correlation with minimum & maximum temperature, respectively ($r=0.489$ and $r=0.645^*$) whereas, non-significant negative correlation was observed with relative humidity ($r=-0.488$). The rainfall showed significant negative correlation ($r=-0.645^*$), whereas sunshine had highly significant positive correlation ($r=0.856^{**}$) during Rabi season 2021-22. The present findings are similar with Kumar [19] found that that jassid population had positive and highly significant correlation with maximum temperature while minimum temperature. The present study is in conjunction with the findings of Anandmurthy et al. [20] revealed that maximum temperature showed a significant positive correlation with jassid population in summer cowpea.

3.2.2 Aphids (*A. craccivora*)

The incidence of aphids had positive non-significant correlation with temperature (minimum and maximum) ($=0.363$ and $=0.437$, respectively) whereas, non-significant negative correlation with relative humidity ($=-0.328$), rainfall had highly significant negative correlation ($=-0.697^{**}$), while sunshine had highly significant positive correlation ($=0.765^{**}$) during Rabi season 2021-22. The present findings are similar with Kataria and Kumar [11] the population was showing positive with high temperature. The present findings are also similar with Chandra et al. [21] who found that aphids were non-significant positive correlation with maximum temperature

and significant negative with rainfall during both the years in French bean crop.

3.2.3 Whitefly (*B. tabaci*)

The observed incidence of whitefly had non-significant positive correlation with temperature (minimum and maximum) ($=0.430$ and $=0.539$, respectively) whereas, non-significant negative correlation with relative humidity ($=-0.506$), significant negative correlation ($=-0.624^*$) with rainfall, highly significant positive correlation ($=0.811^{**}$) with sunshine during Rabi season 2021-22. The Similar work had been reported by Ghosh [22] who reported that maximum temperature and minimum relative humidity showed non-significant positive correlation with whitefly population. The present findings are similar with Singh et al. [23] who revealed that the whitefly, *B. tabaci* population with maximum temperature significant positive correlation ($r = 0.764$) but minimum temperature, relative humidity and rainfall revealed negative correlation.

3.2.4 Leaf miner (*L. trifolii*)

The incidence of leaf miner showed positive highly significant correlation with temperature (minimum and maximum) ($=0.802^{**}$ and $=0.880^{**}$), respectively whereas, significant positive correlation with relative humidity ($=0.582^*$), significant negative correlation ($=-0.597^*$) with rainfall, highly significant positive correlation ($=0.875^{**}$) with sunshine during Rabi season 2021-22. The present findings are similar with Mazumdar et al. [24] found that in French bean and Cowpea there was significantly positive relation of temperature, whereas wind speed showed negative relation to Agromyzid infestation on cultivated crops.

3.2.5 Pod borer (*H. armigera*)

The pod borer incidence had positive non-significant and highly significant correlation with temperature (minimum and maximum) ($=0.570$ and $=0.691^{**}$, respectively) whereas, non-significant negative correlation with relative humidity (-0.489), highly significant negative correlation ($=-0.628^{**}$) with rainfall, highly significant positive correlation ($=0.847^{**}$) with sunshine during Rabi season 2021-22. The present findings are partially similar of the Tamang et al. [25] who found that Pod borer population exhibited highly significant positive correlation with minimum temperature.

Table 1. Incidence of pest spectra in French bean under the influence of weather factors during *Rabi* 2021-22

SMW	Date	Temperature(°C)		Rainfall (mm)	Relative humidity (%)	Sun shine (hrs.)	Jassids /Cage	Aphids/10 cm terminal shoot	White fly/ Cage	Leaf miner damage (%)	Larval pop. of pod borer/ plant
		Min.	Max.								
51	17-23 Dec 2021	5.90	21.60	0.00	77.30	5.00	0.80	5.80	2.60	1.50	0.00
52	24-31 Dec 2021	9.40	20.90	15.00	81.50	2.60	0.50	2.60	1.50	2.70	0.00
1	1-7 Jan 2022	9.30	19.40	14.20	82.50	2.60	0.30	3.50	0.90	5.20	0.30
2	08-14 Jan 2022	10.60	20.20	11.60	87.50	2.20	0.40	4.20	0.60	4.70	0.70
3	15-21 Jan 2022	5.70	15.90	0.00	86.75	2.00	1.20	15.90	4.30	8.20	1.80
4	22-28 Jan 2022	8.30	17.30	8.80	87.45	2.90	0.90	9.50	1.20	10.70	0.90
5	29Jan-04Feb 2022	8.90	19.60	3.00	86.70	4.90	1.00	17.50	2.60	12.30	1.20
6	05-11 Feb 2022	8.20	20.60	0.00	86.35	4.90	3.80	28.00	8.50	14.50	3.50
7	12-18 Feb 2022	9.90	24.00	0.00	80.40	8.30	4.20	39.50	16.20	21.50	4.30
8	19-25 Feb 2022	13.50	25.60	0.00	77.70	8.10	5.50	45.20	20.70	28.70	6.50
9	26 Feb -5 Mar 2022	11.40	27.10	0.00	79.60	9.00	7.20	30.50	15.60	35.30	8.70
10	05-11Mar 2022	12.50	29.10	0.00	77.30	7.80	4.40	25.20	10.70	38.20	6.30
11	12-18 Mar 2022	16.20	32.00	0.00	77.10	8.30	2.70	15.30	5.20	40.50	3.70

SMW= Standard Meteorological Weeks

Table 2. Effect of abiotic factors on incidence of pest spectra in French bean during *Rabi* 2021-22

Insect Pests	Weather Parameters				
	Temperature(°C)		Rainfall (mm)	Relative Humidity (%)	Sunshine (hrs.)
	Min.	Max.			
Jassid	0.489	0.645*	-0.645*	-0.488	0.856**
Aphid	0.363	0.437	-0.697**	-0.328	0.765**
Whitefly	0.430	0.539	-0.624*	-0.506	0.811**
Leaf minor	0.802**	0.880**	-0.597*	0.582*	0.875**
Pod borer	0.570	0.691**	-0.628**	-0.489	0.847**

*Significant at the 0.05 level, **Significant at the 0.01 level




Fig. 1. Incidence of pest spectra in French bean under the influence of weather factors during Rabi 2021-22

4. CONCLUSION

The incidence of insect pests like aphid, whitefly, Jassid, leaf miner, pod borer was initiated after sowing till harvesting. The highest incidence of Jassid was observed on French bean crop at 9th SMW (7.20 per cage) and lowest 1st SMW (0.30 per cage), Aphid 8thSMW (45.20 aphids per 10cm terminal shoot) and 52thSMW (2.60 aphids per 10cm terminal shoot). The highest incidence of whitefly observed in 8th SMW (20.70 per cage) and lowest 2nd SMW (0.60 per cage). The highest incidence of Leaf miner was recorded in 11th SMW (40.50% leaf damage) and lowest in 51th SMW (1.50% leaf damage). The highest incidence of pod borer was noticed in 9th SMW (8.70 larvae per plant) and lowest in 1st SMW (0.30 larvae per plant). The Jassid population showed highly significant positive correlation with sunshine and positive significant correlation with maximum temperature and positive non-significant correlation with minimum temperature whereas, significant negative correlation with rainfall and negative non-significant correlation with relative humidity. The aphid population showed highly significant positive correlation with sunshine and positive non-significant correlation with minimum and maximum temperature. The whitefly population showed highly significant positive correlation with sunshine and positive non-significant correlation with minimum and maximum temperature. The leaf miner population showed positive highly significant correlation with minimum and maximum temperature and significant positive correlation with relative humidity. The pod borer population showed highly significant correlation with maximum temperature and sunshine. The information generated in the present research can be successfully utilized to formulated effective insect pest management strategies in French bean.

ACKNOWLEDGEMENTS

The authors are highly thankful to the Head, Department of Entomology, College of Agriculture, Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya (U.P.) India for providing essential facilities and support during the experimentation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bitocchi E, Nanni L, Bellucci L, Rossi M, Giardini A, Zeuli PS, Logozzo G, Stougaard J, Mcclean P, Attene G and Papa R. Mesoamerican origin of the common bean (*Phaseolus vulgaris* L.) is revealed by sequence data. Proc. Nat. Acad. Sci. 2012;109(14):788-796.
2. Duke JA. Handbook of world economic importance. USDA, Beltsville, Maryland, Plenum, New York and London; 1981.
3. FAOSTAT. Available:<http://www.fao.org/faostat/en/#data/QC>.2018
4. Graham PH, Vance CP Legumes: Importance and constraints to greater use. Plant Physiol. 2003;131(3):872-877.
5. Mondal A, Shankar U, Abrol DP, Kumar A, Singh AK. Incidence of major insect pests on French bean and relation to environmental variables. Indian J. Entomol. 2018;80(1):51-55.
6. Sahu A, Singh PS, Singh SK. Assessment of host plant resistance in certain French bean (*Phaseolus vulgaris* L.) genotypes against major insect pests. J. of Exp. Zool. India. 2021;24(2):1167-1170.
7. Anonymous. Consensus document on the biology of common bean (*Phaseolus vulgaris* L.) Series on Harmonisation of Regulatory Oversight in Biotechnology No. 59. OECD Env. Health and Safety Div. 2015;47:24.
8. Nderitu JH, Wambua EM, Olubayo F, Kasina JM and Waturu CN. Evaluation of French beans (*Phaseolus vulgaris* Cultivars and Breeding lines for Resistance to Thrips (*Thysanoptera*: Thripidae) pest in Kenya. J. Entomol. 2007;4:202-209.
9. Jones DR. Plant viruses transmitted by whiteflies. Eur. J. Plant Pathol. 2003;109(3):195-219.
10. Oyewale RO, Bamaiyi LJ. Management of cowpea insect pests. Scholars Academic Journal of Biosciences. Scholars Academic Journal of Biosciences (SAJB). Sch. Acad. J. Biosci. 2013;1(5):217-226.
11. Kataria R, Kumar D. Population dynamics of *Aphis craccivora* (Koch) and its natural enemies on bean crop in relation to weather parameters in Vadodara, Gujarat, India. Legume Res. 2016;9(1):36-41.
12. Sahoo S, Giraddi RS. Insect pest spectrum of french bean *Phaseolus vulgaris* (pole type) grown under net

- house. Indian Journal of Entomology Online published. 2021;e21177.
13. Singh PS, Yadav NK. Seasonal abundance of insect pests on mung bean and its correlation with abiotic factors. J. Entomol. Res. 2013;37(4):297-299.
 14. Kishor DR, Prasad R, Moses S, Singh PP. Population dynamics of aphid and pod borer on lentil and their natural enemies during Rabi season 2017 at Pusa, Samastipur. Curr. J. App. Sci. Tech. 2019;32(2):1-6.
 15. Rani JB, Hanumantharaya L. Population dynamics of insect pests of French bean under hill zone of Karnataka. Adv. Life Sci. 2016;5(5):1951-1956.
 16. Khaliq N, Shankar U, Rather BA, Kumar Y. Seasonal incidence of pea leaf miner *Chromatomyia horticola* Goureau Infesting Garden Pea. Indian. J. Entomol. 2022;1-3.
 17. Mahesh JN, Dharmanna DK, Jayappa AH. Succession of major insect pests of french bean (*Phaseolus vulgaris* Linn.) in Relation to Crop Stage. Adv. Life. Sci. 2016;5(5):1906-1909.
 18. Jakkaray BC, Sharanabasappa H, Adivappar N. Seasonal incidence of lepidopteran pests in French bean (*Phaseolus vulgaris* L.). Int. J. Ecol. Envi. Sci. 2020;2(4):312-315.
 19. Kumar S. Management of insect pest complex of tomato with combination insecticides. Thesis, M.Sc. A.N.D. University of Agriculture and Technology, Ayodhya. 2021;25-38.
 20. Anandmurthy T, Parmar GM, Arvindarajan G. Seasonal incidence of major sucking pests infesting cowpea and their relation to weather parameters. Int. J. Plant Prot. 2018;11(1):35-38.
 21. Chandra U, Singh K, Singh HM, Kumar R. Population dynamics of sap feeders in urdbean [*Vigna mungo* (L.) Hepper] and their correlation with abiotic factors. Int. J. Plant Prot. 2010;3(2):274-275.
 22. Ghosh SK. Incidence of whitefly (*Bemisia tabaci* Genn.) and their sustainable management by using biopesticides. Proceedings of the 4th ISOFAR scientific conference. 'Building Organic Bridges', at the Organic World Congress, Istanbul, Turkey. 2014;623-626.
 23. Singh M, Bairwa DK, Jat BL. Seasonal incidence of sucking insect pests of green gram. J. Entomol. Zool. Stud. 2019; 7(2):654-658.
 24. Mazumdar S, Bhuiya BA. Correlation between climatic factors and leaf miner (*Insecta: Agromyzidae*) infestation on three vegetable crops in Chittagong, Bangladesh. J. Asiat. Soc. Bangladesh Sci. 2015;41(1):1-5.
 25. Tamang S, Venkatarao P, Chatterjee M, Chakraborty G. Population dynamics of major insects pest of mung bean and correlation with abiotic under terai agroclimatic zone of West Bengal. 2017;12(2):893-897.

© 2023 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107897>