



The Characteristic, Occurrence of Aflatoxin and Associated Risk with Human Health

Sharique Ahmad ^a[¶], Subuhi Anwar ^a[#] and Pushendra D. Pratap ^b[†]

^a Department of Pathology, Era's Lucknow Medical College and Hospital, Era University, Lucknow, Uttar Pradesh – 226003, India.

^b Department Biochemistry, Era's Lucknow Medical College and Hospital, Era University, Lucknow, Uttar Pradesh - 226003, India .

Authors' contribution

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/MRJI/2022/v32i71333

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92468>

Review Article

Received 08 August 2022
Accepted 14 October 2022
Published 20 October 2022

ABSTRACT

Aflatoxins are toxins produced by the fungus *Aspergillus parasiticus* and *Aspergillus flavus*. They are colorless, cannot be detected under normal light, odorless and contaminated foods most often do not have any special or bad smell. Environmental factors such as the temperature, and vectors causing grain damage have an important effect on favoring the colonization of fungus and the production of aflatoxins. They contaminate agriculture products worldwide affecting their quality, safety & security. The effects on safety & security cause health threats among people and animals on their long-term exposure through consumption because they are mutagenic, teratogenic, genotoxic, and carcinogenic. Humans and animals can develop the disease known as aflatoxicosis as a result of aflatoxins known as Acute and Chronic primary aflatoxicosis. Aflatoxins' long-term effects have been shown to decrease normal immune response also it can cause growth retardation in babies and affect nutritional deficiency. It can disrupt such enzymes that can affect hormones, endocrine glands, and neurotransmitters which may influence a person's cognitive abilities, memory, and learning, restlessness, muscular tremors, seizures, absentmindedness, tremors, uncoordinated movement of muscles, and aberrant agitation are caused by deficiencies in the neurotransmitters.

[¶]Professor;

[#]Research Assistant;

[†]Research Scholar;

*Corresponding author: E-mail: Diagnopath@gmail.com;

Aflatoxin toxicity in humans can range from acute or chronic conditions to liver damage such as liver carcinoma, internal bleeding, edema, and instant death. Acute toxicosis can cause gastrointestinal dysfunctions in humans and animals. Aflatoxins have been shown to have detrimental negative impacts on vascular fragility and cardiovascular health, tissue bleeding, and teratogenic consequences. According to reports, aflatoxins have severe and detrimental impacts mostly on the respiratory systems it can also cause lung cancer.

Keywords: Aflatoxicosis; food security; safety; fungi; one health.

1. INTRODUCTION

Aflatoxins are toxins biosynthesized by fungi *Aspergillus parasiticus* and *Aspergillus flavus*. Which are abundant in nature hence lying in the category of mycotoxins. The name aflatoxins were derived from a combination of Afla and toxins where 'A' in afa stands for *Aspergillus* & 'fla' stand for *flavus* [1]. Chemically, aflatoxins are derivatives of difurano coumarins possessing a pentanone ring joined to the coumarin nucleus by a bifuran group. [2]. Aflatoxins were first discovered in England during the 1960s when an estimated 100,000 turkeys died after they had been fed with feeds contaminated with the fungi *Aspergillus flavus*. Environmental factors have a significant impact on aflatoxin contamination. [3]. An increase in temperature by every 2°C in European nations proved to raise the risk of aflatoxin production in cereals and the probability of contamination of cereals such as maize is likely to keep increasing for the next thirty years due to ideal meteorological circumstances [4,5]. Fungi colonization on agriculture products the initial point through

which aflatoxins enter the chain are favored by their nature (grain damage), moisture, and the environmental temperatures Fig. 1. Fungi grow in areas with a temperature range of 12-48°C producing aflatoxins at a 25-37 °C temperature range, together with a moisture content of 80-85 percent [6,7].

During this temperature range, the agriculture produces are experiencing a high rate of respiration. One of the outcomes water vapor will lead to an increase in the moisture content providing a favorable environment for fungi colonization and synthesis of aflatoxins when it increases beyond 14%. The temperature range also favors the expression of genes by the fungi for the production of aflatoxins.

In countries, where agricultural production is very high, the environmental temperatures lay between 40°N and 40°S of the equator the range with which fungus development and aflatoxin formation are favored hence exposing the environment and its components to aflatoxins effects.




Fig. 1. Cycle for synthesis of aflatoxins

2. TYPES OF AFLATOXINS

Aflatoxins belong to a difuranocoumarins group and are categorized into difurocoumaro cyclopentenone and difurocoumarolactone groups as per their chemical structure summarized in Table 1. Aflatoxins are of different types however, only four are major one's aflatoxins AFB₁, AFB₂, AFG₁ & AFG₂ Fig. 2. Aflatoxins B1 & B2 are hydroxylated when consumed by animals to M1 {AFM₁} & M2 {AFM₂} respectively [8]. AFB₁ and AFB₂ are

found in crops and or their products whereas AFM₁ and AFM₂ are found in animal by-products. AFB₁ & AFB₂ are synthesized by the fungi *Aspergillus flavus* whereas AFG₁ & AFG₂ are synthesized by the fungi *Aspergillus parasiticus*.

B & G stand for the blue fluorescent and green bright color created on the chromatographic plates when under UV light whereas the numbers 1 & 2 indicate major and minor compounds. B is produced by the flavus species and G by the parasitic species.

Table 1. List of the major aflatoxins produced by *Aspergillus* species

Sl. No.	Name of Toxin	Group	Species
1	Aflatoxin B1 (AFB1)	Difurocoumarocyclopentenone series	<i>A. flavus</i> <i>A. arachidicola</i> <i>A. bombycis</i> <i>A. minisclerotigenes</i> <i>A. nomius</i> <i>A. ochraceoroseus</i> <i>A. parasiticus</i> <i>A. pseudotamarii</i> <i>A. rambellii</i> <i>Emericella venezuelensis</i>
2	Aflatoxin B2 (AFB2)		<i>A. arachidicola</i> <i>A. flavus</i> <i>A. minisclerotigenes</i> <i>A. nomius</i> <i>A. parasiticus</i>
3	Aflatoxin B2a (AFB2a)		<i>A. flavus</i>
4	Aflatoxin M1 (AFM1)		<i>A. flavus</i> <i>A. parasiticus</i> Metabolite of aflatoxin B1 in humans and animals and comes from a mother's milk
5	Aflatoxin M2 (AFM2)		Metabolite of aflatoxin B2 in milk of cattle fed on contaminated foods
6	Aflatoxin M2A (AFM2A)		Metabolite of AFM2
7	Aflatoxicol (AFL)		<i>A. flavus</i> Metabolite of AFB1
8	Aflatoxicol M1		Metabolite of AFM1
1	Aflatoxin G1 (AFG1)	Difurocoumarolactone series	<i>A. arachidicola</i> <i>A. flavus</i> <i>A. minisclerotigenes</i> <i>A. nomius</i> <i>A. parasiticus</i>
2	Aflatoxin G2 (AFG2)		<i>A. arachidicola</i> <i>A. flavus</i> <i>A. minisclerotigenes</i> <i>A. nomius</i> <i>A. parasiticus</i>
3	Aflatoxin G2A (AFG2A)		Metabolite of AFG2
4	Aflatoxin GM1 (AFG1)		<i>A. flavus</i>

Sl. No.	Name of Toxin	Group	Species
5	Aflatoxin GM2 (AFGM2)		Metabolite of AFG2
6	AFGM2A		Metabolite of AFGM2
7	Aflatoxin B3 (AFB3)		Transformation of AFG1
8	Parasiticol (P)		<i>A. flavus</i>
9	Aflatrem		<i>A. flavus</i> <i>A. minisclerotigenes</i>
10	Aspertoxin		<i>A. flavus</i>
11	Aflatoxin Q1 (AFQ1)		Major metabolite of AFB1 <i>in vitro</i> liver reparations of other higher vertebrates




Fig. 2. Structures for the types of aflatoxins

3. CHARACTERISTICS OF AFLATOXINS

Aflatoxins are colorless and cannot be detected under normal light and can be made visible under UV light (basic detection method), they are odorless and contaminated food most often does not have any special or bad smell. Sometimes grains can smell moldy due to fungal contamination when the moisture content of the contaminated product is still high.

They are insoluble in water, solubility in solvent polarities like chloroform and methane, and they are unstable in extreme pH (<3 or >10).

Aspergillus flavus fungi are green and can multiply their population under stressful conditions. Penetration and colonization of fungi in crops are also supplemented by the injuries caused by vectors such as insects and nematodes [9].

They decompose at temperatures of 237-306⁰C and hence do not decompose quickly [10]. Therefore, limiting the consequences of aflatoxin requires either stopping the development of the fungi or detoxifying contaminated feeds and meals [11].

4. THE PREVALENCE OF AFLATOXIN CONTAMINATION IN AGRICULTURAL PRODUCTS

Contamination of food and feeds with aflatoxins worldwide presents major food and feed safety issues in areas of the world where humidity is high and temperatures are warm which encourages fungi growth Fig. 3. Fungi colonization and contamination usually happen during the pre-harvesting, harvesting, and post-harvesting sessions leading to the synthesis of aflatoxins.




Fig. 3. Aflatoxins flow along the feeding cycle leading to health

According to FAO, more than a 25percent of total the global agriculture commodities are contaminated by aflatoxins resulting in health issues when consumed. Aflatoxin contamination also results in a loss of market value [12]. Mycotoxin contamination in Sub-Saharan Africa among agriculture products is a great threat to health where more than 250,000 deaths occurring annually are hepatocellular carcinoma-related resulting from aflatoxin contamination alone [13].

In a country like Uganda, more than 60% of the population carries out farming in which mycotoxins remain a scourge and a previously unheard-of continue to decrease in the economic and dietary benefit of agriculture products [14]. The economic growth of many developing countries like Uganda greatly depends on the trading of agriculture products and due to the poorly equipped pre-harvesting, during harvesting and post-harvesting handling by farmers and the favoring environmental conditions leads to an increase in aflatoxins contamination hence high levels beyond the acceptable quantities [15].

According to WHO, in Africa, the aflatoxins burden accounts for about 40% of daily diseases. In 1974, there was an outbreak of Hepatitis in India due to aflatoxins which were found in their staple food maize. In Kenya, there are reported cases of disease outbreaks due to the consumption of maize a staple food that also is a

source of animal feeds contaminated with aflatoxins. In 2013, countries in Europe reported contamination of milk with aflatoxin, which was acquired by the animals through the feeds. Many countries cross-check the level of aflatoxins levels in imported products to avoid ill effects on their population.

5. RISK ASSOCIATED WITH AFLATOXINS

Aflatoxins are toxins (poisonous substances) produced within the cells of organisms. They are thus among the most toxic chemicals that have an impact on how plants, animals, and people are physiologic. This is referred to as aflatoxicosis hence a One Health concern.

Recognizing the connections between people, animals, plants, and their common environment, One Health is a coordinated, multisector, and transdisciplinary strategy working to achieve optimal health outcomes. Aflatoxins affect food safety & security one of the issues One Health program focuses on rectifying.

This causes health threats among people and animals on their long-term exposure through consumption because of the nature of its carcinogenic, mutagenic, teratogenic, and genotoxic effects demonstrated in Fig. 4 [16,17]. All vertebrate species are susceptible to aflatoxicosis with animals being more susceptible at a lower aflatoxin concentration as compared to People [6]. The aflatoxins in the occurrence of




Fig. 4. Overview of AFB1 metabolism and excretion in humans and animals

feeds & foods of animals result in a decline in productivity (of meat, eggs & milk), reduction in weight gain, and an increase in disease incidences due to their immunosuppression abilities and the damage to vital organs [18].

On consumption of agriculture products contaminated with aflatoxins by both animals and humans, health issues arise. When animals feeding on agricultural produce are eaten or their products are eaten by humans, these aflatoxins will be expressed in different tissues ending up in human bodies thus being hazardous.

Aflatoxins in humans affect the liver, and kidneys, cause spleen enlargement, decrease protein & fats digestion as well as absorption, cause impaired carbohydrate breakdown, cause reduced sperm count & infertility, birth defects, higher incidence of cancer, decreased resistance & increased susceptibility to infections.

6. CHEMICAL AND BIOLOGICAL EFFECTS OF AFLATOXINS

Aside from work related to structure elucidation, little systematic research has been carried out to find out the behavior & reactivity of aflatoxins, and agriculture products when treated with chemicals.

Aflatoxin B1 undergoes catalytic hydrogenation, which results in the adoption of 3moles of hydrogen and the formation of a tetra hydro deoxy derivative. When the hydrogenation process is interrupted after 1mole of hydrogen is taken up, aflatoxin B₂ is produced in a quantifiable quantity [19].

Aflatoxins can contaminate various agricultural products such as maize, g. nuts, barley, wheat, rice, sorghum, millet, soybeans, cassava, oilseeds, fruits, eggs, meat, animal feeds, and milk [20,21].

Aflatoxins enter the food and any step in the feed chain, from pre-harvest through consumption [22,23] causing many reactions within the cells. When the fungi colonize the plants in presence of favorable conditions, aflatoxins are produced which compromise the self-defense & growth of the plants and contaminate the crop seeds affecting the quality of the yield [9].

Once produced within cells, aflatoxins cause programmed cell death (apoptosis), and inhibit nucleic acid synthesis leading to a decrease in protein synthesis hence stunt growth; mutations hence affect the cell membrane stability leading to cell damage.

7. EFFECT OF AFLATOXINS ON HEALTH

A collection of naturally produced carcinogens known as aflatoxins is recognized to contaminate various foods used by both humans and animals [24]. Food products that have aflatoxins are a prevalent issue in the tropics and subtropics regions all over the world, particularly being developed nations with subpar sanitation systems and environments where fungi grow in environments with high humidity and warm temperatures [25]. Humans and animals can develop the disease known as aflatoxicosis as a result of aflatoxins. (I) Acute primary aflatoxicosis, which is one of its two main manifestations, is caused when moderate to high quantities of aflatoxins are eaten demonstrated in Fig. 5. Hemorrhage, abrupt liver injury, edema, and changes in digestion, food absorption, and metabolism are just a few of the symptoms of certain acute illness episodes that may occur [26]. (II) Consumption of Aflatoxin exposure in low to moderate amounts results in chronic primary aflatoxicosis. The impacts are typically subtle and difficult to spot. Some of the typical symptoms include poor food conversion and slower development rates, whether or not an overt aflatoxin illness has developed [27].

Aflatoxins' long-term effects have been shown to decrease normal immune responses by either lowering T cell or phagocytic activity quantity or functions, as seen in animal models of the disease. In a dosage response association between aflatoxins and growth rate in babies and

children, aflatoxins have also been documented to affect nutrition [28]. In animal studies, aflatoxins also modify nutrients like vitamin A or D, rendering them inaccessible to the body's normal physiology and resulting in nutritional deficits [29]. Aflatoxins are harmful to prenatal exposure as well, as through breast milk; they are transferred from mother to child. It has been discovered that AFB₁, in particular, interferes with the enzymes and substrates required for the production of several hormones, causing the various endocrine glands to malfunction [30].

Aflatoxin toxicity in humans can range from chronic or acute liver damage conditions such as liver carcinoma, internal bleeding, edema, and instant deaths [31]. AFB₁, AFB₂, and AFM were absorbed into the liver, bile duct, spleen, heart, muscle, and kidney. Mutation, cancer, immunodeficiency, lung damage, and birth abnormalities are all caused by these additional reactions [32]. Animals exposed to low dietary amounts of aflatoxins have immune suppression, reduced reproduction, liver damage, and decreased milk supply. In addition to fatty livers, and kidneys as well as cerebral edema and heart involvement, the symptom of aflatoxicosis in animal life may also be defined by nausea, stomach cramps, pulmonic embolism, coma, loss of consciousness, and death [33]. Acute toxicosis can cause gastrointestinal dysfunctions like ascites, intense pain, stomach cramps, bleeding in diarrhea, reduced feed intake, visual loss, circling, ear trembling, frothing at the lips, cumulative damage, and deaths in




Fig. 5. Aflatoxins disease pathway in humans

dairy and beef cattle, Other symptoms include tiredness, weight loss, depression, and sharp decreases in milk production weight loss, bleeding, [34].

The mitochondrial DNA, composition, and purpose of oxidative phosphorylation in brain cells are also abnormally affected by aflatoxins [35].

Acute treatment with AFB1 decreases regional brain acetylcholinesterase enzymes, which may influence a person's cognitive abilities, memory, and learning. Dopamine, serotonin, and the precursor's tyrosine and tryptophan levels are all affected by aflatoxin.

Neurological symptoms such as neurocognitive deterioration, altered sleep patterns, and brain damage indicators including dullness and uneasiness, muscular tremors, seizures, memory loss, seizure disorders, loss of motor control, and abnormal feelings are caused by deficiencies in these neurotransmitters [36]. Aflatoxins have been shown to have detrimental negative impacts on vascular fragility and cardiovascular health, tissue bleeding, and teratogenic consequences [37]. According to reports, aflatoxins have severe and detrimental impacts, mostly on the respiratory system. Also, one organ system with essential functioning components in continuous and direct interface with the environment is the respiratory system. Many people who work in the food industry are exposed to aflatoxins, particularly AFB1. These individuals have been linked to higher rates of upper respiratory tract and lung cancer [38]. Higher levels of aflatoxins have reportedly been found in the sperm cells of infertile men in people who have regularly consumed meals infected with the toxin [39].

Liver cancer (HCC): AFB1 was classified as a Group 1 carcinogen in 2002 because it causes the creation of DNA adducts that aid in the development of liver cancer [40]. According to research, AFB1 is a significant cause of hepatocarcinoma, with 4.6–28.2% of hepatocellular carcinoma (HCC) cases worldwide being linked to AFB1 exposure [41]. In addition, those exposed to AFB1 have a 30-fold increased risk of developing HCC if they have Hepatitis B virus (HBV) [42]. Acute hepatitis is brought on by high exposure levels, and liver cancer is then brought on by long-term exposure [43]. The genotoxic effects of AFB1, on the other hand, are still unknown. Recently, some authors treated the

human HL7702 hepatic cells with Microcystin-LR (MC-LR) and AFB1 and showed that MC-LR and AFB1 co-exposure: I induced DNA damage; and (ii) increased the activity of Superoxide Dismutase and Catalase, the levels of glutathione, and the expression of APE The link between AFB1 exposure and liver cancer is undeniable, so researchers are looking at how to create a diagnostic tool for early recognition of the consequences of AFB1 [44].

Moreover, liver tissues and blood from F344 rats exposed to AFB1 for 4 weeks were used to examine the impact of AFB1 on miRNA expression during the start phase of carcinogenesis. In both liver tissues and blood, these findings showed that AFB1 up-regulated miR-122-5p, 34a-5p, and 181c-3p more than controls, indicating that epigenetic modifications were brought about by AFB1 during the first stage of carcinogenesis [45]. Epigenetic processes are generally dysregulated in HCC patients following exposure to AFB1. Human hepatocytes actually exhibited six up-regulated and hypomethylated genes after AFB1 exposure, including Cyclin K (CCNK), Diaphanous Related Formin 3 (DIAPH3), Histone Cluster 1 H2B Family Member f (HIST1H2BF), RAS Oncogene 27A (RAB27A), Proliferating Cell Nuclear Antigen (PCNA), and Thioredoxin Reductase 1 (TXNRD1), which can be used as early markers [46].

Lung cancer: Workers who are exposed to pulmonary AFB1 exposure from grain dust are at risk of developing lung cancer [47]. As a result, AFB1 is classified as a pulmonary carcinogen, and several epidemiological studies have concentrated on the link between AFB1 and lung malignancies. Some authors looked at the AFB1 detoxification in lobectomy patients' lung tissues in 1996. Their research proved that lung cytosols produced a DNA-binding metabolite as a result of being exposed to AFB1. Its cytosolic activation can increase human lung susceptibility to AFB1 and is strictly associated with the activities of PGH synthase (PHS) and lipoxygenase (LOX) [48].

AFB1 can also cause oxidative DNA damage and lesions known as 8-oxo-7,8-dihydro-20-deoxyguanosine (8-oxodG) that can be fixed with the treatment of 8-oxoguanine glycosylase (OGG1). As a result, some researchers investigated the impact of OGG1 deficiency on cancer and oxidatively damaged DNA caused by AFB1. One dose of either DMSO or AFB1 was

administered to OGG1 null mice. OGG1 null mice treated with AFB1 lost weight and died, even though the incidence of lung cancer did not rise. The K-ras mutation pattern in lung tumor DNA was not consistent with AFB1-initiation. As a result, OGG1 status had no discernible impact on the DNA damage or tumorigenesis caused by AFB1, however, the deletion of OGG1 alleles led to an increase in sensitivity to additional AFB1 toxicity-related effects [49]. The effects of AFB1 on Src kinase and insulin receptor substrate (IRS) in lung cancer cells have recently been assessed, and cell migration was observed. After AFB1 therapy, the lung cancer cell lines A549 and SPCA-1 underwent a Western blot examination to assess IRS expression and Src, Akt, and ERK phosphorylation. According to this research, AFB1: (i) up-regulate IRS1 and IRS2; (ii) increases Src, Akt, and ERK1/2 activation; and (iii) stimulates lung cancer cell migration, which is blocked by saracatinib [50].

Gastrointestinal Cancer (GIC): Numerous research has examined the relationship between aflatoxins and the digestive system because, after exposure to contaminated food, the digestive system typically serves as the first line of defense against carcinogens. Harrison et al. hypothesized in 1993 that even in affluent nations, exposure to human aflatoxin could pose a risk to organs other than the liver. The authors looked for an AFB1-DNA adduct by analyzing normal and tumor samples from various organs (colon, rectum, breast, cervix, and liver) to support this claim. It's interesting to note that every sample tested positive, and colorectal tumor tissue in particular displayed a greater adduct concentration than normal tissues from the same patients [51].

The effects of AFB1 alone and in combination with other mycotoxins were assessed in Caco-2 cells. AFB1 was the third most cytotoxic mycotoxin that was evaluated, with a cytotoxic effect at a concentration of 19.28 M [52]. To confirm its impact on genotoxicity and DNA damage, AFB1 (3-5 M) was applied to HCT116 colorectal cancer cells in a different investigation. According to this study, AFB1 treatment: (i) slightly increased H2AX phosphorylation in HCT116 cells by inducing an ATM response; (ii) did not activate P53, CHK1, or CHK2, and (iii) did not block the G1 phase of the cell cycle [53]. Authors in 2016 confirmed that AFB1 exposure causes MDM2 up-expression, a p53-negative modulator, in murine intestinal cancer cells [54].

8. CONCLUSION

Information on the contamination and colonization of fungi and the negative impacts associated with it through the production of aflatoxins in agriculture products and most of the developing countries are still lacking due to a lack of public awareness of their effects on the quality of agriculture products and their potentially hazardous effects on human and animals health. Therefore focusing on increasing public awareness of aflatoxins and their effects on agriculture, health, and economic growth is required.

In addition, target genes were altered by AFB1 and AFM1 in light of the paucity of evidence indicating whether genes, proteins, or miRNAs might be employed as damage markers owing to exposure. According to the network analysis, the thirteen proteins (CYP3A4, TP53, GSMT1, MDM2, CAT, OGG1, IRS1, IRS2, SRC, AKT1, MAPK1, MAPK3, and PDK1) that are regulated by AFM1/AFB1 are connected both directly and indirectly through nodes like NR112 and GNMT. However, the research on miRNA prediction has shown that twelve miRNAs can specifically target the genes that are known to be affected by AFM1.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Assaf JC, Nahle S, Chokr A, Louka N, Atoui A, El Khoury A. Assorted methods for decontamination of aflatoxin M1 in milk using microbial adsorbents. *Toxins (Basel)*. 2019;11(6).
2. Klich MA, Tang S, Denning DW. Aflatoxin and ochratoxin production by *Aspergillus* species under ex vivo conditions. *Mycopathologia*. 2009;168(4):185-91.
3. Guchi E. Implication of aflatoxin contamination in agricultural products. *Am J Food Nutr*. 2015;3(1):12-20.
4. Battilani P, Toscano P, Der Fels-Klerx V, Moretti A, Camardo LM, Brera C, Robinson T. *Sci Rep*. 2016. Aflatoxin B1 contamination in maize in Europe increases due to climate change;6(1):1.
5. Moretti A, Pascale M, Logrieco AF. Mycotoxin risks under a climate change scenario in Europe. *Trends Food Sci Technol*. 2019;84:38-40.

6. Coppock RW, Christian RG, Jacobsen BJ. Aflatoxins. In: Veterinary toxicology. Academic Press; 2018. p. 983-94.
7. Pitt JI, Hocking AD. Fungi and food spoilage. New York: Springer. 2009;519:388.
8. Inan F, Pala M, Doymaz I. Use of ozone in detoxification of aflatoxin b1 in red pepper. J Stored Prod Res. 2007;43(4): 425-9.
9. Kumar A, Pathak H, Bhadauria S, Sudan J. Aflatoxin contamination in food crops: causes, detection and management: a review. Food Prod Process Nutr. 2021;3(1):1-9.
10. Rustom IYS. Aflatoxin in food and feed: occurrence, legislation and inactivation by physical methods. Food Chem. 1997;59(1):57-67.
11. Jasutiene I, Garmiene G, Kulikauskiene M. Pasteurization and fermentation effects on aflatoxin M1 stability. Milchwissenschaft. 2006;61(1):75-9.
12. Jalili M. A review on aflatoxins reduction in food. Iran J Health Saf Environ. 2016;3(1):445-59.
13. Echodu R, Maxwell Malinga G, Moriku Kaducu J, Ovuga E, Haesaert G. Prevalence of aflatoxins, ochratoxin and deoxynivalenol in cereal grains in Northern Uganda: implication for food safety and Health. Toxicol Rep. 2019;6:1012-7. Available:<https://doi:10.1016/j.toxrep.2019.09.002>
14. Omara T, Nassazi W, Omute T, Awath A, Laker F, Kalukusu R et al. Aflatoxins in Uganda: an Encyclopedic Review of the Etiology, Epidemiology, Detection, Quantification, Exposure Assessment, Reduction and control. Int J Microbiol. 2020;2020:4723612.
15. Smith JE. Aflatoxins. In: Handbook of plant and fungal toxicants. CRC Press. 2020;269-85.
16. Peles F, Sipos P, Györi Z, Pfliegler WP, Giacometti F, Serraino A et al. Adverse effects, transformation and channeling of aflatoxins into food raw materials in livestock. Front Microbiol. 2019;10:2861.
17. Ráduly Z, Szabó L, Madar A, Pócsi I, Csernoch L. Toxicological and medical aspects of Aspergillus-derived mycotoxins entering the feed and food chain. Front Microbiol. 2019;10(2908):2908.
18. Robens JF, Richard JL. Aflatoxins in animal and human health. Rev Environ Contam Toxicol. 1992;127:69-94.
19. Wogan GN. Chemical nature and biological effects of aflatoxins. Bacteriol Rev. 1966;30(2):460-70.
20. Tola M, Kebede B. Occurrence, importance and control of mycotoxins: a review. Cogent Food Agric. 2016;2(1): 1191103.
21. Lizarraga-Paulin E, Miranda-Castro S, Moreno-Martinez E, Torres-Pacheco I, Lara-Sagahon AV. Novel methods for preventing and controlling aflatoxins in food: A worldwide daily challenge. Aflatoxins-recent Advances and future Prospects Razzaghi-Abyaneh M, editor. 2013;93-128.
22. Torres AM, Barros GG, Palacios SA, Chulze SN, Battilani P. Review on pre- and post-harvest management of peanuts to minimize aflatoxin contamination. Food Res Int. 2014;62:11-9.
23. Norlia M, Jinap S, Nor-Khaizura MAR, Radu S, Samsudin NIP, Azri FA. Aspergillus section Flavi and aflatoxin: occurrence, detection and identification in raw peanuts and peanut-based products along the supply chain. Front Microbiol. 2019;10(2602):2602.
24. SAB, AA. Mycotoxins in food in West Africa: Current situation and possibilities of controlling it. Afr J Biotechnol. 2003;2(9):254-63.
25. Bennett JW, Klich MA. Mycotoxins. Clin Microbiol Rev. 2003;16(3):497-516:Article CAS.
26. Aflatoxins NL. National Library of Medicine. Hazardous substance data base. Toxnet (National Data Network); 2002.
27. WHO hazardous chemicals in humans and environmental health. Geneva, Switzerland: International Programme on Chemical safety. World Health Organisation. 2000;7:9. Available:http://whqlibdoc.who.int/hq/2000/WHO_PCS_00.1.pdf
28. Peraica M, Radić B, Lucić A, Pavlović M. Diseases caused by molds in humans. Bull World Health Organ. 1999;77(9): 754-66.
29. World Health Organization. Collaboration between the World Health Organization and the National Institute of Environmental Health Sciences: Highlights from 30 years of partnership.
30. Barrett JR. Liver cancer and aflatoxin: New information from the Kenyan outbreak. Environ Health Perspect. 2005;113(12).

31. Murthy TR, Jemmali M, Henry Y, Frayssinet C. Aflatoxin residues in tissues of growing swine: effect of separate and mixed feeding of protein and protein-free portions of the diet. *J Anim Sci.* 1975;41(5):1339-47.
32. Thrasher JD. Aflatoxicosis in animals. *Aflatoxins and health*; 2012 Available:www.alphaboostjuice.com/AFLA_TOXICOSIS_IN_ANIMALS.pdf
33. Fapohunda SO, Awoyinka OA, Olajuyigbe OO, Ezekiel CN, Esiaba I. Enzyme-related aflatoxin production in vital organs of rats fed with *Aspergillus* species-inoculated rat chow. *J Biol Environ Sci.* 2007;1(1).
34. Verma RJ. Aflatoxin cause DNA damage. *Int J Hum Genet.* 2004 Dec 1;4(4):231-6.
35. Coulombe Jr RA. Nonhepatic disposition and effects of aflatoxin b1. *The Toxicol Aflatoxins Hum Health Vet Agric Significance.* 1993:89-101.
36. Weekley LB, O'Rear CE, Kimbrough TD, Llewellyn GC. Differential changes in rat brain tryptophan, serotonin and tyrosine levels following acute aflatoxin b1 treatment. *Toxicol Lett.* 1989;47(2):173-7.
37. Harriet AM. Is indoor mold contamination a threat to health? *J Environ Health.* 2003;62(2):0022892 62 2:0022-892.
38. Coulombe Jr RA. Nonhepatic disposition and effects of aflatoxin b1. *The Toxicol Aflatoxins Hum Health Vet Agric Significance.* 1993:89-101.
39. Gupta RC. Aflatoxins, ochratoxins and citrinin. *Reproductive and developmental Toxicology.*
40. Nugraha A, Khotimah K, Rietjens IMCM. Risk assessment of aflatoxin b1 exposure from maize and peanut consumption in Indonesia using the margin of exposure and liver cancer risk estimation approaches. *Food Chem Toxicol.* 2018;113:134-44.
41. Kucukcakan B, Hayrulai-Musliu Z. Challenging role of dietary aflatoxin b1 exposure and hepatitis B infection on risk of hepatocellular carcinoma. *Open Access Maced J Med Sci.* 2015;3(2):363-9.
42. Liu Y, Wu F. Global burden of aflatoxin-induced hepatocellular carcinoma: A risk assessment. *Environ Health Perspect.* 2010;118(6):818-24.
43. Rapisarda V, Loreto C, Malaguarnera M, Ardiri A, Proiti M, Rigano G; et al. Hepatocellular carcinoma and the risk of occupational exposure. *World J Hepatol.* 2016;8(13):573-90.
44. Liu W, Wang L, Zheng C, Liu L, Wang J, Li D et al. Microcystin-LR increases genotoxicity induced by aflatoxin b1 through oxidative stress and DNA base excision repair genes in human hepatic cell lines. *Environ Pollut.* 2018;233:455-63.
45. Livingstone MC, Johnson NM, Roebuck BD, Kensler TW, Groopman JD. Profound changes in miRNA expression during cancer initiation by aflatoxin b1 and their abrogation by the chemopreventive triterpenoid CDDO-Im. *Mol Carcinog.* 2017;56(11):2382-90.
46. Rieswijk L, Claessen SM, Bekers O, van Herwijnen M, Theunissen DH, Jennen DG et al. Aflatoxin b1 induces persistent epigenomic effects in primary human hepatocytes associated with hepatocellular carcinoma. *Toxicology.* 2016;350-352:31-9.
47. Hayes RB, Van Nieuwenhuize JP, Raatgever JW, ten Kate FJW. Aflatoxin exposure in the industrial setting: an epidemiological study of mortality. *Food Chem Toxicol.* 1984;22(1):39-43.
48. Donnelly PJ, Stewart RK, Ali SL, Conlan AA, Reid KR, Petsikas D et al. Biotransformation of aflatoxin b1 in human lung. *Carcinogenesis.* 1996;17(11):2487-94.
49. Mulder JE, Turner PV, Massey TE. Effect of 8-oxoguanine glycosylase deficiency on aflatoxin b1 tumorigenicity in mice. *Mutagenesis.* 2015;30(3):401-9.
50. Cui A, Hua H, Shao T, Song P, Kong Q, Luo T et al. Aflatoxin b1 induces Src phosphorylation and stimulates lung cancer cell migration. *Tumour Biol.* 2015;36(8):6507-13.
51. Harrison JC, Carvajal M, Garner RC. Does aflatoxin exposure in the United Kingdom constitute a cancer risk? *Environ Health Perspect.* 1993;99:99-105.
52. Sobral MMC, Faria MA, Cunha SC, Ferreira IMPLVO. Toxicological interactions between mycotoxins from ubiquitous fungi: impact on hepatic and intestinal human epithelial cells. *Chemosphere.* 2018;202:538-48.
53. Gursoy-Yuzugullu O, Yuzugullu H, Yilmaz M, Ozturk M. Aflatoxin genotoxicity is associated with a defective DNA damage response bypassing p53 activation. *Liver Int.* 2011;31(4):561-71.

54. Kim J, Park SH, Do KH, Kim D, Moon Y. Interference with mutagenic aflatoxin b1-induced checkpoints through antagonistic action of ochratoxin A in intestinal cancer cells: A molecular explanation on potential risk of crosstalk between carcinogens. *Oncotarget*. 2016;7(26): 39627-39.

© 2022 Ahmad et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92468>