



Microbiological and Nutritional Quality of Complementary Foods for Children (6 to 24 months) Produced in the City of Man (Côte d'Ivoire)

**Koffi Affoué Carole ^{a*}, Kossonou Yao Kamelé ^a,
Adingra Kouassi Martial-Didier ^b,
Yapi Aka Eugène-Janvier ^a, Kouassi Kouassi Clément ^c
and Rose Koffi-Nevry ^b**

^a *Department, of Food Process and Bioproducts, University of Man (Côte d'Ivoire), UFR Agronomic, Forestry and Environmental Engineering, University of Man, BP 20 Man, Côte d'Ivoire.*

^b *Department of Food Science and Technology, University of Nangui Abrogoua, Côte d'Ivoire.*

^c *Department of Biochemistry and Microbiology, Laboratory of Agrovalorization, Jean Lorougnon Guede University, Côte d'Ivoire.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/MRJI/2022/v32i91344

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93975>

Original Research Article

Received: 01/10/2022

Accepted: 04/12/2022

Published: 16/12/2022

ABSTRACT

This study aims to determine the microbiological and nutritional quality of complementary foods for children produced in the city of Man (instant flour and porridge). To do this, a structured survey of nannies and vendors, coupled with microbiological and physicochemical analyzes was carried out.

*Corresponding author: E-mail: carole.koffi@univ-man.edu.ci, kofficar@yahoo.fr;

To this end, a questionnaire survey was conducted among nannies and porridge sellers in order to collect information on the use of complementary foods and to assess the state of cleanliness of the sales environment. A study of physicochemical (carbohydrates, lipids, humidity, ash, minerals) and microbiological (total aerobic mesophilic flora, yeasts and molds, total coliforms, *Escherichia coli*, *Staphylococcus sp.* and *Salmonella sp.*) characteristics using standard methods and referenced was then conducted to assess the nutritional value and health status of the porridges (Millet, Kokobaka, Anagobaka) and flours from the Focolari center consumed in Man. The survey data showed that the majority of the nannies (61.1%) give a complementary food before the age of introduction (6 months), with regard to the saleswomen (100%) unaware of the proportion of the components and their nutritional intake as well as 54.8% of them have an unhealthy work place and environment. The microbiological analysis carried out on the 50 samples revealed a potential contamination of spoilage and pathogenic germs. The contamination rate of Anagobaka porridge is very worrying unlike other porridges and flours, especially with the presence of *salmonella*, the high rate of *Escherichia coli* and *Staphylococcus aureus* respectively ($3.91 \times 10^3 \pm 0.09$ and $5.34 \times 10^4 \pm 0.10$ CFU/mL.) Regarding the nutritional analysis, there is a content of carbohydrates (70.17-85.08%), lipids (6.93-10.40%) and iron (30.64-64.34%) higher than those recommended by the FAO/WHO standard in the 6 types of samples analyzed. However, the zinc content is below this standard in all the samples tested. However, zinc deficiency is one of the main causes of stunted growth in children. All these results have shed light on the existence of a potential health risk and possible nutritional deficiencies in infants and young children who consume locally produced complementary foods.

Keywords: Microbiological quality; complementary food; physico-chemical characteristics; germs; Man (Côte d'Ivoire).

1. INTRODUCTION

Malnutrition remains a public health problem throughout the world. Nearly 112 million children under five worldwide are underweight and 178 million are stunted [1]. In developing countries, child malnutrition is one of the main causes of public health and social well-being problems, it is the cause of half of child deaths [2]. In Côte d'Ivoire, approximately 45 % of annual deaths of children under 5, i.e. 42,000 deaths per year or 115 deaths per day, are due to malnutrition [3]. All these forms of malnutrition are more severe during the weaning period, the so-called complementary feeding period (6 to 24 months) [4].

During this period, it is necessary to bring new foods in liquid or semi-liquid form to supplement the contributions of breast milk. These new and first foods given to children are called complementary foods and must be of satisfactory nutritional, health and organoleptic quality [5]. Indeed, according to the recommendations of the WHO and UNICEF, infants must receive exclusive breastfeeding up to 6 months. However, beyond 6 months, breast milk alone is not enough to meet their nutritional needs. They should therefore receive complementary feeding while continuing to breastfeed for up to 2 years or more [5]. These complementary foods must be

foods of nutritional value appropriate to the age of the child and uncontaminated from both a microbiological and toxicological point of view. Thus, the first complementary foods are most often porridge made from cereals, roots or tubers, rich in carbohydrates and low in protein [6]. Porridge is sold in precarious hygienic conditions, which constitutes a major public health risk for consumers, especially children under five [7].

Like several local products, infant flour, the main complementary food, is produced essentially using an artisanal or semi-artisanal process under conditions of insufficient sanitation by millers with a relatively low level of education [8]. In addition, in Côte d'Ivoire, imported infant flours exist on the market as well as those produced locally as complementary foods. These complementary foods sold in supermarkets and large commercial chains are inaccessible to a large part of the population and are generally not used by low-income households because they are relatively expensive [8].

However, infant flours and porridges made traditionally are accessible to all, but they have a very low nutritional value and cause problems of malnutrition and nutritional deficiencies [9].

Studies have focused on complementary feeding of young children in Côte d'Ivoire [9,10]. However, at the level of the city of Man, no study has yet been undertaken concerning the microbiological and nutritional quality of complementary foods.

In addition, according to multiple indicator cluster survey [11], the West of Côte d'Ivoire (area where the city of Man is located) is an area with a high prevalence of malnutrition, making the bed of several childhood pathologies including malaria, respiratory infections acute and diarrhoea. In the west of Côte d'Ivoire, 26 % of children under 5 are chronically malnourished compared to 21.6 % nationally and 6 % of children in this area of Côte d'Ivoire with this same age bracket are emaciated.

These different situations have aroused the interest of carrying out this study which aims to know the formulation and evaluate the nutritional and microbiological quality of complementary foods produced in the city of Man (Côte d'Ivoire).

2. MATERIALS AND METHODS

2.1 Materials

2.1.1 Plant material

The plant material consists of 6 types of flour composed as follows: a flour made from maize (*Zea mays*), sesame and cocoa (*Theobroma cacao*) (FMSC); one based on maize (*Zea mays*) and soy (*Glycine max*) (FMS); another based on maize (*Zea mays*), soy (*Glycine max*), cocoa (*Theobroma cacao*), sesame (*Sesamum indicum*), rice (*Oriza sativa*), millet (*Pennisetum glaucum*) and banana (*Musa paradisiaca*) (FM5B); one based on maize (*Zea mays*), ginger (*Zingiber officinale*) and/or pepper (*Capscicum annum*) used for "Kokobaka porridge (KP)", a flour made of maize (*Zea mays*) only: "Anagobaka porridge (AP)". And the last one based on millet (*Pennisetum glaucum*) used for "millet porridge (MP)". Among these flours, it should be noted that three (FMS, FMSC and FM5B) come from the Focolari Medico-Social Center of Man (Côte d'Ivoire).

2.1.2 Technical material

The technical equipment used to carry out this work consists of survey sheets and usual microbiology and biochemistry laboratory equipment. Plat Count Agar (PCA) culture medium was used for the enumeration of Total

Mesophilic Aerobic Flora (TMAF) and Baird Parker (BP) agar was used for the enumeration of *Staphylococcus sp.* As for the count of *Salmonella sp.* and *Escherichia coli (E. coli)*, it was done with the respective media *Salmonella-Shigella (SS)* and Violet Red Bile Lactose (VRBL). VRBL medium was also used for the enumeration of coliforms. Buffered Peptone Water (EPT) broth was used for preparation of the stock solution and for the various dilutions.

2.2 Methods

2.2.1 Survey methodology

2.2.1.1 Sampling

2.2.1.1.1 Choice of neighborhoods

The main neighborhoods visited were Camp Sea, Cafop, Kôkô, Trade, Dioulabougou and Blockauss, which are home to the majority of vendors selling porridge made from local products. These neighborhoods were selected following a pre-survey carried out by our research team.

2.2.1.1.2 Choice of the Focolari medico-social center in Man (Côte d'Ivoire)

This survey aims to provide information on infant foods offered to children in the city of Man. This center regularly produces flour that it markets, which is why we chose flour from this center for our work.

2.2.1.1.3 Choice of porridge saleswomen

The purpose of this survey was to define the ingredients used to make infant porridge offered to children in the city of Man and to highlight the preparation conditions and the state of cleanliness of the sales environment, the saleswoman, the equipment, the presence or absence of flies on the food or all around the point of sale. The list of the different traders selected was provided from a pre-survey. An exhaustive sampling of all the porridge vendors in these six neighborhoods selected for this study was done. There are 31 vendors surveyed, including 6 in Camp Sea, 4 in Cafop, 7 in Kôkô, 3 in Trade, 8 in Dioulabougou and 3 in Blockauss.

2.2.1.1.4 Choice of mothers (nannies)

The size (n) of the samples (mothers of children) per neighborhood was obtained using the

formula of FAO [12] for a non-exhaustive independent sample as follows:

$$n = \frac{t^2 \times p(1-p)}{e^2} \quad (1)$$

n = minimum size sought for the sample;
 t = 95% confidence level (typical value of 1.96);
 p = estimated prevalence of wet nurses in the study area, p estimated at 50% since the total number of wet nurses in the area is not known; e = 5% margin of error (value of 0.05).

Thus, the minimum size of nurses sought was 90 nurses. Therefore, the selection of individuals was made by the snowball technique. The criterion used is that of the representativeness of the sampling of actors surveyed (nurses) by district. Thus, this study made it possible to survey 90 nurses, that is 15 nannies per neighborhood.

2.2.1.1.5 Conduct and data collection of the survey

The methodology adopted with porridge saleswomen and nannies (mothers of children) was that of a questionnaire. This questionnaire was established using the SphinxPlus.V5.TuiTe software. To this end, to put the interlocutor in confidence and obtain reliable data, the investigators used politeness, declined their identities and presented the framework of their work. Interviews only begin when the nannies and vendors consent to the interview. Sometimes, when the respondent does not speak French, the interview was conducted in the local language. Interpretation in the local language was done with the help of an interpreter proficient in this language. The interview begins with recording information about the respondent's identification and basic information: name, sex, age, nationality and level of education. Then the questions on the practices

and knowledge of complementary feeding, the contribution of these foods to young children, were addressed with regard to mothers of children. From the saleswomen surveyed, relevant information was collected, in particular respect for good hygiene practices, the mode of installation of the vendor (outdoors, under a shelter or itinerant), the formulation of complementary foods. In addition, the state of cleanliness of the sales environment, the saleswoman, the equipment, the presence of flies on the food or all around the place of sale and the existence of a place to deposit garbage has been addressed.

2.2.2 Complementary food flour collection

Different types of infant flours were sampled in sterile sachets at a rate of 500 g per sachet. The samples were taken and analyzed progressively following five (05) successive productions for each infant flour at the level of the production structure (Medico-Social Center Focolari of Man). As for the porridge, they were purchased and contained in sterile bags through 3 sectors (2 districts per sector) of the city of Man. Five (05) samples of each type of mixture were analyzed at 2-day intervals. A total of 50 samples were used for this study, consisting of fifteen (15) 500 g flour sachets and 35 approximately 300 g sachets of 3 different porridges (Table 1). Each sample was put in a cooler and transported to the laboratory for analysis.

2.2.3 Manufacture of complementary foods

The artisanal processes developed by Aboua et al. [13] were used to set up the different flours of this work. The processes for preparing the porridges and flours studied are represented by Figs. 1, 2 and 3 as practiced by the Focolari center and most of the porridge saleswomen surveyed.

Table 1. Number of samples taken in several districts of the city of Man

Samples	Camp Sea / Cafop	Kôkô / Trade	Dioulabougou / Blockauss	Focolari
KP	5	5	5	None
MP	5	5	5	None
AP	None	5	None	None
FMS	None	None	None	5
FMSC	None	None	None	5
FM5B	None	None	None	5

Total sample number = 50

KP: Kokobaka porridge; MP: Millet porridge; AP: Anagobaka porridge; FMS: Infant maize and soy flour; FMSC: Flour made from maize, sesame and cocoa; FM5B: Infant flour made from maize, sesame, soy, millet, rice, cocoa and banana; None = 0




Fig. 1. Millet porridge production diagram

2.2.4 Methods of microbiological analysis of flours and porridges

The microbiological analysis of infant flours and porridges is based on techniques for counting, isolating and identifying microorganisms likely to alter the hygienic and/or nutritional qualities of flours and porridges. These are germs such as faecal and total coliforms, *Escherichia coli*, *Staphylococcus sp.*, yeasts and molds, total flora and *Salmonella*. The identification of a germ, qualitative aspect is characterized by the absence or the presence of the latter. On the other hand, the counting of a germ is carried out by counting the colonies followed by a method of calculation and therefore represents the quantitative aspect of it.

2.2.4.1 Preparation of culture media, stock suspension and dilutions

The culture media were prepared according to the manufacturer's instructions.

2.2.4.1.1 Preparation of stock suspension and tenfold dilutions

The preparation of the samples, the stock suspension and the decimal dilutions was carried out according to the international standard ISO 6887-1 (1999). The samples were analyzed immediately after collection. A quantity of 25 g or 25 ml of sample is withdrawn aseptically into a sterile stomacher bag to which 225 ml Buffered Peptone Water (EPT) is added. The mixture was

homogenized and left for 15-20 min for revivification of microorganisms. The solution obtained corresponds to the stock suspension. From this stock suspension, a series of successive decimal dilutions was carried out. One milliliter of the stock suspension was taken using a pipette and then introduced into a test tube containing 9 mL of Buffered Peptone Water. This corresponds to the 10^{-1} solution. The operation continued until the 10^{-5} dilution was obtained.

2.2.4.1.2 Research and enumeration of the different bacterial flora

After the incubation period mentioned in the standard specific to each germ, the count of the characteristic colonies for each dish containing

less than 300 colonies and 30 colonies at least and between 15 and 150 colonies (for coliforms) was carried out. The number N of colony forming units (CFU)/g present in the sample analyzed and considered as an average of successive dilution is given by the following formula:

$$N = (C_1 + C_2) / V(n_1 + 0.1n_2) d \quad (2)$$

$C_1 + C_2$ = sum of the characteristic colonies of two boxes of successive dilutions counted

V = volume of inoculum applied to each dish

d = dilution rate corresponding to the first dilution retained

n_1 = number of dishes retained at the first dilution

n_2 = number of dishes retained at the second dilution




Fig. 2. Production diagram of Anagobaka and Kokobaka porridges




Fig. 3. Diagram of flour production at the Focolari medical-social center in Man

2.2.4.1.3 Enumeration of mesophilic aerobic flora

The enumeration of mesophilic aerobic flora was carried out according to the ISO international standard 4833 (2003). Inoculation was done by exhaustion streaking on Plate

Count Agar (PCA) previously poured into Petri dishes and then incubated in an oven at 37 °C for 24 h ± 2 h. After the incubation period the colonies were counted. The characteristic colonies appeared whitish or yellowish.

2.2.4.1.4 Search for *Staphylococcus* sp.

For the research of *Staphylococcus* sp. suspected pathogens was carried out according to ISO international standard 6888-1 (2003); 0.1 mL of stock solution and of each dilution were inoculated by streaking on Petri dishes containing BP medium (Baird Parker) and then incubated at 37 °C. for 48 h. Dishes containing 15 and 150 characteristic colonies were retained. The black, shiny, domed colonies surrounded by a white precipitate and a clearing halo were counted.

2.2.4.1.5 Enumeration of total coliforms and *Escherichia coli*

Total and thermotolerant coliforms were counted respectively according to the standard International ISO 4832 (2006) and ISO 16649-2 (2001). Inoculation was done on bile agar with crystal violet and neutral red (VRBL agar) then incubated at 37 °C for total coliforms and at 44 °C for thermotolerant coliforms, in an oven for 24 h ± 2 h. Characteristic colonies were counted after the incubation period. Thus, the dishes containing 15 and 150 characteristic colonies at the level of 2 successive dilutions were retained.

2.2.4.1.6 Enumeration of yeasts and molds

The culture medium used is Sabouraud poured into Petri dishes. 1 mL of each dilution was inoculated on Petri dishes and then incubated at 25 °C. for 2 days. Yeast colonies are milky in appearance while mold colonies are filamentous.

2.2.4.1.7 Search for *Salmonella* sp.

Research of *Salmonella* sp. was carried out according to ISO international standard 6579/A1 (2007). The search for *Salmonella* in food involves essential steps: pre-enrichment, enrichment, isolation and confirmation.

- **Pre-enrichment**

The stock solution containing 25 g of sample mixed in 225 mL of EPT then homogenized, was incubated at a temperature of 37 °C for 24 ± 2 h.

- **Selective enrichment**

0.1 mL of the solution from the pre-enrichment was inoculated into a test tube containing 10 mL of rappaport vassiliadis broth. The mixture was homogenized and then incubated at 37 °C for 24 ± 2 h.

- **Isolation**

The isolation is done from the cultures obtained from the selective enrichment medium. Petri dishes containing *Salmonella-Shigella* culture medium were streaked. The plates thus inoculated were incubated for 24 hours at 37°C. The typical colonies of *Salmonella* sp. are black with a transparent halo on *Salmonella-Shigella* agar and small in size (2 to 4 mm in diameter).

- **Confirmation**

Confirmation is made by purification of the colonies in Petri dishes containing the *Salmonella-Shigella* medium then incubated at 37 °C for 24 hours.

2.2.4.1.8 Interpretations of microbiological data

At the end of the microbiological analysis, the interpretation of the results was made according to a 3-class sampling plan for TMAF, *E. coli*, total coliforms, *Staphylococcus* sp. suspected pathogens, yeasts and molds and a 2-class sampling plan for *Salmonella* sp.

The N values (CFU/g or mL) calculated for each flora studied according to the samples of each type of porridge and flour, were compared with the normative reference of the microbiological criteria applicable to foodstuffs intended for infants and young children (Table 2).

- For the 3-class plan, when the values obtained are lower than the minimum microbiological criterion set by germ according to standard NFV08-010-Mars (1996) and up to three (3) times this criterion, the product is of microbiological quality satisfactory (SMQ) ($m \leq N \leq 3m$).
- The product is of acceptable microbiological quality (AMQ) for the values obtained between three (3) and ten (10) times the criterion ($3m < N \leq 10m$).
- The microbiological quality is unsatisfactory (UMQ) ; when the germ count values obtained are greater than ten (10) times the criterion ($N > 10m$).
- For *salmonella* sp., the presence of *salmonella* indicates that the sample is of unsatisfactory microbiological quality (UMQ). The sample is of microbiological quality satisfactory (acceptable microbiological quality) if they are absent.

Table 2. Microbiological standards for infant flour and porridge (NFV08-010- March 1996)

Microorganisms	Criteria CFU/g or mL	
Total aerobic mesophilic flora	10 ⁴	10 ⁵
Yeasts and molds	10 ³	10 ⁴
Total coliforms	10 ³	10 ⁴
<i>Escherichia coli</i>	1	10
<i>Staphylococcus sp</i>	10	10 ²
<i>Salmonella sp</i>	Absent in 25 g or mL	

2.2.5 Biochemical analysis methods for flours and porridges

The water content of the flours was obtained according to AOAC method [12] using an oven at a temperature between 103 and 105 °C until a constant weight was obtained. The water content corresponding to the variation in weight of the flour during the operation. The ash content was determined by dry incineration according to AOAC method [14]. Total lipids were extracted from samples homogenized in chloroform-methanol-water (1:2:1, v/v/v) as described by Folch et al. [15] and determined by gravimetry. As for total carbohydrates, they were quantified by the method of Munson and Walker [16]. Regarding the content of minerals, it was determined by the X-ray fluorescence spectrometry technique described by Thirion-Merle [17].

3. RESULTS

3.1 Survey Results

3.1.1 Survey of mothers/children

At the end of the survey carried out among mothers with children aged between 6-24 months, the results obtained are shown in Table 3. A proportion of 20% of the mothers questioned know what a complementary food against 80% of them who ignore it. Among those who use a complementary food, 61.1% of mothers claim to have introduced this food before the age of 6 months, 27.8% of them introduced it from 6 months and 11.1 % beyond 6 months (Table 3). The majority of nannies (mothers) surveyed (32.2%) declared that Anagobaka porridge is the most given to children, followed by millet porridge (24.4%).

The survey revealed that most complementary foods are purchased by mothers at the market or supermarket, i.e. 39.5%. Those who design these foods themselves at home are 27.6%. A proportion of 17.8% say they buy these foods in

the neighborhood, 11.8% in a health center (Table 3).

3.1.2 Survey carried out among sellers

The survey carried out among the vendors revealed that the level of education of the vendors questioned is distributed as follows: 38.7 % of these women have a primary education level; 32.3 % of them have not been to school; 16.1 % of these women have attended Koranic school and 12.9% have a secondary education. No saleswoman has a higher level of education (Table 3). Concerning the knowledge of complementary food, 74.2% of saleswomen are unaware of the usefulness of these complementary foods against 25.8 % of them who know it (Table 3). Regarding the knowledge and practice of hygiene of saleswomen along the production line, 88.9 % of saleswomen know and practice hygiene, unlike 11.1 % of saleswomen (Table 3). The state of cleanliness of the sales environment based on the abundant presence of flies on the food or all around the place of sale reveals that 54.8% of the saleswomen work in an unhealthy environment against only 3.2% of those who sell in a safe environment.

Several relevant data were collected through this survey sheet addressed to vendors of locally produced complementary food in the city of Man. Among these data we have the mode of installation of the saleswoman, the duration of sale of the complementary food.

3.2 Results of Bacteriological Analyzes

3.2.1 Bacterial load of complementary foods (porridges and flours)

A total of 50 samples were analyzed. Tables 4, 5, 6 and 7 give a summary of the average contamination level of the porridge and flour germs of all the samples for each germ. The average microbial load of the Anagobaka porridge of the samples analyzed revealed an unsatisfactory microbiological quality (UMQ) for

Table 3. Socio-economic characteristics and data on complementary foods distributed in Man

	Survey results data					
	Mothers		Saleswomen			
Complementary food knowledge	Yes	20%	Educational level	Primary	38,70%	
	No	80%		Not in school	32,30%	
Age of start of consumption of the complementary food	<6 month	61,10%	Level of knowledge of complementary food	Koranic school	16,10%	
	6 month	27,80%		Secondary	12,90%	
	> 6 month	11,10%		Oui	74,20%	
Use of locally manufactured complementary food	Yes	94,40%	Knowledge and practice of hygiene	Non	25,80%	
	No	5,60%		Yes	88,90%	
Food preference given to children	Anagobaka porridge	32,20%	Proportion of components and nutritional value of complementary foods	No	11,10%	
	Porridge from the Focolari medical and social center	20%		Yes	0%	
	Kokobaka porridge	18,90%		No	100%	
	millet porridge	24,40%		Poor	54,80%	
	rice porridge	3,30%		Acceptable degree	41,90%	
	other food	1,10%	Sanitary level of the place of sale	Good	3,20%	
	Advice for the use of the complementary food	Feeders	62,50%			
		Health center	19,20%			
		Friends and parents	13,30%			
		Culture	4,20%			
Other		0,80%				
Place of supply of complementary foods	Market or supermarket	39,50%				
	At home	27,60%				
	The neighborhood	17,80%				
	Health center	3,30%				

Table 4. Contamination level of millet porridge

Germs studied		Samples of millet porridge taken from quarters (n=5 per quarter)			Microbiological standards	
		Camp Sea/Cafop	Kôkô/Trade	Dioulabougougou/ Blockauss		
TMAF	Charge CFU/mL	4,79.10 ⁴ ± 0,10	8.92.10 ⁴ ± 2.12	8.58.10 ⁴ ± 0.09	10 ⁴	10 ⁵
	Quality	AMQ	AMQ	AMQ		
Yeasts and molds	Charge CFU/mL	0	0	0	10 ³	10 ⁴
	Quality	SMQ	SMQ	SMQ		
Total coliforms	Charge CFU/mL	4.14x10 ³ ± 0.07	4.58x10 ⁴ ± 1.05	3.58x10 ⁴ ± 0.05	10 ³	10 ⁴
	Quality	AMQ	UMQ	UMQ		
Escherichia coli	Charge CFU/mL	0	0	0	1	10
	Quality	SMQ	SMQ	SMQ		
Staphylococcus sp.	Charge CFU/mL	2.21x10 ³ ± 0.07	3.21x10 ³ ± 0.04	5.33x10 ³ ± 0.16	10	10 ²
	Quality	UMQ	UMQ	UMQ		
Salmonella sp.	Absent in 25 mL	Absent	Absent	Absent	Absent in 25 mL	
	Quality	SMQ	SMQ	SMQ		

Table 5. Contamination level of Kokobaka porridge

Germs studied		Samples of Kokobaka porridge taken from quarters (n=5 per quarter)			Microbiological standards	
		Camp Sea/Cafop	Kôkô/Trade	Dioulabougougou/ Blockauss		
TMAF	Charge CFU/mL	8.52x10 ⁴ ± 0.08	2.78x10 ⁵ ± 0.14	3.99x10 ⁵ ± 0.34	10 ⁴	10 ⁵
	Quality	AMQ	UMQ	UMQ		
Yeasts and molds	Charge CFU/mL	0	9.09x10 ² ± 0.48	2.27x10 ² ± 1.08	10 ³	10 ⁴
	Quality	SMQ	SMQ	SMQ		
Total coliforms	Charge CFU/mL	5.81x10 ⁴ ± 1.18	6.55x10 ⁴ ± 0.78	6.78x10 ⁴ ± 0.57	10 ³	10 ⁴
	Quality	UMQ	UMQ	UMQ		
Escherichia coli	Charge CFU/mL	0	0	0	1	10
	Quality	SMQ	SMQ	SMQ		
Staphylococcus sp.	Charge CFU/mL	6.26x10 ³ ± 0.35	1.88x10 ⁴ ± 0.07	2.95x10 ⁴ ± 0.04	10	10 ²
	Quality	UMQ	UMQ	UMQ		
Salmonella sp.	Absent in 25 mL	Absent	Present	Present	Absent in 25 mL	
	Quality	SMQ	UMQ	UMQ		

Table 6. Contamination level of Anagobaka porridge

Germs studied	Samples of Anagobaka porridge taken from quarter (n=5 per quarter)			
		Kôkô/Trade	Microbiological standards	
TMAF	Charge CFU/mL	$3.43 \times 10^5 \pm 0.22$	10^4	10^5
	Quality	UMQ		
Yeasts and molds	Charge CFU/mL	$2.41 \times 10^4 \pm 0.13$	10^3	10^4
	Quality	UMQ		
Total coliforms	Charge CFU/mL	$4.08 \times 10^4 \pm 0.48$	10^3	10^4
	Quality	UMQ		
<i>Escherichia coli</i>	Charge CFU/mL	$3.91 \times 10^3 \pm 0.09$	1	10
	Quality	UMQ		
<i>Staphylococcus sp.</i>	Charge CFU/mL	$5.34 \times 10^4 \pm 0.10$	10	10^2
	Quality	UMQ		
<i>Salmonella sp.</i>	Absent in 25 mL	Present	Absent in 25 mL	
	Quality	UMQ		

Table 7. Contamination level of Focolari flour

Germs studied	Flour samples taken at CMS Focolari (n=5 per type of flour)				
		FM5B	FMS	FMSC	Microbiological standards
TMAF	Charge CFU/mL	$1.90 \times 10^4 \pm 0.13$	$7.83 \times 10^4 \pm 0.38$	$1.79 \times 10^5 \pm 1.24$	10^4 10^5
	Quality	AMQ	AMQ	UMQ	
Yeasts and molds	Charge CFU/mL	$9.09 \times 10^2 \pm 0.59$	$1.65 \times 10^3 \pm 0.78$	$1.72 \times 10^4 \pm 1.04$	10^3 10^4
	Quality	SMQ	SMQ	UMQ	
Total coliforms	Charge CFU/mL	$4.97 \times 10^3 \pm 0.04$	$3.62 \times 10^3 \pm 1.12$	$5.58 \times 10^3 \pm 0.07$	10^3 10^4
	Quality	AMQ	AMQ	AMQ	
<i>Escherichia coli</i>	Charge CFU/mL	0	0	0	1 10
	Quality	SMQ	SMQ	SMQ	
<i>Staphylococcus sp.</i>	Charge CFU/mL	$1.42 \times 10^2 \pm 0.31$	$1.04 \times 10^2 \pm 0.14$	$2.82 \times 10^3 \pm 0.59$	10 10^2
	Quality	UMQ	UMQ	UMQ	
<i>Salmonella sp.</i>	Absent in 25 mL	Absent	Absent	Absent	Absent in 25 mL
	Quality	SMQ	SMQ	SMQ	

each type of germ, thus testifying to a very high level of contamination. Thus, for the total aerobic mesophilic flora the average load is 3.43×10^5 CFU/mL, yeasts and molds have a charge of 2.41×10^4 CFU/mL, total coliforms showed a load of 4.08×10^4 CFU/mL, *Escherichia coli* revealed an average load of 3.91×10^3 CFU/mL, *Staphylococcus aureus* revealed an average load of 5.34×10^4 CFU/mL and *Salmonella* was found in the samples.

3.2.2 Total mesophilic aerobic flora (TMAF)

The microbiological quality analysis revealed an average load (N) of the TMAF of acceptable

microbiological quality (AMQ) for the millet porridge. These values range from 4.97×10^4 to 8.58×10^4 CFU/mL. Concerning the Kokobaka porridge, the results revealed a AMQ of 8.52×10^4 CFU/mL for the Camp Sea/Cafop sector and an unsatisfactory microbiological quality (UMQ) for the Kôkô/trade and Dioulabougou/Blockauss sectors with average loads of 2.78×10^5 and 3.99×10^5 CFU/mL, respectively. Focolari flours showed a satisfactory microbiological quality (QMS) with a value of 1.90×10^4 CFU/g. At Camp Sea/Cafop, the analysis gave an acceptable microbiological quality (AMQ) of a load of 7.83×10^4 CFU/mL. The Kôkô/Trade sector and an unsatisfactory microbiological quality (UMQ)

with a load of 1.79×10^5 for maize-soya-cocoa flour (FMSC).

3.2.3 Yeasts and molds

The results of the microbiological analysis show a satisfactory microbiological quality (QMS) for all the porridges of millet, Kokobaka and Focolari flours. The millet porridges analyzed have an average load of 0 CFU/mL in all sectors, that of the Kokobaka porridges varies from 0 to 2.27×10^2 CFU/mL. Focolari flours have a load of 90.9 CFU/g; 1.65×10^3 CFU/g and 30.3 CFU/g respectively for FMS, FM5B and FMSC.

3.2.4 Total coliforms

For total coliforms, the contamination level of the millet porridge is of acceptable microbiological quality with a load of 4.14×10^3 CFU/mL. For the Camp Sea/Cafop sector, the microbiological quality is unsatisfactory with an average microbial load of 4.58×10^4 CFU/mL and 3.58×10^4 CFU/mL respectively in the Kôkô/ Trade and Dioulabougou/ Blockauss sectors. Kokobaka porridge, on the other hand, testifies to a microbiological quality that is unsatisfactory in all sectors with a load of 5.81×10^4 ; 6.55×10^4 CFU/mL and 6.78×10^4 CFU/mL at Camp Sea/Cafop, Kôkô/Trade and Dioulabougou/ Blockauss. Conversely, all the flours from Focolari have an acceptable microbiological quality with an average load equal to 4.79×10^3 CFU/g, 3.52×10^3 CFU/g and 5.58×10^3 CFU/g respectively for FMS, FM5B and FMSC.

3.2.5 *Escherichia coli*

Concerning *E. coli*, the results of microbiological analysis show a satisfactory microbiological quality (QMS) for all the porridges of millet, Kokobaka and Focolari flours. The average load is 0 CFU/mL in all sectors for porridge and also 0 CFU/g for all flours.

3.2.6 *Staphylococcus aureus*

The results of the microbiological analysis show a satisfactory microbiological quality (SMQ) for all the porridges of millet, Kokobaka and Focolari flour concerning *Staphylococcus aureus*. The millet porridges analyzed have an average load which varies from 2.21×10^3 CFU/mL (Camp Sea/Cafop) at 5.33×10^3 CFU/mL (Dioulabougou/ Blockauss). The Kokobaka porridges revealed an average load ranging from 6.26×10^3 CFU/mL at

2.95×10^3 CFU/mL. The load of the Focolari flours has a load of 1.42×10^3 CFU/g ; 2.82×10^3 CFU/g and 3.03×10^1 CFU/g respectively for FMS, FM5B and FMSC.

3.2.7 *Salmonella*

Concerning *Salmonella*, results of microbiological analysis show a satisfactory microbiological quality for all the millet and Kokobaka porridges and the Focolari flours analysed. An absence of salmonella was found in all samples.

3.3 Results of Physicochemical Analyzes

3.3.1 Macronutrient content

The carbohydrate content of all the samples analyzed respects the recommended standard which is 68 g/100g Dry Weight (DW). Regarding the lipids content, all the flours from the Focolari medical-social center and flour of Kokobaka porridge (FKP) respect the standard set, unlike the flours for porridge of millet and Anagobaka which do not respect this standard with respective values averages of 7.80 ± 0.08 g/100g DW and 6.93 ± 0.09 g/100g DW which is below 8 g/100g DW. The moisture content of all the flours from the Focolari Center analyzed do not meet the set standard, which is the coating of the flours of millet porridge (8.74 ± 0.21 %), Kokobaka (6.55 ± 0.17 %) and Anagobaka (13.15 ± 0.5 0%), the norm being 5%. In terms of ash content, all the samples do not meet the standard set, such as the FM5B flour from Focolari, which has a value of 3.21 ± 0.24 % which is much higher than the 2.9 % standard (Table 8).

3.3.2 Micronutrient content

The phosphorus content of Focolari flours varies from 194.05 mg/100g DW (FM5B) to 503.45 mg/100g DW (FMS). Regarding porridge flour, this content is 178.93 mg/100g, 505.57 mg/100g and 271.77 mg/100g respectively for flour, millet porridge (FBM), Kokobaka (FKP) and Anagobaka (FAP). The potassium content of flours from Focolari ranges from 295.74 mg/100g DW (FMSC) to 605.80 mg/100g DW (FMS). Potassium content of porridge flours vary from 23.06 mg/100g DW to 49.23 mg/100g DW. The calcium contents vary from 192.71 mg/100g DW to 441.70 mg/100g DW and from 21.15 mg/100g DW to 35.52 mg/100g DW for the flours from the medical-social center and those from the

Table 8. Macronutrients parameters

Physico-chemical parameters (g/100g DW)				
Samples	Carbohydrates	Lipids	Moisture (%)	Ash
FMS	75.86 ± 0.06	8.10 ± 0.11	4.90 ± 0.08	1.22 ± 0.19
FMSC	70.17 ± 0.15	9.50 ± 0.50	4.86 ± 0.05	2.50 ± 0.09
FM5B	77.45 ± 0.11	10.40 ± 0.31	4.59 ± 0.15	3.21 ± 0.24
FMP	80.14 ± 0.31	7.80 ± 0.08	8.74 ± 0.21	1.75 ± 0.10
FKP	85.08 ± 0.17	8.04 ± 0.05	6.55 ± 0.17	1.05 ± 0.05
FAP	79.25 ± 0.12	6.93 ± 0.09	13.15 ± 0.50	1.50 ± 0.13
FAO/WHO (2006) Standard	68	8	5	2.9

FKP: Flour of Kokobaka porridge; FMP: Flour of Millet porridge; FAP: Flour of Anagobaka porridge; FMS: Infant maize and soy flour; FMSC: Flour made from maize, sesame and cocoa; FM5B: Infant flour made from maize, sesame, soy, millet, rice, cocoa and banana.

Table 9. Micronutrients of the samples studied

Micronutrients in flours (mg/100g DW)						
Samples	P	K	Ca	Fe	Zn	Cu
FMS	503.45±0.01	305.8±0.01	192.91±0.01	52.04±0.03	2.56±0.02	13.43±0.01
FMSC	333.08±0.02	295.74±0.01	323.61±0.01	30.64±0.01	2.2±0.01	9.3±0.01
FM5B	505.57±0.01	413.14±0.01	441.7±0.01	35.02±0.01	1.76±0.02	7.8±0.02
FBM	178.93±0.01	49.23±0.01	35.52±0.04	64.34±0.02	2.92±0.01	1.32±0.01
FBK	194.07±0.01	34.86±0.02	21.15±0.02	63.56±0.01	1.26±0.02	1.31±0.01
FBA	271.77±0.01	23.06±0.01	30.01±0.01	48.24±0.01	2.6±0.01	0.38±0.02
FAO/WHO (2006) Standard	281.2	408.7	341.2	8.5	3.7	0.1

FKP: Flour of Kokobaka porridge; FMP: Flour of Millet porridge; FAP: Flour of Anagobaka porridge; FMS: Infant maize and soy flour; FMSC: Flour made from maize, sesame and cocoa; FM5B: Infant flour made from maize, sesame, soy, millet, rice, cocoa and banana.

porridges respectively. The iron content varies from 30.64 mg/100g DW to 52.04 mg/100g DW for the Focolari flours and from 48.24 mg/100g DW to 64.34 mg/100g DW for the vendors' porridge flours. The zinc contents varies from 1.76 mg/100g DW to 2.56 mg/100g DW for the Focolari flours and from 1.26 mg/100g DW to 2.92 mg/100g DW for the porridge flours. The copper content varies from 7.80 mg/100g DW to 13.43 mg/100g DW and from 0.38 mg/100g DW to 1.31 mg/100g DW respectively for Focolari flours and porridge flours. The results of micronutrient contents of the various flours analyzed show that, with the exception of iron and copper which comply with the standard, the mineral contents of the flours used as a complementary food are generally lower than the recommended standards (Table 9).

4. DISCUSSION

At the end of the results of the survey carried out among mothers and vendors, it appears that 80 % of mothers do not know the concept of complementary food. Indeed, more than half of

these mothers (62.5 %) self-advise on complementary food against only 12.9 % of them who have recourse to a health agent. The role of the health worker is indeed to explain the importance of the complementary food and therefore to provide some nutritional education to nursing mothers. These results coincide with those of Azagoh et al. [18] who found similar results where 74 % of mothers were unaware of the notion of complementary feeding during their study on the knowledge and practices of mothers of children aged 6 to 18 months relating to weaning. However, these results differ from those of Diallo [19] who worked on knowledge of mothers' attitudes and practices on the feeding of children from 0 to 23 months in the health district of Niafunké in Mali. Indeed, 84.8% of the women surveyed in this health district do not know the concept of complementary food.

In addition, the food most given to children is Anagobaka porridge (32.2 %), followed by millet porridge (24.4 %). In general, the most used complementary foods are foods based on cereals and legumes. Indeed, these cereals and

legumes are locally available and therefore easily accessible to nannies who use them to make meals for children. This assertion is supported by Koné et al. [6] who worked on the formulation and physicochemical characterization of infant flour composed of: dehydrated attiéké-cashew kernel. Similarly, Mühlemann [9] confirmed the use of porridges made from cereals, tubers and roots during his study on complementary feeding of children during their first two years of life.

In addition, the sellers have no knowledge of the proportion of the components and the nutritional intake of the complementary food they sell. This observation is the same at the level of the Focolari medico-social center, where the flour samples are collected. This is dangerous for consumers as fragile and vulnerable as children aged between 6 and 24 months. Aissi et al. [20] found this same fact in their study. This is why the Codex Alimentarius requires disclosure of food content in food labeling [21].

These complementary foods are introduced into the life of the child according to the appreciation of the mother. Indeed, 61.1 % said they had given the child the complementary food before the age of six (06) months. Against only 27.8 % of mothers who claim to start this diet from six (06) months. These results are not consistent with those promoted by the WHO [5] which recommends that complementary foods should only be introduced into the diet of young children from the age of six months. The ignorance of the notion of complementary feeding, the advisers (the mothers themselves) could justify this situation. The work carried out by Azagoh et al. [18] had already mentioned that the majority of mothers did not respect the WHO recommendations concerning the start of the introduction of complementary foods in children under 24 months.

Among the nannies (mothers) who use complementary food, only 11.8% buy infant food in a health center. This low proportion of purchases in health centers translates into the fact that mothers do not take their child's feeding advice from health workers [19].

In terms of hygiene, 54.8% of vendors operate in an inappropriate environment and place of sale. On the other hand, 41.9% of those sell in an acceptable sanitary environment. According to Azagoh et al. [18], the level of education has a correlation with the knowledge and practice of hygiene. The level of education of porridge

vendors in the neighborhoods selected for this study is low. Indeed, 32.3% of them have not been to school. This situation could justify the lack of good hygiene practices observed by these porridge sellers.

The results of the microbiological analyzes indicate a high level of contamination of the porridges and supplementary flours. The porridge and flour samples from the Focolari medical-social center in the 3 sectors sampled show an acceptable microbiological quality for the search for total aerobic mesophilic flora (TMAF). Except for the samples from the Kôkô/Trade sector (Kokobaka porridge and Anagobaka porridge), the Dioulabougougou/Blockauss sector (Kokobaka porridge) and the FMSC flour from the Focolari medico-social center in Man. Indeed, the load of the total aerobic mesophilic flora (TMAF) is below or within the upper limit of the standard prescribed by WHO (10^4 - 10^5 CFU/ml). It is important to specify that this flora is an indicator of the general level of hygiene of a product, it provides information on factors such as the environment, cross-contamination during handling, the environment, packaging and product conservation [22]. This allows us to say that the vendors in the Kôkô/Trade sector (Kokobaka porridge and Anagobaka porridge) and Dioulabougougou/Blockauss (Kokobaka porridge) and the FMSC flour from the Focolari center are faced with food safety problems. However, these germs do not have a great impact on the health of the consumer, on the other hand, they cause significant economic losses due to the deterioration of the products. In terms of porridge, the results obtained differ from that of Dossa [22] who found a low presence of total flora in his sample of millet porridge (1.6×10^3 CFU/mL).

Flours from millet, Kokobaka and Anagobaka porridges and those from the Focolari medical-social center showed total coliform levels above the recommended standard in food (10^3 - 10^4 CFU/mL).

The presence of total coliforms could be translated by faecal contamination during the production process, attributable to the lack of hygiene of the saleswomen. The acceptability shows that a certain vigilance is observed during the production of porridge and flour produced by some vendors and producers. These results corroborate those of Noutais [23] who found the presence of coliforms in a sample (22×10^3

CFU/mL) from his study on the evaluation of the microbiological quality of two types of fermented porridge made from sorghum. According to Houssou et al. [24], milling would be a critical step, it depends on the sunshine and the level of sanitation of the places and influences the microbiological quality of the flours. The results.

Staphylococcus sp. is present in all the samples with an unsatisfactory microbiological quality. The results obtained could be explained by handling during the sale and the absence of wearing professional sales clothing. The presence of these germs could be due to a lack of hygiene (direct contact of the saleswomen with the product) during production and marketing, *staphylococcus sp.* is a vector of oral and cutaneous contamination. These bacteria are the cause of food poisoning, despite being part of the commensal flora of humans. The results obtained are similar to those of N'Goran-Aw et al. [8] and Nouais [23].

Escherichia coli and *Salmonella sp.* were absent in all the samples of millet porridge, Kokobaka and infant flours (FMS, FM5B and FMSC) unlike the Anagobaka porridge which revealed an unsatisfactory microbiological quality for *E. coli* and the presence of *Salmonella*. These porridges therefore demonstrate satisfactory microbiological quality. *Escherichia coli* is called "faecal contamination control" germ, it indicates that there is dirty handling and is characteristic of contamination from faeces. *Salmonella sp.* is itself a pathogenic germ, its presence concludes that the product is dangerous for consumption and would be due to poor processing conditions (contaminated environment or insufficient cooking) [25].

Concerning the macronutrient contents and the physicochemical parameters, the total carbohydrate content of all the flours analyzed is very high and does not respect the standard set (68 g/100g DW) by FAO/WHO [4]. The lipid content of all the samples practically meets the standard set by FAO/WHO [4] except that of the Anagobaka porridge flours. The lipids content of this flour is certainly lower than the norm but differs from that obtained by Anigo et al. [26] in their work on complementary food gruels formulated from malted cereals, soybeans and groundnut in North-western Nigeria. Lipids play an important role in satiety and weight gain. A food low in fat and energy promotes weight loss [25]. The moisture content of all flours is lower than the norm except those of millet porridge

(8.74 %), Kokobaka porridge (6.55 %) and Anagobaka porridge (13.15 %). The high flour moisture content of the Anagobaka porridge far above the recommended standard could be due to the long soaking time of the maize kernels and less drying of the flour. The low moisture content of flours increases their shelf life and, from a microbiological point of view, limits the development of microorganisms, with the exception of molds [4].

In terms of the micronutrient contents of the flours analyzed, the phosphorus contents of all the flours of the Focolari medical-social center are higher than the recommended standard, unlike the flours of the porridge where the phosphorus contents are below the standard prescribed by FAO/WHO [4]. These results differ from those of Kpan et al. [10] who worked on the nutritional quality of Anagobaka weaning flours marketed in the markets of four municipalities in the district of Abidjan, for Focolare flours and are identical for porridge flours. The potassium and calcium contents of all the flours analyzed do not meet the recommended standard except infant flour made from maize, sesame, oy, millet, rice, cocoa and banana (FM5B). This result could be due to the diversity of raw materials used in the formulation of this flour unlike other flours. The work carried out by Kpan et al. [10] presented potassium and calcium levels below the norm in their study. Potassium is necessary for the regulation of cell water balance, the utilization of carbohydrates and the construction of proteins. Calcium is the mineral par excellence because it ensures the rigidity of the bones and promotes the growth of children [27]. The iron and copper contents of all the types of flour analyzed meet the set standard. These results differ from those of Badham et al. [28] (for iron) and are identical to those of Kpan et al. [10] (for copper). Iron is involved in the constitution of enzymes and the transport of oxygen. An iron deficiency causes a decrease in physical abilities and a drop in resistance to infections. The most vulnerable people are women and children under two [28]. The zinc content of all types of flour analyzed is below the set standard. Zinc deficiency is one of the main causes of stunted growth in children.

5. CONCLUSION

This study carried out in Man (Côte d'Ivoire) revealed that the nannies of this locality give food other than their breast milk to their child before the age of six (06) months. The locally produced

complementary food given the most to children is Anagobaka porridge, followed by that of millet, then porridge made from flour from the Focolari medical-social center and Kokobaka porridge. However, the microbiological analysis of these food supplements produced in Man, revealed a potential contamination of these by various germs of alterations and pathogens. Above all, the contamination rate of Anagobaka porridge is very worrying especially with the presence of *Salmonella* and the high rate of *Escherichia coli*, *Staphylococcus sp* unlike other porridges (millet, kokobaka) and flours (FMS, FMSC, FM5B). The analysis of the physicochemical composition of the flours revealed that these porridges and flours from the Focolari medical-social center contain high levels of carbohydrates, iron and copper which do not meet the standards prescribed for infant food. However, the lipid contents of the samples comply with the FAO/WHO recommendations for a complementary food except for the Anagobaka porridge sample. However, the zinc content is below this standard in all the samples tested. However, zinc deficiency is one of the main causes of stunted growth in children. The results of this study have shed light on the existence of a potential health risk and possible nutritional deficiencies in infants and young children who consume these locally produced complementary foods in Man (Côte d'Ivoire).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. WHO, Nutrition for infants and young children: sixty-third world health assembly. World Health Organization, Washington, D.C. 2017;9.
2. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, Onis M, Ezzati M, Grantham MS, Katz J, Martorell R, Uauy R. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013;382(9890): 427-451.
3. *Lancet*, Maternal and Child Nutrition. Executive Summary of The *Lancet* Maternal and Child Nutrition Series. 2013;1-12.
4. FAO/WHO, Joint FAO/WHO Food Standards Programme. Report of the twenty -seventh session of the codex committee on nutrition and foods for special diets. ALINOM. 2006;105.
5. WHO, UNICEF. Global Strategy for Infant and Young Child Feeding. 2002;37.
6. Koné S, Soro D, Koffi EK. Formulation and physicochemical characterization of infant flour made up of: dehydrated attiéké-cashew kernel. *International Journal of Innovation and Applied Studies*. 2019;25(2):700 – 708.
7. Kagambèga B, Hama C, François T, Adama S, Cheikna Z, Yves T, Aly SS. Traditional fermented cereal-based porridges in Burkina Faso: diversity, technologies of production and microorganisms with associated probiotic potential. *Science Technology Review, Synthesis*. 2019;25(2):12-24.
8. N'Goran-AW EBZ, Doudjo S, Sadat A, David AK, Emmanuel AN. Assessment of the physico-chemical and microbiological characteristics of a traditional donut made from fermented millet (gnomy) marketed in the city of Yamoussoukro (Côte D'Ivoire). *European Scientific Journal*. 2017;13(9):227-241.
9. Mühlemann P. Iron nutrition during the first two years of life. A study on weaning practices in the Ivory Coast; 1998.
10. Kpan SE, Yapo CA, Boga GL, Diane K, Djaman AJ, N'guessan JD. Nutritional quality of Anagobaka weaning flours marketed in the markets of four communes in the District of Abidjan. *Africa Science*. 2018;14(6):413 – 427.
11. MICS, Multiple Indicator Cluster Survey – Côte d'Ivoire. 2016;1-442. [Online] Available:<https://mics-surveysprod.s3.amazonaws.com> (October 18, 2022).
12. FAO. Conducting small nutrition surveys field manual, Issue 5 of Nutrition and Agriculture, Rome. 1992;5:1-180.
13. Aboua F, Nemlin J, Kossa A, Kamenan A. Traditional processing of some cereals grown in Côte d'Ivoire. *Cereals in hot regions*. AUPELF-UREF, Eds John Libbey Eurotext, Paris. 1989;223-229.
14. AOAC. Official methods of analysis. Association of Official Analytical Chemists Ed., Washington DC. 1990;1-684.
15. Folch J, Lees M, Stanley GH. A simple method for the isolation and purification of total lipids from animal tissues. *Journal of Biological Chemistry*. 1957;226:497–509.
16. Munson LS, Walker PH. The unification of reducing sugar methods. *American*

- Chemistry Society Journal. 1906;28: 663-686.
17. Thirion-Merle V. X-ray fluorescence spectrometry. 2014;1-7. [Online] Available:<https://hal.archives-ouvertes.fr> (18 October 2022).
 18. Azagoh KR, Enoh J, Niangue B, Cissé L, Oulai S, Andoh J. Knowledge and practices of mothers of children from 6 to 18 months relating to weaning management: case of the general hospital of Marcory. Mali Medical. 2013;XXVIII(4): 1-4.
 19. Diallo AM. Knowledge, attitudes and practices of mothers on the diet of children from 0 to 23 months and their nutritional status in pediatrics / Ureni Ducsfref de Niafunke from December 2018 to February 2019, Thesis of Medicine in Odontostomatology, University of Sciences, Techniques and Technologies of Bamako, Mali. 2020;1-109.
 20. Aissi AK, Koungblenou SD, Dougnon V, Klotoe JR, Bankole H, Deguenon Y, Degbey C, Montcho S, Fanou B, Fah L, Edoh PA, Loko F. Evaluation of the sanitary quality of *Moringa oleifera* Lam leaf powders marketed for the benefit of People Living with HIV in Cotonou (Benin). International Journal of Biological and Chemical Sciences. 2013;7(4):1461-1473.
 21. FAO/WHO. Food labelling, full codex alimentarius texts. Joint FAO/WHO Food Standards Programme; 2001.
 22. Dossa GSC. Microbiological evaluation of a local millet porridge "Bita" with a galactogenic effect marketed by the company "mille et une porridges" in Cotonou". Professional degree in food technology engineering, University of Abomey-Calavi, Benin. 2019;1-59.
 23. Noutais L. Evaluation of the microbiological quality of two types of fermented sorghum-based porridges, marketed by the company "Mille et une porridge" in Co tone". Professional degree in food technology engineering, University of Abomey-Calavi, Benin. 2018;1-46.
 24. Houssou PAF, Ahoyo ARN, Metohou ER, Dansou V, Djivoh H, Hotegni AB, Mensah GA. Evaluation of the quality of yêkè-yêkè (corn couscous) and gambarilifin (refined flour of maize) during storage. Ivorian Science Technology Review. 2016;27: 136-150.
 25. Tabo DA. Study of the contribution of avian salmonella to human salmonellosis in Chad: Case of the capital city, N'Djamena. Doctoral thesis at the Institute of Life and Environmental Sciences and Industries (AgroParisTech), University of Paris Tech. 2013;1-152.
 26. Anigo KM, Ameh DA, Ibrahim S, Danbauchi SS. Nutrient composition of complementary food gruels formulated from malted cereals, soybeans and groundnut for use in North-western Nigeria. African Journal of Food Science. 2010;4:65-72.
 27. Charturvedi VC, Shrivastava R, Upreti RK. Viral infections and trace element: A complex trace element. Current Science. 2004;87:1536 – 1554.
 28. Badham J, Zimmermann MB, Kramer K. The guide to nutritional anemia. Switzerland: Sight and Life; 2007. [Online] Available: <http://www.sightandlife.org/> (18 October 2022)

© 2022 Carole et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/93975>