



Assessing the Impact of Gas Flaring on Tea Plants in the South of Kothaloni OCS in the District of Dibrugarh, Assam, India

Ripsita Phukan ^{a*}, Indra Prasad Sahewalla ^a and Diganta Kumar Bora ^a

^a *Department of Tea Husbandry and Technology, Assam Agricultural University, Jorhat, Assam, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i1130943

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85416>

Original Research Article

Received 22 January 2022

Accepted 26 March 2022

Published 29 March 2022

ABSTRACT

Effect of gas flaring on physiology and growth of tea plants was done by using Randomized Complete Block Design (RCBD) and analysis was done accommodating two seasons and five levels of distances during September 2019-March 2020. Plant samples were collected from (40-50) metres, (50-60) metres, (60-70) metres, (70-80) metres and control site (150-160) metres away from the tea garden situated south of kothaloni OCS in rainy and autumn seasons. The experimental plot was laid out at a distance of 40 metres away from the flare pit. The mean value of plant parameters namely relative turgidity, tea leaf area, specific leaf weight, stomatal count, caffeine content, chlorophyll content, polyphenol content and plucking point density were decreased near the gas flaring while mean value of water saturation deficit of tea leaves recorded highest at distances closest to the flaring site.

Keywords: Gas flaring; seasons; distance.

1. INTRODUCTION

Many oil drilling sites of upper Assam are located near the tea estates. The excess or unwanted gases which are produced during different operations such as crude oil extraction get burnt through a device. This process can be called as

"gas flaring", and the resulting gas is called "gas flare" [1,2] and is generally used as a safety measure to release pressure [3].

Gas flare consists of a mixture of different gases and its composition greatly depends on the gas source that is entering to the system of flare.

*Corresponding author: E-mail: ripsitaphukan@gmail.com;

During oil-gas production, associated gases that are released mainly contain natural gas and natural gas is consisted of more than 90% methane (CH₄) with ethane, a small amount of other hydrocarbons and inert gases such as N₂ and CO₂ [1]. Gas flare from refineries and other different operations in general contain a mixture of hydrocarbons, carbon dioxide (CO₂) and sometimes hydrogen [1]. Being among the Green House Gases, CO₂ and methane (CH₄) contribute to the increase of the temperature of the atmosphere when directly released into the air [1]. So, gas flare significantly impacts the microclimate in vicinity of the flare sites [4]. This gas also contributes to the acid rain, with its effects on vegetation are more severe near the flare site [5]. The acid rain causes vegetation loss too [6].

India is one of the leading countries in the world with respect to global tea consumption, production and exports. The state of Assam in India lying on either side of the Brahmaputra river is one of the most prolific tea producing regions and also enriched with natural resources like natural gas, oil and coal. . On the whole, the crude oil is one of the valuable natural resources of Dibrugarh district in the state of Assam (India), where the petroleum industry plays an important role in the process of its industrialization. This district is also the place of intense practice of the tea [*Camellia sinensis* (L) O. Kuntze] culture. This is also true for Kothaloni in Dibrugarh district , particularly South of Kothaloni Oil Collection Station (OCS). However, there is any information concerning to the effects of gas flaring on tea plants in this area. Knowing the great socio-economic role of tea in this area, it is important to carry out investigations on this fact. It is the purpose of this study.

2. MATERIALS AND METHODS

2.1 Characteristics of the Study Area

The experimental site located in South of Kothaloni OCS in No. 1 Naharani, Dibrugarh district of Assam, India is situated at 27°37'97"N latitude and 95°09'30"E longitude and at an elevation of 108 m above mean sea level.

2.2 Experimental Techniques

2.2.1 Sampling sites

The study was done during September 2019-March 2020. Experimental Design selected was Randomized Complete Block Design accomodating five distances namely D1(40-50

metres, D2(50-60) metres, D3(60-70) metres, D4(70-80) metres and control site DC(150-160) metres away from the gas flaring site and two seasons namely rainy and autumn seasons. A pond was there in between the flare site and the selected tea garden and so the experimental plot was laid out at 40 metres away from the flare pit. The flaring site was surrounded by a concrete wall. Rainy and autumn seasons were selected to know its seasonal variations in different plant parameters.

2.2.2 Measurement of physiological parameters

2.2.2.1 Relative turgidity (RT) and Water saturation Deficit (WSD)

RT was obtained according Weatherley method [7]. Its expression is given by:

$$RT(\%) = \frac{\text{Average fresh weight (mg)} - \text{Average dry weight (mg)}}{\text{Average saturation weight (mg)} - \text{Average dry weight (mg)}} \times 100$$

As concerned to WSD, it was obtained following Weatherley method [7] and its expression is:

$$WSD(\%) = \frac{\text{Average saturation weight (mg)} - \text{Average fresh weight (mg)}}{\text{Average saturation weight (mg)} - \text{Average dry weight (mg)}} \times 100$$

2.2.2.2 Specific leaf weight

Fifty leaves without the petiole were selected from each planting material and their individual leaf area was measured. Specific leaf weight of tea leaves can be calculated by using the following formula:

$$SLW(g/cm^2) = \frac{\text{Dry weight of leaf (mg)}}{\text{Total leaf area (cm}^2\text{)}}$$

2.2.2.3 Stomatal count

The estimation of the stomatal count was determined by leaf impression method [8]

2.2.2.4 Leaf area of tea leaves(cm²)

Individual leaf was taken and tracing was done over graph paper and the covered grids by the leaf are counted to give the area.and done using millimeter graph paper method [9]. It is expressed in cm².

2.2.2.5 Chlorophyll content

The chlorophyll extraction was done using methanol given by Taylor [10]. Buds and two

leaves weighing 0.5g were taken and chopped into pieces. A liquid mixture was made by adding 25 ml of methanol slowly to the chopped pieces placed in mortar and pestle. In 25 ml conical flask, 2ml of supernatant was transferred and conical flask was covered with carbon paper which was kept separately in a dark place for about 30 minutes. The mixture was then centrifuged at 5000 rpm for 5 minutes. In 2 ml of the mixture, 8 ml methanol was added after centrifuge to make a volume of 10 ml. Then spectrophotometric observation of diluted solutions was recorded in different wavelengths namely 653 nm for chlorophyll-a and 666 nm for chlorophyll-b. It was expressed as mg/g.

The sample content in chlorophyll a was obtained following this expression:

$$\text{Chl-a} = 15.65A_{666} - 7.34A_{653}$$

That in chlorophyll b was by

$$\text{Chl-b} = 27.05A_{653} - 11.21A_{666}$$

A_{666} ; A_{653} the absorbance of the solution obtained after treatment at 653 and 666 nm respectively.

2.2.2.6 Caffeine content

Caffeine estimation was done by boiling 200 ml of water containing 20 g of two leaves and a bud for 30 minutes in a 400 ml beaker and should be filtered while hot. The filtrate was collected in another clean beaker and of Lead acetate of 10% aqueous solution was added in the filtrate that was taken in a clean beaker followed by centrifuged at 5000 rpm for 5 minutes. The volume of the supernatant was reduced to 25 ml by boiling and cooled it at room temperature. The mixture was extracted with 25 ml chloroform in a separating funnel after cooling the reaction. The chloroform was distilled off from the chloroform extract on a water bath using a water condenser. By using a spatula, the dried residue was scrapped from the flask and weighed was taken. Weight of fresh tea leaves and weight of crude caffeine in gram gives the amount of caffeine. The total caffeine in tea leaves was expressed in percentage (%).

2.2.2.7 Total polyphenol content

The shoots were separated into bud, first leaf and second leaf, oven-dried and then crushed to make powder and polyphenol content was estimated with Folin-Ciocalteu reagent, based on

the reaction between phenols and phosphomolybdate that results in the blue complex formation [11]. It was expressed in percentage (%).

2.2.2.8 Plucking point density

The method used for counting plucking points was same as the method described by Barua and Dutta [12]. A light bamboo grid of 10 cm square sides was placed on the top of the bush and in the randomly selected plant, the number of plucking points at the the end of the season was collected.

3. RESULTS AND DISCUSSIONS

3.1 Relative Turgidity

Relative turgidity of tea leaves had shown significant differences among distances and seasons. Relative turgidity was found to be lower near the gas flaring site which increased with distances away from the gas flare (Table 1). This might be due to high temperature stress that decreased fresh and dry weight of leaves of tea plant which reduced its relative turgidity. This was agreed with the findings of Naz [13], who reported that relative turgidity of potato plants were successively reduced by increased heat stress. In this study, relative turgidity of tea leaves had shown decreased mean value in autumn season as compared to rainy season. This might be due to increased water stress in the season of autumn which led to reduction in relative turgidity of tea leaves. These results corroborates the findings of Waheed [14] who reported that relative turgidity of tea leaves were successively decreased during drought stress.

3.2 Water Saturation Deficit

Water saturation deficit was significantly varied among distances. Among the distances, D2, D3, D4 and DC were significant to D1. There was no significant variations observed in rest of the distances. Higher water saturation deficit near the gas flaring site as compared to far away distances (Table 2) might be due to high temperature stress that reduced tea plants' relative water content and it successively increased water saturation deficit of tea. The relative humidity around the gas flare was relatively reduced and might desiccate leaves of plants that are vulnerable to desiccation [15]. In this study, water saturation deficit of tea leaves

was increased in autumn season than rainy season. This might be due to water stress intensified by dry season that led to increase in water saturation deficit of tea leaves.

3.3 Number of Stomata

Number of stomata of tea leaves varied significantly among distances. Among the distances, variations were observed between all the distances except D3 and D4, low mean value were recorded near the flaring site (Table 3). Significant variation was observed among seasons and autumn season recorded the lowest mean value. This might be due to presence of pollutants in the surrounding environment of gas flare that changed biochemical composition along with morphology of plants leading to the nitrogen limitation for the plant. In a similar findings of Sarma [16], who reported that stomata of *Cassia tora* in Sivasagar district of Assam was reduced due to CO₂ stressed environment.

3.4 Leaf Area of Tea Leaves

It was observed that tea leaf area varied significantly among distances. Among the distances, D4, DC are significant to D1. Between rest of the distances no significant variations were observed. Leaf area of tea leaves were recorded to be lower near the gas flaring site (Table 4). This might be due to decreased stomatal conductance and reduced nitrogen content in plant growing under enriched air pollutants level for which the photosynthetic activities might be suppressed that led to reduction in leaf area of tea. The findings of Sarma [16] also reported reduction in leaf area of *Cassia tora* near the gas flaring site. However, tea leaf area had shown no significant differences in seasons.

3.5 Specific Weight of Tea Leaves

Specific leaf weight varied significantly among distances. Within the distances, significant variations were found between D1 and D3, D1 and D4, D1 and DC, D2 and D3, D2 and D4, D2 and DC. Specific weight of tea leaves were recorded to be lower near the gas flaring site (Table 5). This might be due to the reason that plants growing under enriched CO₂ stress might reduced leaf area of tea plants resulted in lowering its specific weight near gas flaring site.

The findings of Sarma [16] also reported reduction in leaf area of *Cassia tora* near the gas flaring site.

Seasons had shown significant variation where autumn season recorded the lowest mean value. This might be due to the seasonal variation of air pollutants and decreased photosynthetic activities during post monsoon period [16].

3.6 Chlorophyll Content

It was found that chlorophyll a content (mg/g) and chlorophyll b content (mg/g) were significantly varied along distances. Chlorophyll content a and b were recorded low in the vicinity of gas flaring site (Table 6). Combination of both drought and heat stress might be the reason of low chlorophyll content near the gas flaring site. Chlorophyll content might be reduced due to heat alone [17] and shortage of water might disrupt chloroplast structure which might led to destruction of Chlorophyll. This is in confirmation with the findings of Isichei and Sanford [15] who reported Chlorophyll content of *Eupatorium odoratum* was decreased in the distances nearer to gas flaring site.

Chlorophyll a and chlorophyll b contents had shown significant variation along seasons. Mean chlorophyll a and mean chlorophyll b contents were recorded highest in rainy season having 1.66 mg/g and 0.67 mg/g respectively.

3.7 Caffeine Content

Caffeine content in tea leaves had shown significant variations with distances. Distances nearer to gas flaring site had shown lowest value as compared to distances away from the gas flaring site (Table 7). This might be due to drought stress that inhibited the expression of genes related to caffeine biosynthesis and thereby reducing the accumulation of Caffeine in *Camellia sinensis*. This corroborates the findings of Song [18], Lee [19] and Zhang [20] who reported caffeine levels decreased with an increase in light.

3.8 Polyphenol Content

Significant variation was seen in polyphenol content (%) of tea leaves along distances and the significant variations were observed between all the distances except D2 and D3. Polyphenol content was observed to be lower in distances closest to the gas flaring site and also

observed lowest in autumn season as compared to rainy season (Table 8). This might be due to the alteration of biochemical constituents necessary for tea quality by drought stress.

Erturk [21] also reported significant decrease in total polyphenol content that reduced leaf quality of *Camellia sinensis* as affected by drought stress.

Table 1. Relative turgidity of tea leaves as affected by gas flaring

Seasons Distances (m)	Relative turgidity (%)			SE(m)	Significance
	Rainy season (S1)	Autumn season (S2)	Mean		
D1(40-50)	86.45	72.90	79.68		
D2(50-60)	90.32	76.04	83.18		
D3(60-70)	88.34	78.68	83.51		
D4(70-80)	89.36	78.09	83.72		
DC(150-160)	89.86	79.16	84.51		
Mean	88.86	76.97			
Factors	C.D.	SE(d)			
Distance(D)	2.48	1.17	0.83		S
Season(S)	1.57	0.74	0.52		S
Distance X Season(DXS)	N/A	1.659	1.17		NS

*S= Significant at 5% probability level; NS = Non Significant; N/A= Not Applicable

Table 2. Water saturation deficit of tea leaves as affected by gas flaring

Seasons Distances (m)	Water saturation deficit (%)			SE(m)	Significance
	Rainy season (S1)	AutumAutumn season(S2)	Mean		
D1(40-50)	13.58	27.45	20.51		
D2(50-60)	9.73	23.86	16.79		
D3(60-70)	11.30	21.54	16.42		
D4(70-80)	10.39	21.67	16.03		
DC(150-160)	11.00	21.01	16.01		
Mean	11.20	23.10			
Factors	C.D.	SE(d)			
Distance(D)	2.46	1.16	0.82		S
Season(S)	1.56	0.77	0.52		S
Distance X Season(DXS)	N/A	1.65	1.16		NS

*S= Significant at 5% probability level; NS = Non Significant; N/A= Not Applicable

Table 3. Stomatal count of tea leaves as affected by gas flaring

Seasons Distances (m)	Stomatal count (number/mm ²)				SE(m)	Significance
	Rain Rainy season(S1)	Autum Autumn season(S2)	Autumn	Mean		
D1(40-50)	20.00	19.33		19.66		
D2(50-60)	21.00	20.33		20.66		
D3(60-70)	21.66	21.33		21.50		
D4(70-80)	22.00	21.00		21.50		
DC(150-160)	23.00	22.66		22.83		
Mean	21.53	20.93				
Factors	C.D.	SE(d)				
Distance(D)	0.49	0.23		0.16		S
Season(S)	0.31	0.15		0.10		S
Distance X Season(DXS)	N/A	0.37		0.23		NS

*S= Significant at 5% probability level; NS = Non Significant; N/A= Not Applicable

Table 4. Leaf area of tea as affected by gas flaring

Seasons Distances(m)	Leaf area (cm ²)		Mean	Significance
	Rainy season (S ₁)	Autumn season (S ₂)		
D ₁ (40-50)	38.66	34.08	36.37	
D ₂ (50-60)	45.89	36.75	41.32	
D ₃ (60-70)	44.49	43.76	44.12	
D ₄ (70-80)	49.76	49.18	49.47	
D _C (150-160)	50.15	49.37	49.76	
Mean	45.79	42.62		
Factors	C.D.	SE(d)	SE(m)	
Distance(D)	8.78	4.15	2.93	S
Season(S)	N/A	2.62	1.85	NS
Distance X Season (DXS)	N/A	5.87	4.15	NS

*S= Significant at 5% probability level; NS = Non Significant ; N/A= Not Applicable

Table 5. Specific leaf weight of tea leaves as affected by gas flaring

Seasons Distances (m)	Specific leaf weight (g/cm ²)		Mean	Significance
	Rainy season (S ₁)	Autumn season (S ₂)		
D ₁ (40-50)	0.008	0.007	0.007	
D ₂ (50-60)	0.008	0.007	0.007	
D ₃ (60-70)	0.011	0.010	0.010	
D ₄ (70-80)	0.012	0.010	0.011	
D _C (150-160)	0.012	0.010	0.011	
Mean	0.010	0.009		
Factors	C.D.	SE(d)	SE(m)	
Distance(D)	0.002	0.001	0.001	S
Season(S)	0.001	0.001	0.000	S
Distance X Season (DXS)	N/A	0.001	0.001	NS

*S=Significant at 5% probability level; NS=Non Significant; N/A=Not Applicable

Table 6. Chlorophyll content of tea as affected by gas flaring

Seasons Distances (m)	Chlorophyll a content (mg/g)		Mean	Chlorophyll b content (mg/g)		Mean	
	Rainy season (S ₁)	Autumn season (S ₂)		Rainy season (S ₁)	Autumn season (S ₂)		
D ₁ (40-50)	1.00	0.93	0.96	0.43	0.41	0.42	
D ₂ (50-60)	1.19	1.15	1.17	0.53	0.51	0.52	
D ₃ (60-70)	1.58	1.39	1.48	0.67	0.57	0.62	
D ₄ (70-80)	2.18	2.12	2.15	0.86	0.73	0.79	
D _C (150-160)	2.35	2.28	2.31	0.88	0.82	0.85	
Mean	1.66	1.57		0.67	0.60		
Factors	C.D.	SE(d)	SE (m)	Significance	C.D.	SE(d)	SE (m) Significance
Distance	0.13	0.06	0.04	S	0.05	0.02	0.02 S
Season	0.08	0.04	0.03	S	0.03	0.02	0.01 S
Distance X Season (DXS)	N/A	0.09	0.06	NS	N/A	0.03	0.02 NS

*S= Significant at 5% probability level; NS = Non Significant ; N/A= Not Applicable

Table 7. Caffeine content of tea as affected by gas flaring

Seasons Distances (m)	Caffeine content (%)		Mean
	Rainy season (S ₁)	Autumn season (S ₂)	
D ₁ (40-50)	2.38	2.33	2.35
D ₂ (50-60)	2.46	2.44	2.45
D ₃ (60-70)	3.01	2.71	2.86
D ₄ (70-80)	3.29	3.19	3.24
D _C (150-160)	3.54	3.43	3.48
Mean	2.93	2.82	
Factors	C.D.	SE(d)	SE(m) Significance
Distance(D)	0.10	0.05	0.03 S
Season(S)	0.06	0.03	0.02 S
Distance X Season (DXS)	N/A	0.07	0.05 NS

*S= Significant at 5% probability level; NS = Non Significant ; N/A= Not Applicable

Table 8. Total polyphenol content of tea leaves as affected by gas flaring

Seasons Distances (m)	Total polyphenol content (%)		Mean	Significance
	Rainy season (S ₁)	Autumn season (S ₂)		
D ₁ (40-50)	21.72	21.03	21.37	
D ₂ (50-60)	22.24	21.78	22.01	
D ₃ (60-70)	22.79	22.06	22.42	
D ₄ (70-80)	23.13	22.80	22.96	
D _C (150-160)	23.48	23.06	23.27	
Mean	22.67	22.14		
Factors	C.D.	SE(d)	SE(m)	
Distance(D)	0.49	0.23	0.16	S
Season(S)	0.31	0.15	0.10	S
Distance X Season (DXS)	N/A	0.33	0.23	NS

*S= Significant at 5% probability level; NS = Non Significant; N/A= Not Applicable

Table 9. Plucking point density of tea as affected by gas flaring along

Seasons Distances (m)	Plucking point density (No./2500 cm ²)		Mean	
	Rainy season (S ₁)	Autumn season (S ₂)		
D ₁ (40-50)	32.00	30.66	31.33	
D ₂ (50-60)	33.66	32.66	33.16	
D ₃ (60-70)	37.00	36.33	36.66	
D ₄ (70-80)	40.33	38.66	39.49	
D _C (150-160)	42.00	41.00	41.50	
Mean	36.99	35.86		
Factors	C.D.	SE(d)	SE(m)	Significance
Distance(D)	1.55	0.73	0.52	S
Season(S)	0.98	0.46	0.33	S
Distance X Season (DXS)	N/A	1.04	0.73	NS

*S= Significant at 5% probability level; NS = Non Significant; N/A= Not Applicable

3.9 Plucking Point Density of Tea Bush

It had shown significant differences with respect to distance. Plucking point density had also shown significant variation in seasons and was higher in rainy season (Table 9). Plucking point density of tea bush was lowest in distances nearer to the gas flaring site. Air pollutants was higher in the surrounding environment of gas flare [22] which might reduce chlorophyll content of tea leaves that affected growth of tea bush reducing its plucking point density. This study finds similarity with the findings of Sarma [16].

4. CONCLUSION AND PERSPECTIVE

Gas flaring has effects in physiology of tea plants with respect to growth in the area under study where the studied plant parameters had shown an increasing trend with distances away from the gas flaring site except water saturation deficit of tea leaves which increased with the distances nearer to the gas flaring site. Further research may be carried out to observe the impact of gas flaring on tea yield, soil biological properties, pest and disease infestation, etc.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Emam EA. Gas flaring in industry: an overview. *Petroleum & Coal*. 2015;57(5): 532-555.
- Sharma KK, Hazarika S, Kalita B, Sharma B. Effect of Flaring of Natural Gas in Oil fields of Assam on Rice cultivation. *Indian Journal of Environmental Science and Engineering*. 2011; 53(3): 289-298.
- Ghadyanlou F, Vatani A. Flare gate recovery methods for olefin plants. *Chemical Engineering*. 2015;122(5):66-70.
- Ite AE, Udo JI. Gas Flaring and Venting Associated with Petroleum Exploration and Production in the Nigeria's Niger Delta. *American Journal of Environmental Protection*. 2013; 1(4):70-77.
- Ezenwaji EE, Okoye AC, Otti VI. Effects of gas flaring on rainwater quality in Bayelsa State, Eastern Niger-Delta region, Nigeria. *Journal of Toxicology and Environmental Health Sciences*. 2013;5(6): 97-105.
- Amadi AN. Impact of Gas-Flaring on the Quality of Rain Water, Groundwater and Surface Water in Parts of Eastern Niger Delta, Nigeria. *Journal of Geosciences and Geomatics*. 2014; 2(3):114-119.
- Weatherly PE. Studies in the water relations of the cotton plant. *New Phytol*. 1950;49:8197.
- Beakbane AB, Majumder PK. A relationship between stomatal density and growth potential in apple rootstocks. *Journal of Horticultural Science*. 1975; 50(4):285-289.
- Pandey SK, Singh H. A simple, cost-effective method for leaf area estimation. *Journal of Botany*. 2011;(3):1-6.
- Taylor S, Baker D, Owuor P, Orchard J, Othieno C, Gay C. A model for predicting black tea quality from the

- carotenoid and chlorophyll composition of fresh green tea leaf. *Journal of the science of Food and Agriculture*. 1992; 58(2):185-191
11. Bray HC, Thorpe WV. Analysis of phenolic compounds of interest in metabolism. *Methods of Biochemical Analysis*. 1954;1: 27-52.
 12. Baruah DN, Dutta KN. Distribution of shoots on the plucking surface of a tea bush and its relation to spacing. *Two and A Bud*. 1973;18:8-11.
 13. Naz N, Durrani F, Shah Z, Khan NA, Ullah I. Influence of heat stress on growth and physiological activities of potato (*Solanum tuberosum* L.). *International Journal of Experimental Botany*. 2018; 87: 225-230.
 14. Wahhed A, Hamid FS, Shah AH, Ahmad H, Khalid A, Abassi FM, Ahmad N, Aslam S, Sarwar S. Response of different tea (*Camellia sinensis* L.) clones against drought stress. *Journal of Materials and Environmental Science*. 2012;3(2):395-410.
 15. Isichei AO and Sanford WW. The Effects of Waste Gas Flares on the Surrounding Vegetation in South-Eastern Nigeria. *Journal of Applied Ecology*. 1976;13(1): 177-187.
 16. Sarma B, Chanda SK, Bhattacharyya PR. Dynamics of morpho-physiological characteristics of *Cassia tora* L. (Fabaceae) in oil field pollution stress environment. *International Journal Environmental Science and Technology*. 2019;16:4091-4102.
 17. Treshow M. *Environment and Plant Response*. McGraw-Hill, New York; 1970.
 18. Song R, Kelman D, Johns K, Wright A. Correlation between leaf age, shade levels, and characteristic beneficial natural constituents of tea (*Camellia sinensis*) grown in Hawaii. *Food Chemistry*. 2012; 133:707-714.
 19. Lee LS, Choi JH, Son N, Kim SH, Park JD, Jang DJ. Metabolomic analysis of the effect of shade treatment on the nutritional and sensory qualities of green tea. *Journal of Agricultural and Food Chemistry*. 2013; 61:332-338.
 20. Zhang Q, Shi Y, Ma L, Yi X, Ruan J. Metabolomic analysis using ultra-performance liquid chromatography-quadrupole-time of flight mass spectrometry (UPLC-Q-TOF MS) uncovers the effects of light intensity and temperature under shading treatments on the metabolites in tea. *Plos One*. 2014; 9(11):1-10.
 21. Erturk Y, Ercisli S, Sengul M, Eser Z, Haznedar A, Turan M. Seasonal variation of total phenolic, antioxidant activity and minerals in fresh tea shoots (*Camellia sinensis* var. *sinensis*). *Pakistan Journal of Pharmaceutical Sciences*. 2010;23(1):69-74.
 22. Bhattacharyya PR. Studies on the effect of oil field pollution upon nutritional quality of "Som" (*Machilus bombycina*) and the biology and certain biochemical aspects of Muga silkworm. Ph.D. Thesis, Gauhati University, Assam; 1994.

© 2022 Phukan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/85416>