



Evaluating the Potential of Phytochemicals as Natural Substitute for Synthetic Antioxidant: A Review

Mohammad Hossein Maleki ^a, Milad Daneshniya ^{a*}, Zahra Latifi ^b,
Younes Pirouz Zarrin ^c, Maral Behzadnia ^d and Naghmeh Morakabati ^e

^a Department of Food Science and Engineering, Faculty of Industrial and Mechanical Engineering, Qazvin Branch, Islamic Azad University, Qazvin, Qazvin Province, Iran.

^b Young and Elite Researchers Club of Islamic Azad University, Sari Branch, Mazandaran Province, Iran.

^c Department of Veterinary Medicine, Faculty of Veterinary, Islamic Azad University, Tabriz Branch, East Azerbaijan Province, Iran.

^d Department of Food Science and Technology, College of Pharmaceutical Sciences, University of Medical Sciences, Islamic Azad University, Tehran, Iran.

^e Department of Food Science and Technology, Faculty of Food Science and Technology, Noor Branch, Islamic Azad University, Noor, Mazandaran Province, Iran.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRB/2022/v10i130214

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/84048>

Review Article

Received 17 November 2021

Accepted 22 January 2022

Published 25 January 2022

ABSTRACT

This study aimed at investigating the antioxidant potential of flavonoids as a subgroup of polyphenols and ultimately entirely and partially substitute the synthetic antioxidant with natural them. All relevant databases were searched for the terms including 'phytochemical', polyphenol', 'flavonoid', 'natural additive', as well as 'antioxidant activity' combined with *in vivo* and *in vitro* phrases. Free radicals are among the damaging factors to macromolecules. The oxidative reaction caused by these radicals is one of the reasons for food spoilage, which causes unpleasant odor, loss of taste, damaged tissue and appearance, and loss of nutritional value of exposed material.

*Corresponding author: E-mail: miladdaneshnia@gmail.com; miladdaneshnia@qiau.ac.ir.

The common antioxidants employed in foods include Butylated hydroxytoluene (BHT), Butylated hydroxyanisole (BHA), Tert-butyl hydroquinone (TBHQ), and Propyl gallate (P.G.). Synthetic antioxidants have adverse effects such as causing mutation and carcinogenicity in the human body despite their high performance; hence, many studies have been carried out on replacing them with natural antioxidants. Phytochemicals and a widespread group of them known as polyphenols possess a high antioxidant activity. A major subset of polyphenols, Flavonoids, are secondary metabolites in plants with many applications. These compounds are potential antioxidants because of their capabilities, such as scavenging the free radicals donating hydrogen atoms, electrons, and chelate metal cations. The antioxidant mechanism of action of flavonoids is transferring hydrogen toward free radicals. Accordingly, the more the flavonoid structure makes the hydrogen transfer faster and easier, the more the flavonoid's antioxidant power will be. Therefore, the antioxidant activity of the flavonoids with hydroxy groups in their structure is higher among the different flavonoids. Besides health promotion and some disease prevention effects, various in vitro studies have shown that flavonoids possess a high antioxidant activity that is competitive with synthetic antioxidants. However, to make it commercially available, cheaper and high-performance extraction methods of flavonoids will have to be developed.

Keywords: Antioxidant; flavonoids; food preservation; phytochemical; polyphenol.

1. INTRODUCTION

Most food products are made up of various chemical components that are easily oxidized. Lipids (fats, oils, and waxes) in general have significant potential to lose electrons. Their auto-oxidation in food products initiated by metal ions exposure, metallo-protein catalysts, light, heat, or ionizing radiation can deteriorate the color, flavor, texture, quality, and safety of food. Food fats are chemically made of triglycerides, and oxidation occurs at the unsaturated sites of the triglycerides, resulting in rancidity [1,2]. Free radicals are the most frequent oxidants in biological systems. These are atoms, molecules, or ions with one or more unpaired electrons, turning them to highly reactive and unstable. In order to stabilize themselves, unpaired electrons in these free radicals seek out and acquire electrons from other substances. However, the first assault pairs the odd electron, another free radical is produced in the process, resulting in a chain reaction. The free radicals cause considerable damages to the macromolecules in organisms (of bodies), including DNA, proteins, lipids, and carbohydrates. The damages caused by free radicals are also referred to as oxidative damages [3-6]. Free radicals are frequently derived from oxygen, nitrogen, and sulfur molecules in biological systems. These free radicals are constituents of the reactive oxygen species (ROS), reactive nitrogen species (RNS), and reactive sulfur species (RSS) molecular families. Free radicals such as superoxide anion ($O_2^{\bullet-}$), per hydroxyl radical (HO_2^{\bullet}), hydroxyl radical (O.H.), nitric oxide, and other species like hydrogen peroxide (H_2O_2), singlet oxygen (O_2), hypochlorous acid (HOCl), and peroxyxynitrite

(ONOO⁻) are forms of ROS. Nitric oxide is converted to RNS by reacting with $O_2^{\bullet-}$ to generate ONOO⁻. By reacting thiols with ROS, RSS is quickly produced. The production of ROS occurs as a result of cellular metabolism and functional activities [1,7].

Antioxidants are one of the factors which prevent oxidative damages. The structure of antioxidants is such that although they release hydrogen atoms, the reactivity of these atoms is extremely low [8]. In general, antioxidants are divided into two groups - synthetic and natural. Because synthetic antioxidants are cheaper, more accessible, having high efficiency and stability, they were gradually considered as alternatives to their natural type after World War II [9]. Phenolic compounds among the phytochemical are known to be the most important natural antioxidants, and they are recognized as excellent antioxidants due to their capacity of scavenging free radicals, donating hydrogen atoms, electrons, and chelate metal cations [10,11]. The natural antioxidant can be considered as health-promoting compounds as well as food preservatives. These antioxidants can help to limit or prevent the production of chemical toxins, improving the nutritional value and health benefits of the food products and also extending their shelf life. While called natural, overdose usage of these antioxidants can cause poisoning in consumers; hence legislation for their utilization on the industrial scale should be considered [12]. Nowadays, synthetic antioxidants are commonly used in industries, the most common of which include Butylated hydroxytoluene (BHT), Butylated hydroxyanisole (BHA), Tert-butyl hydroquinone (TBHQ), and

Propyl gallate (P.G.). Many other suggested compounds have antioxidant activity, of which only a few can be used in foods. Ethoxyquin (EMQ) is a synthetic antioxidant with a non-phenolic structure, the use of which is not permitted for human consumption compared to other antioxidants mentioned earlier; however, it is used for food preservation in animal feeds [13]. Synthetic antioxidants are available for the industries with an economically profitable price which consequently leads to lower price of the product for costumers; they possess the effective potential to preserve the product and extend the shelf life, the common type of them are applicable for a wide range of the products and have a specified dosage for utilization in each product as well as known changes associate with their addition. Nevertheless, synthetic antioxidants can also initiate or development of numerous diseases such as cancer, cardiovascular diseases, type 2 diabetes mellitus, cataract, rheumatoid arthritis, or different neurodegenerative diseases. They can damage DNA and induce mutations, resulting in the onset of neoplasia. Furthermore, these antioxidants may cause lipid and blood cholesterol levels to surge in the body lead to the development of enzymes for its metabolization in the liver, posing a risk of destroying vital chemicals like vitamin D, acting as inducers of urticaria and eczematous dermatitis [14,15].

The use of antioxidants in foods has been limited by the regulatory laws of one country or international standards [16], so the consumption of synthetic antioxidants is limited due to their toxic effects and carcinogenicity. Since the carcinogenicity of synthetic antioxidants has been observed, the necessity of using alternatives without adverse effects is becoming increasingly important [17]. Therefore, in recent years, the demand for natural antioxidants, especially of plant origins, has increased because of growing concern on consumption health due to their toxicological effects [18-20]. Hence, this review has been conducted to investigate the antioxidant activity of phytochemicals, emphasizing *in vitro* studies to shed light on the potential of flavonoids to be utilized as natural antioxidants in food and examine their capability to substitute the synthetic antioxidants.

2. METHODOLOGY

This review was conducted to investigate the antioxidant activity of flavonoids and their

potential to entirely or partially substitute synthetic antioxidants in food products. All relevant databases were searched for the terms including 'phytochemical', 'polyphenol', 'flavonoid', 'natural additive', as well as 'antioxidant activity' combined with *in vivo* and *in vitro* phrases. It should also be mentioned that in the process of selecting relevant papers, priority was given to the research providing *in vitro* information regarding the antioxidant activity of phytochemicals. Data was collected through search using Scopus, Pubmed, Web of Science, Science Direct, Google Scholar databases, and related books and reliable websites, as well as prominent conferences and congresses with various publication dates ranging from 1981 to 2021.

3. DISCUSSION

Phytochemicals are chemical compounds and bioactive substances of plants that are naturally found in plants associated with biological activities beneficial for human health. These compounds are categorized by their chemical structures and include polyphenols (flavonoids and non-flavonoid polyphenols), carotenoids, phytosterols, betalains, and organosulfur. Phytochemical compounds cause the color, odor, and taste of plants to be changeless and play an important role in the plant defense system. More than 5000 pigments in plants are considered phytochemicals. These compounds can reveal their biological activity through various mechanisms such as antiviral, antibacterial properties, antioxidant, anti-inflammatory, anticancer, modulation of enzyme activities, stimulation of the immune system, gene expression, regulation of cholesterol synthesis, and blood pressure. They can also improve health and prevent diseases [21, 22]. Studies have shown that the presence of these compounds in the fruits and vegetables-rich diets help in delaying aging, reducing inflammation and oxidative stress risk and also decreasing the risk of chronic diseases such as cardiovascular diseases, arteriosclerosis, cancer, diabetes, cataract, neurological diseases, and disorders of the cognitive function [23-25]. Fruits, vegetables, legumes, and whole grains are the primary sources of phytochemicals in the diet [26]. There are different methods for extracting phytochemical compounds. These compounds can be extracted from plants through physical and chemical methods such as ultrasonication, super-critical fluid extraction, solvent extraction, and cold pressing [21,27]. It should be pointed

out that many phytochemicals are insoluble in aqueous solutions and do not have acceptable oil solubility. These compounds are chemically unstable and will decompose rapidly if being exposed to the external environment and adjacent to oxygen, temperature, light, pH, and some other reactive substances.

Phenolic compounds or polyphenols are important groups of phytochemical compounds in plants [28, 29]. Phenolic compounds have excellent antioxidant characteristics and are often found in fruits and vegetables [30]. These compounds include flavonoids, flavanols, anthocyanins, anthraquinones, benzoyl acetyl, and their derivatives. Polyphenols have two general classes, one is flavonoids, and the other is phenolic acids. Different parts of the plant produce compounds that are mainly phenolic and are known as natural antioxidants. Compounds such as flavonoids, coumarins, tocopherols, carotenoids, organic acids, and derivatives of cinnamic acid are phenolic antioxidants from the plant source. In this category of compounds, the hydroxyl group is directly bonded to an aromatic hydrocarbon [31]. There are at least 8000 known structures of phenolic compounds, the simplest of which is Phenol shown in Fig. 1 [28]. Phenolic compounds are known for their antioxidant activity that depends on the structure, especially the number and location of hydroxyl groups and the nature of substituents in aromatic rings [32].




Fig. 1. Phenol, the most straightforward component of phenolic compounds

Phenolic compounds are referred to as health promotion compounds and have the capability of preventing chronic heart disease [33, 34]. The antioxidant characteristics of phenolic compounds depend on their capability of giving electrons to trap free radicals by the formation of stable phenoxyl compounds [35].

Plants have always been excellent food sources for the consumption of valuable bioactive compounds [36]. These natural antioxidants have been extracted from plants in the form of essential oils and extract from various sources such as fruits (grapes, pomegranates, dates, kinnow), vegetables (broccoli, potatoes,

drumstick, pumpkin, Indian turmeric, nettle), medicinal plants, and spices (tea, rosemary, oregano, cinnamon, common sage, thyme, peppermint, ginger, clove) and have been investigated for reducing fat oxidation [37-42]. Phenolic compounds are discovered in combination with saccharides (monosaccharides and polysaccharides) bonded to one or more phenolic groups. Although some phenolic compounds are ubiquitous, others are specific to particular plant families and are found in particular plant organs or at specific stages of plant growth [43]. Important phenolic compounds in plants that have been reported to possess antioxidant activity in *In vitro* and *in vivo* studies are listed in Table 1.

3.1 The Structure of Flavonoids and Their Antioxidant Activity

Flavonoids are the largest group of phytochemicals and are also considered as the most widespread group of polyphenols found in fruits and vegetables. Flavonoids have excellent antioxidant properties and can exhibit their antioxidant activity by scavenging free radicals and reactive oxygen species, chelating the metals, and preventing the oxidation of low-density lipoproteins (LDLs) [30,79]. Flavones are the most basic structures of flavonoids. From the structural point of view, all flavonoids possess a C6-C3-C6 carbon skeleton in which the carbon atoms are arranged in three phenolic rings A, B, and C, and the C ring normally contains oxygen [80, 81]. Fig. 2 indicates a Phenylbenzopyrane as the basic structure of flavonoids.




















Fig. 2. Basic structure of flavonoids (Phenylbenzopyrane)






Flavonoids have different families, each of which has different members. The flavonoid families include Flavones, Isoflavones, Flavonols, Flavanones, Dihydroflavonols, Flavan-3-ols (monomers), Proanthocyanidins (oligomeric flavan-3-ols), and Anthocyanidins. These families, along with their members, are shown in Table 2 [82].







Table 1. The phenolic compounds with high antioxidant activity






Name	Chemical formula	Structure	Reference(s)
Gallic acid	$C_7H_6O_5$		Gallic acid prevents the melanogenesis, rancidity, and spoilage of fats and oils due to its antioxidant nature, facilitating its application as food additives in various eatable materials like baked goods, candy, chewing gums, and cosmetics [44,45].
<i>p</i> -Coumaric acid	$C_9H_8O_3$		<i>p</i> -coumaric acid occurs widely in the cell walls of graminaceous plants. It decreases low-density lipoprotein peroxidation, shows antioxidant and antimicrobial activities, and plays a vital role in human health. It is a good antioxidant and a good antimicrobial; therefore, it is a natural alternative instead of synthetic additives [46]
Rosmarinic acid	$C_{18}H_{16}O_8$		A significant number of herbal preparations and food supplements containing rosmarinic acid are marketed with claims for beneficial health effects. Due to lipid peroxidation and bacterial growth inhibition, Rosmarinic acid is approved for use as a natural antioxidant and/or preservative in the food industry [47].
Caffeic acid	$C_9H_8O_4$		Natural phenolic antioxidants, including caffeic and ferulic acids, gained attention as photoprotective agents. Caffeic acid is a α -tocopherol protective in low-density lipoprotein. Furthermore, their combination with other products, such as chlorogenic and caftaric acids, showed more potent antioxidant activity in various systems [48].
Protocatechuic acid	$C_7H_6O_4$		Protocatechuic acid is a type of widely distributed naturally occurring phenolic acid. Protocatechuic acid has structural similarities with gallic acid, caffeic acid, vanillic acid, and syringic acid, which are well-known antioxidant compounds. Hence, it could therefore be used in the pharmacological or food industry as a natural antioxidant [49].

Carnosol	$C_{20}H_{26}O_4$		<p>Several <i>in vitro</i> studies were reviewed regarding the antioxidant activity of the carnosol and carnosic acid. Using the 2,2-diphenyl-1-picrylhydrazyl method, these bioactive compounds were validated for their antioxidant activity. Also, using the thiobarbituric acid, superoxide anion, and lipid-free radicals scavenging activity assays and Rancimat methods (determination of oxidative stability of fat), carnosol has been reported to inhibit lipid peroxidation through the lipid-free radical scavenging mechanism. These studies have shown the antioxidant potential of carnosol and carnosic acid, whose properties are closely related to other biological activities, such as cytoprotective and anticancer, primarily due to their ability to neutralize reactive oxygen species [50].</p>
Carnosic acid	$C_{20}H_{28}O_4$		
Flavanol (catechin)	$C_{15}H_{14}O_6$		<p>It has been revealed that catechin is antioxidant at all concentrations and attributed the antioxidant behavior to iron chelation. Consequently, catechin has been a widely used antioxidant in combination with other biopolymers for the development of active food packaging [51,52].</p>
Flavanone (naringenin)	$C_{15}H_{12}O_5$		<p>Naringenin has exhibited higher antioxidant capacity and hydroxyl and superoxide radical scavenger efficiency than naringin. Additionally, naringenin has shown greater effectiveness in the protection against oxidative damage to lipids in a dose-dependent manner [53].</p>
Flavonol (quercetin)	$C_{15}H_{10}O_7$		<p>Due to the capacity of quercetin to scavenge free radicals and bind transition metal ions, quercetin is considered a powerful antioxidant. Because of these qualities, quercetin can prevent lipid peroxidation [54].</p>

Flavone (Apigenin)	$C_{15}H_{10}O_5$		Apigenin is a widely distributed flavonoid responsible for antioxidant potential and chelating redox-active metals. The antioxidant mechanism of apigenin includes oxidant enzymes inhibition, modulation of redox signaling pathways, reinforcing enzymatic and nonenzymatic antioxidants, metal chelation, and free radical scavenging [55].
Anthocyanin (delphinidin)	$C_{15}H_{11}O_7^+$		Anthocyanins are polyphenols with the known antioxidant activity which can be responsible for various biological activities. The antioxidant capacity of plant-derived anthocyanins has been demonstrated with various assay methods, including ORAC, FRAP, TEAC, and DPPH. [56].
Isoflavone (genistein)	$C_{15}H_{10}O_5$		Genistein has been widely studied as an antioxidant, with some research claiming that it may have the ability to scavenge reactive oxygen and nitrogen species (ROS and RNS) effectively [57].
Rosmanol	$C_{20}H_{26}O_5$		Comparison between rosmanol and synthetic antioxidants has shown that rosmanol possesses an antioxidant capacity four times higher than synthetic antioxidants of BRT and BRA [58].
Epicatechin	$C_{15}H_{14}O_6$		Catechin and epicatechin are two flavan-3-ol stereoisomers that, according to their structure, have comparable radical scavenging and antioxidant activity. Both have the ability to bind the iron ion and scavenge hydroxyl, peroxy, and 2,2-diphenyl-1-picrylhydrazyl (DPPH) radicals [59].
Epicatechin gallate(ECG)	$C_{22}H_{18}O_{10}$		Evaluation of radical scavenging activity of some catechins <i>in vitro</i> revealed that in each of the assays, including DPPH, ABTS, and FRAP after epigallocatechin gallate, epicatechin gallate had the highest potential [60].

Epigallocatechin gallate (EGCG)	$C_{22}H_{18}O_{11}$		Epigallocatechin gallate effectively protected against supercoiled DNA nicking caused by peroxy and hydroxyl radicals. In scavenging the 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical, epigallocatechin gallate had a synergistic impact with α -tocopherol, demonstrating a direct free radical scavenging potential [61].
Rutin	$C_{27}H_{30}O_{16}$		Rutin has exhibited strong DPPH radical scavenging activity. In addition, rutin possesses effective inhibition property of lipid peroxidation. The antioxidant activity of rutin is competitive with various antioxidants such as BHT and Vc (ascorbic acid) [62].
Eugenol	$C_{10}H_{12}O_2$		Antioxidant activity of eugenol has been evaluated with different assays such as DPPH, ABTS, and DMPD are competitive with inhibiting potential for lipid peroxidation of standard antioxidants including BHA, BHT, α -tocopherol, and Trolox [63].
Carvacrol	$C_{10}H_{14}O$		The antioxidant activity of carvacrol has been shown through a series of <i>in vitro</i> assays, including hydroxyl radical, superoxide, nitric oxide, DPPH, and ABTS radical scavenging assays [64].
Thymol	$C_{10}H_{14}O$		Different reports regarding antioxidant properties of thymol are due to its ability to donate H-atoms from phenol hydroxyl groups, which could react with peroxy radicals to produce stabilized phenoxyl radicals and, consequently, terminate lipid peroxidation chain reactions [65].

Safrole	$C_{10}H_{10}O_2$		The antioxidant activity of safrole oil has been reported to be comparatively lower than the Trolox reference standard, which is known to have high antioxidant activity. However, free radicals scavenging activity of safrole is still considered eminent [66]
Menthol	$C_{10}H_{20}O$		Different studies have reported that the antioxidant activity of some of the plant-derived essential oils is related to menthol; for instance, the radical scavenging activity of peppermint essential oil is associated with the presence of menthol and menthone, which contains the hydroxyl radical (·O.H.) [67, 68].
Myristicin	$C_{11}H_{12}O_3$		While evaluating the myristicin with DPPH and ABS methods revealed its antioxidant activity inadequacy compared to BHA, the synergetic effect of myristin with other antioxidants can be considered [69].
Eucalyptol	$C_{10}H_{18}O$		Previous researches have shown that there might be a direct relationship between the high scavenging activity of essential oil over BHT and the high content of Eucalyptol in the essential oil [70, 71].
Terpineol	$C_{10}H_{18}O$		Due to various biological activities, terpineol is a proper compound to be improved the shelf life of food products, as it has been shown to be effective for enhancing the antioxidant activity of chitosan biopolymer to inhibit free radicals mediated deterioration [72].
<i>p</i> -Cymene	$C_{10}H_{14}$		Previous investigations on DPPH free radical scavenging by some plant-derived essential oils have revealed that <i>p</i> -Cymene, as one of the main constituents, plays a prominent role in the antioxidant activity of the essential oils [73].

Cinnamaldehyde	C_9H_8O		Cinnamaldehyde enhances the antioxidant defense against reactive oxygen species and reduces the levels of lipid peroxidation products, and increases the activities of antioxidant enzymes [74].
Piperine	$C_{17}H_{19}NO_3$		Piperine interacts readily with highly oxidizing radicals and binds redox-active metal ions, indicating that it could potentially function as an effective antioxidant [75].
Propyl gallate	$C_{10}H_{12}O_5$		As it has become determined by free radical scavenging assays, the propyl gallate compound has the potential to inhibit lipid oxidation which is competitive with TBHQ [76].
Resveratrol	$C_{14}H_{12}O_3$		In different studies, resveratrol has exhibited a robust reducing power, chelating activity on Fe^{2+} , free radical-scavenging, hydrogen peroxide scavenging, and hydroxyl radical scavenging activities [77].
Diosmin	$C_{28}H_{32}O_{15}$		The <i>in vitro</i> studies on diosmin showed its free radical scavenging activity. The hydroxyl free radical scavenging and superoxide free radical scavenging activities of diosmin have been evidenced by the <i>in vitro</i> studies [78].

Each of these flavonoids can possess completely distinct antioxidant activity due to their specific structure. Many *in vitro* and *in vivo* studies utilizing natural flavonoids have been conducted to confirm the relationships between flavonoid structure and antioxidant activity. The hydroxyl groups (C in Fig. 3), ortho-dihydroxy arrangement in the B ring (A in Fig. 3), C2-C3 unsaturated bond combined with C-4 carbonyl group in the C skeleton (B in Fig. 3), and O-methylation have all been identified as distinct chemical structures related to flavonoids antioxidant activities [83, 84].

The free hydroxyl group will donate its hydrogen atom to a radical molecule, stabilizing it and generating a flavonoid phenoxyl radical that is relatively stable. This stable molecule can then react with a second radical (Alkoxy) to form a stable quinone structure. Furthermore, the location of the hydroxyl groups, rather than the overall quantity of hydroxyl groups, has a significant impact on antioxidative action. Scavenging of both oxygen and nitrogen-centered free radicals is dominated by the B-ring hydroxyl structure. Hydroxyl groups on this nucleus provide hydroxyl, peroxy, and peroxy nitrite radicals hydrogen and an electron, stabilizing them and forming a fairly stable flavonoid radical. Moreover, the twisting angle of the B-ring in relation to the remainder of the molecule has a significant impact on its capacity to scavenge free radicals [85-88].

3.2 Regulations Regarding Synthetic and Natural Antioxidants

Many scientists claim that there should be a shift away from synthetic antioxidants due to the growing interest in natural food antioxidants; however, it has not yet been achieved, and thus the natural or synthetic origin of the various antioxidants currently used in the food industry is not specified in the official tables showing the amounts and permissions for the use of each additive in each type of food [89]. The Codex Alimentarius Commission is the worldwide standard-setting authority in charge of regulating and establishing the framework of food regulations that guide the use of antioxidants in foods. Since food regulatory systems and legal frameworks in respect to the use of antioxidants as food additives differ by country, these criteria are neither required nor immediately applicable [90]. The European Food Safety Authority (EFSA) of the European Union and the Food and Drug Administration (FDA) of the United States of America are the two primary guiding

authorities in charge of regulating the licensing of food additives across the world [91]. Under the E.U., regulation E.C. No 1129/2011 specifies a list of antioxidants, categorized according to their E-numbers, that are regulated in the section of "other food additives." The natural antioxidants recognized as food additives by the E.U. include rosemary extracts (E392), ascorbic acid (E300), and tocopherols (E306–E309), according to this regulation [92]. Food products and ingredients are monitored by various authorities in the United States of America, the most relevant of which is the U.S. Food and Drug Administration (FDA). The FDA's regulations for food items are listed in Title 21 of the Code of Federal Regulations of the United States. There is no distinct category for natural antioxidants in the Code of Federal Regulations (21 CFR 182; 21 CFR 172), as there is in European regulations [93-95]. Although natural antioxidants such as ascorbic acid and tocopherols are specifically designated for use in food, the U.S. regulation is far more comprehensive than the European one and includes additional substances that fall into other categories but yet have demonstrated antioxidant activity. Several of the compounds allowed for use as coloring adjuncts, spices, or natural flavorings, such as phloretin glycoside (flavonoids), carotene, and astaxanthin (carotenoids), or extracts of rosemary or sage, have recognizable antioxidant potential. These antioxidants - phloretin, astaxanthin, and sage extract are not listed as food additives in any category in E.U. regulations, although they are classified for technical applications in U.S. standards [96, 97]. Considering the mentioned regulations, it is evident that regardless of utilization of natural or synthetic antioxidants, the optimal choice of antioxidant for each food matrix should be determined on a case-by-case basis, as *in vitro* antioxidant activities are not always reproduced in the food, and there are frequent changes due to food processing or interactions with other food matrix components that have antioxidant or prooxidant activities [98].

3.3 Antioxidants in Food Preservation

Once lipid oxidation in a food matrix is analyzed over time, there is frequently a lag phase during which the deposition of lipid oxidation by-products is moderated. This lag phase is associated with the presence of antioxidants in the food matrix, which limit the synthesis of free radicals that target fatty acids, as well as the low free radical generation that precedes the concentration of hydroperoxides and β -scission

reactions [89]. The objective of the food industry is to extend the lag period during which the concentrations of the components that cause rancidity taste are below human detection limits. When the food matrix lacks natural antioxidants and/or includes significant endogenous prooxidants, the most typical technique employed by food manufacturers to limit oxidation is to add antioxidants directly to the food matrix [98]. In the context of food science, the phrase "antioxidant" refers to substances that prevent lipid peroxidation and other oxidative reactions, hence preserving the freshness and shelf-life extension of foods. The method of action is the same whether the antioxidant is natural or synthetic, and it comprises free radical scavenging, metal chelating, and singlet oxygen quenching [99, 100]. Due to their stability, low cost, and wide availability, synthetic antioxidants are frequently employed. Phenolic antioxidants are the most extensively used synthetic antioxidants in the food industry, with Butylated hydroxyanisole (BHA) (E320), Butylated hydroxytoluene (BHT) (E321), Propyl gallate (P.G.) (E310), and Tert-butyl hydroquinone (TBHQ) (E319) being the most common. In Table 3, some critical information regarding the chemical structure as well as health-related concerns about the consumption of these antioxidants has been revealed [89, 101-110]. Despite the fact that these synthetic antioxidants are widely used and tightly regulated, there are concerns about their safety owing to overdose usage and/or misuse; for instance, combining various antioxidants may exacerbate their toxic effects. High concentrations of chemical synthetic antioxidants may induce DNA damage or *in vitro* toxicity in some tissues [111-114]. The food industry is attempting to reduce the use of synthetic compounds by substituting natural alternatives in response to consumer concerns that they are being exposed to potentially toxic chemical synthetic compounds through their regular diet. Though, just because plant antioxidants are natural, it does not mean they are safe by default; toxicity tests are always required for each case [111].

3.4 The Structure of Flavonoids and Their Antioxidant Properties

Flavonoids are active polyphenol compounds that are found in almost all fruits and vegetables. Aglycones, glycosides, and methylated derivatives are different forms of flavonoids. Aglycones are produced when the glycosyl group in a glycoside is replaced by a hydrogen atom. Glycosides are composed of two parts — sugars

(glycons) and non-sugars (aglycones) — and play different roles in plants [6]. Various species of flavonoids differ in terms of oxidation rate and C-ring substituent; in other words, the presence of one double bond, one carbonyl group, and one hydroxy group in the pyranil C ring is used to classify them into species and subspecies. The substituents of A and B rings with hydroxy groups are used to identify the available members in a species. Flavonoids in plants are often derived from glycosylate and play a role in the production of shining blue, reddish-yellow, and reddish-orange shades in leaves, flowers, and fruits. Regardless of the variety of fruits and vegetables, flavonoids are found in seeds, nuts, buds, spices, herbal medications, and some drinks such as wine, especially red and tea wines, and in smaller quantities in beer. The antioxidant properties are among the most important properties of flavonoids. Flavonoids are highly effective antioxidants in general, and it has been observed that they protect the body against cardiovascular diseases by reducing the oxidation of low-density lipoproteins (LDLs). In many studies, flavonoid-rich plant extracts have been used to prevent food oxidation [7]. Despite the many benefits of synthetic antioxidants compared to natural ones, such as their low prices, the use of antioxidants in food has been limited by regulatory laws of one country or international standards [13]. Hence, the use of synthetic antioxidants is limited due to their toxic effects and carcinogenicity. Synthetic antioxidants have a limit of using 0.02% fat of food [6]. Since the carcinogenic properties of synthetic antioxidants have been observed, the necessity of the use of alternatives without adverse effects has become increasingly important [14]. Therefore, in recent years, the demand for natural antioxidants, especially of plant origin, has increased due to the increasing concern among consumers about these synthetic antioxidants considering potential toxicological effects [20]. As a whole, flavonoids are highly effective antioxidants, and there are pieces of evidence that they protect the body against cardiovascular diseases by reducing the oxidation of low-density lipoproteins (LDLs). Flavonoids both prevent lipid peroxidation and act as scavengers of radicals such as superoxides, lipid peroxy, and hydroxyls, and also lead to inactivation of single oxygen molecules and prevention of the activity of lipoxygenases. The high potency of flavonoids in inhibiting free radicals relates to their ability to transfer a hydrogen atom from a hydroxy group to the free radical and ultimately stabilize it:

Table 2. Flavonoids family














Primary structure	R groups alterations		Primary structure	R groups alterations						
Anthocyanidins			Flavan-3-ols (monomers)							
	R1	R2		R1	R2	R3				
	Pelargonidin	H		(+)Catechin	OH	H	H			
	Cyanidin	OH		(-)Epicatechin	H	OH	H			
	Delphinidin	OH		(+)Gallocatechol	OH	H	OH			
	Peonidin	OCH ₃		(-)Epigallocatechin	H	OH	OH			
	Petunidin	OCH ₃		(+)Catechin gallate	OG	H	H			
	Malvidin	OCH ₃		(-)Epigallocatechin gallate	H	OG	H			
				(+)Gallocatechin gallate	OG	H	OH			
				(-)Epigallocatechin gallate	H	OG	OH			
Proanthocyanidins (oligomeric flavan-3-ols)			Isoflavones							
	R1	R2		R1	R2					
	Procyanidin B1	H		Daidzein	H	H				
	EC (4b-8)-C			Genistein	OH	H				
	Prodelfinidin B1	OH		Glycitein	H	OCH ₃				
	EGC-(4b-8)-GC	OH								
Flavonols			Flavones							
	R1	R2		R1	R2	R3	R4	R5	R6	
	Kaempferol	H		Apigenin	H	OH	OH	H	OH	H
	Quercetin	OH		Luteolin	OH	OH	OH	H	OH	H
	Myricetin	OH		Tangeretin	H	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCH ₃
				Nobiletin	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCH ₃
Flavanones			Dihydroflavonols							
	R1			R1						
	Naringin	H		Taxifolin	H					
	Eriodictyol	OH		Ampelopsin	OH					
	Hesperetin	OCH ₃								

Table 3. Chemical structures of food industry antioxidants

Antioxidants	Molecular formula	Carcinogenic potential	Allergenic potential	Chemical structures
Butylated hydroxyanisole (BHA)	$C_{11}H_{16}O_2$	Category 1 (use within limit on body = 5.2 % w/w; face >100% w/w; hands >100 % w/w)	Moderate sensitizer (max dose = 0.0729 $\mu\text{g}/\text{cm}^2/\text{hr}$)	
		Acceptable daily intake (ADI)*: below 1.0 mg/kg body weight		
Butylated Hydroxytoluene (BHT)	$C_{15}H_{24}O$	–	Weak sensitizer (max dose = 14.6 $\mu\text{g}/\text{cm}^2/\text{hr}$)	
		Acceptable daily intake (ADI): 0.25 mg/kg body weight		
tert-Butylhydroquinone (TBHQ)	$C_{10}H_{14}O_2$	–	–	
		Acceptable daily intake (ADI): 0.7 mg/kg body weight		
Propyl Gallate (P.G.)	$C_{10}H_{12}O_5$	–	Moderate sensitizer (max dose = 0.0255 $\mu\text{g}/\text{cm}^2/\text{hr}$)	
		Acceptable daily intake (ADI): 0.5 mg/kg body weight		
Ascorbic Acid (Vitamin C)	$C_6H_8O_6$	Category 3 (use with precaution)	–	
		Acceptable daily intake (ADI): there is no safety concern for the use of ascorbic acid (E300), sodium ascorbate (E301), and calcium ascorbate (E302) as food additives at the reported uses and use levels, and there is no need for a numerical ADI for ascorbic acid and its salts.		

Erythorbic acid	$C_6H_8O_6$	–	–	Acceptable daily intake (ADI): 6 mg/kg body weight	
Ascorbyl Palmitate	$C_{22}H_{38}O_7$	–	Weak sensitizer (max dose = 7.29 $\mu\text{g}/\text{cm}^2/\text{hr}$)	Acceptable daily intake (ADI): considering the available data, the Panel concluded that there is no safety concern for the use of ascorbyl palmitate (E304(i)) and ascorbyl stearate (E304(ii)) as food additives at the reported uses and use levels	
Sodium bisulfite	$NaHSO_3$	–	–	Acceptable daily intake (ADI): 0.7 mg SO_2 equivalent/kg body weight	
alpha-tocopherol	$C_{29}H_{50}O_2$	–	–	Acceptable daily intake (ADI): 11-13 mg/kg body weight	

**Mentioned quantities are specified for adults. For more details or quantity for a specific age group, please refer to the released document at European Food Safety Authority.*





Fig. 3. The flavonoid structure-antioxidant activity relationships



Flavones and catechins are the most powerful flavonoids with antioxidant properties that can protect the body against reactive oxygen species (ROS). According to many studies, the free radical scavenging activity for flavonoids has been reported as follow [6, 115]:

Myricetin > quercetin > rhamnetin > morin > diosmetin > naringenin > apigenin > catechin > 5,7-Dihydroxy-3,3',4'-trimethoxyflavone > robinin > kaempferol > flavones

Many flavonoids have significantly higher antioxidant capacity than vitamins C and E. On an in vitro scale, flavonoids can prevent free radical harm through the mechanisms including direct scavenging of ROS, activation of antioxidant enzymes, metal chelating activity, reduction of α -tocopheryl radicals, inhibition of oxidases, mitigation of oxidative stress caused by nitric oxide, increase in uric acid levels, and increase in antioxidant properties of low molecular antioxidants [116].

Based on the direct scavenging of ROS mechanism, flavonoids may directly scavenge free radicals by donating hydrogen atoms. The following equation is used to render radicals inactive; R^\bullet is a free radical, and FI-O^\bullet is a flavonoid phenoxyl radical [117].




Fig. 4 Scavenging of reactive oxygen species (R^\bullet) by flavonoid. The free radical FI-O^\bullet may react with a second radical, acquiring a stable quinone structure [117]

By activation of antioxidant enzymes approach, flavonoids can induce phase II detoxifying enzymes such NAD(P)H-quinone oxidoreductase, glutathione S-transferase, and UDP-glucuronosyl transferase, all of which are important defense enzymes against electrophilic toxicants and oxidative stress. Furthermore, some effects might be a resetting of a combination of radical scavenging and enzyme activity [116, 118]. Specific flavonoids possess the metal chelating activity and have been shown to chelate iron and copper, eliminating a potential source of free radicals. The catechol moiety in ring B, the 3-hydroxyl and 4-oxo groups in the heterocyclic ring C, and the 4-oxo and 5-hydroxyl groups in the C and A rings are all trace metal-binding sites in flavonoids [117].




Fig. 5. Binding sites for trace metals [117]

In cell membranes and human low-density lipoproteins (LDL), α -tocopherol is a crucial antioxidant that protects lipoprotein molecules from oxidative damage. Based on the reduction of α -tocopheryl radicals mechanism, the flavonoids can assist the α -tocopheryl radical, which is a potential prooxidant, by acting as hydrogen donors. They also have the ability to delay the oxidation of LDL by interacting with the α -tocopheryl radical [119]. Flavonoids can cause inhibition of oxidases, so that inhibiting the enzymes responsible for superoxide ($\text{O}_2 \cdot^-$) production, such as protein kinase C. Flavonoids have also been shown to inhibit cyclooxygenase, lipoxygenase, microsomal succinoxidase, and NADH oxidase [116]. Another oxidase, known as xanthine oxidase, is a molybdenum-containing

enzyme and converts xanthine to uric acid. The transfer of an oxygen atom from the molybdenum core to xanthine is known to occur in this enzyme-catalyzed process [120].

Mitigation of oxidative stress caused by nitric oxide is another way flavonoids apply their antioxidant activity. Although nitric oxide (NO) is necessary for sustaining blood vessel dilatation, excessive levels can cause oxidative damage. NO is created by NO synthases catalyzing the oxidation of L-arginine. In numerous lipopolysaccharide-activated cell lines and cultures, flavonoids inhibited NO generation. This impact was most likely induced by flavonoid suppression of inducible NOS expression rather than inhibition of its activity [116, 121]. An increase in uric acid levels can lead to a surge in plasma total antioxidant capacity. There are significant differences in plasma or serum total antioxidant capacity and plasma flavonoid concentrations. Increased uric acid levels, which are a key contributor to plasma total antioxidant capacity, are likely to be responsible for the considerable rise in plasma total antioxidant capacity found after eating flavonoid-rich meals [122]. An increase in antioxidant properties of low molecular antioxidants may occur by their amalgamation with flavonoids. UVA-induced cellular DNA damage was greatly reduced when flavonoids were combined with β -carotene during preincubation, and the effects were in the order of naringin > rutin > quercetin. According to the findings, combining β -carotene with naringin, rutin, or quercetin may improve the safety of β -carotene [123].

As mentioned before, one of the most important factors is affecting the antioxidant activity of flavonoids is the structure of flavonoids. The presence of hydroxyl and glycosylated groups has a significant effect on the antioxidant properties of flavonoids. The presence of glycosylated groups reduces the antioxidant properties of flavonoids. On the other hand, the presence of hydroxy groups increases the antioxidant properties of flavonoids. For maximum free radical scavenging activity, flavonoids must have a hydroxy group at the 4' and 3' positions of the B ring, a conjugated double bond between carbon 2 and 3, an Oxo group at the 4 position of the C ring, a hydroxy group on carbon 3 in the C ring, and a hydroxy group on carbon 5 in the A ring (refer to Fig. 2). Flavonoids, along with their hydroxy groups, act as free radical scavengers, and the presence of many hydroxy groups, especially in the B ring,

increases their antioxidant activity. The hydroxyls of the B ring are the first active stations in traversing the oxidation chain. The three structural groups are responsible for determining the scavenging activity of free radicals and antioxidant activity of flavonoids: a catechol section in the B ring, a conjugated double bond at the positions 2 and 3, an oxo group with the carbonyl group function in the ring C, and the presence of hydroxy groups at the positions 3 and 5 [115, 124]. Hence, it is noteworthy that flavonoids can exhibit different antioxidant power according to their structure and its relation to the suitable mechanism of the antioxidant activity.

Flavonoids are classified as non-nutrients in the scientific food realm. They have typically been eliminated from food crops because of their inhibitory effects on digestive enzymes, astringency and bitterness, and poor absorption after intake. However, as a result of their inclusion in regular meals, several therapeutic benefits such as antioxidant properties in animal studies, the decrease of cardiovascular disease and high blood pressure, and anti-allergic, anti-inflammatory, and anti-diabetic activities make them important. As a result, flavonoids are now considered third-order functional components (food factors) that have biological regulatory roles [125]. Flavonoids are also important in food preservation and industry. The phrase "sustainable intensification" was created by the Royal Society of London, which denotes current agricultural practices aiming at raising food production while also safeguarding biodiversity and environmental processes. Food waste minimization is one of the most critical elements of "sustainable intensification." As a result, there is a growing demand for postharvest storage technology and food preservatives research [126, 127]. Since today's consumers are concerned about the usage of chemically synthesized preservatives, natural products are being prioritized. At the same time, industries are putting a greater focus on the usage of plant-derived antioxidant and antibacterial components to improve shelf life. In this setting, the antioxidant activity of flavonoids may not only preserve food stability over time but also provide protection against necrotrophic fungi and food-borne diseases. The antioxidant mechanism of flavonoids in food systems is mainly connected to their capacity to prevent lipid autoxidation, which is a primary cause of food quality degradation and shelf-life reduction [128]. Flavonoids can donate hydrogen atoms to lipid radicals, resulting in more stable antioxidant




Fig. 6. The inhibition of the superoxide anion generation catalyzed by xanthine oxidase with antioxidants consisted of three reactions (A, B and C). [A] Inhibition of the xanthine oxidase reaction with antioxidants. [B] Reduction of the xanthine oxidase with antioxidants. [C] Scavenging of the superoxide anion with antioxidants [120]

radicals that are less susceptible to autoxidation. Antioxidant action mechanisms of flavonoids include direct scavenging of ROS, inhibition of ROS formation via chelation of trace elements (e.g., quercetin has iron-chelating and iron-stabilizing properties), or inhibition of free radical-generating enzymes (e.g., glutathione S-transferase, microsomal monooxygenase, mitochondrial succinoxidase, NADH oxidase), and activation of antioxidant defenses (e.g., upregulation of antioxidant enzymes with radical scavenging ability). A synergy of some of these mechanisms, such as radical scavenging activity along with some enzyme function inhibition, may also occur [129-131]. The capacity of flavonoids to lower the susceptibility of fresh fruits and vegetables to particular postharvest infections may also contribute to their antioxidant effect in extending shelf life [126, 132]. Flavonoids, exceptionally high hydroxylated anthocyanins, can prevent the growth of grey mold (produced by *Botrytis cinerea*), causing the dynamics of the ROS burst to be disrupted during infection. Flavonoids may bind and inactivate proteins and may form complexes with bacterial cell walls, giving them antibacterial action against a variety of pathogens [133-135]. Flavonoids have been employed in a variety of dietary applications due to their beneficial characteristics. They have been used as active antioxidant materials in the packaging of oxygen-sensitive foods to extend shelf life and improve bioactive component content [136]. They have also been used to minimize lipid oxidation in cooked pork patties and raw mackerel fillets [137]. Considering all

mentioned information regarding the antioxidant potential of flavonoids, it is evident that, besides their health-promoting effects, they can be utilized as food additives to extend their shelf life.

4. CONCLUSION

As one of the significant subgroups of polyphenols, flavonoids are crucial secondary chemicals known as phytochemicals generated by plants that serve a variety of activities, including growth and development and stress resistance. The high demand and interest in the utilization of flavonoids in food processing, as well as their health-promoting effects, has resulted in the increased acknowledgment of positive aspects of flavonoids for human health. It is evident that as an important subset of the phytochemicals family, flavonoids possess a considerable antioxidant activity that is competitive with synthetic antioxidants. Flavonoids are abundant in vegetables, flowers, and seeds, and approaches to extract these compounds from these natural sources have been developed for use as food additives and preservatives. However, on a commercial scale, the extraction of flavonoids directly from the mentioned sources does not seem to be profitable. Hence, it seems that attempting to find a proper source for extraction, such as plant wastes as well as designing a high-performance extraction method, would be a critical step for these compounds to be applied on food products as preservatives.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Atta EM, Mohamed NH, Silaev AA. Antioxidants: An overview on the natural and synthetic types. *European Chemical Bulletin*. 2017;6(8):365-75. Available: <https://doi.org/10.17628/ecb.2017.6.365-375>
- Latifi Z, Khesht SG, Daneshniya M. Detection of olive oil adulteration in terms of monitoring the U.V. and I.R. spectral patterns during thermal oxidation. 4th International Conference on Applied Researches in Science & Engineering At: Vrije Universiteit-Brussel; 2019. Available: <https://en.civilica.com/doc/1000635/>
- Halliwell B. Free radicals and antioxidants: updating a personal view. *Nutrition reviews*. 2012;70(5):257-65. Available: <https://doi.org/10.1111/j.1753-4887.2012.00476.x>
- Traystman RJ, Kirsch JR, Koehler RC. Oxygen radical mechanisms of brain injury following ischemia and reperfusion. *Journal of Applied Physiology*. 1991;71(4):1185-95. Available: <https://doi.org/10.1152/jappl.1991.71.4.1185>
- Nogala-Kalucka M, Korczak J, Dratwia M, Lampart-Szczapa E, Siger A, Buchowski M. Changes in antioxidant activity and free radical scavenging potential of rosemary extract and tocopherols in isolated rapeseed oil triacylglycerols during accelerated tests. *Food Chemistry*. 2005 1;93(2):227-35. Available: <https://doi.org/10.1016/j.foodchem.2004.09.021>
- Daneshniya M, Maleki MH, Liavali H, Hassanjani M, Keshavarz Bahadori N, Mohammadi M, Jalilvand Nezhad H. Antioxidant activity of flavonoids as an important phytochemical compound in plants. In Proceedings of the 2nd International Congress on Engineering, Technology and Innovation, Darmstadt, Germany; 2020.
- Daneshniya M, Maleki MH, Liavali HM, Hassanjani MR. Phenolic compounds extracted from plants as potential antioxidants. In Proceedings of the 2nd International Congress on Engineering, Technology and Innovation, Darmstadt, Germany 2020.
- Pokorný J. Are natural antioxidants better—and safer—than synthetic antioxidants?. *European journal of lipid science and technology*. 2007;109(6):629-42. Available: <https://doi.org/10.1002/ejlt.200700064>
- Halliwell B, Murcia MA, Chirico S, Aruoma OI. Free radicals and antioxidants in food and in vivo: what they do and how they work. *Critical Reviews in Food Science & Nutrition*. 1995;35(1-2):7-20. Available: <https://doi.org/10.1080/10408399509527682>
- Afanas' ev IB, Dcrozko AI, Brodskii AV, Kostyuk VA, Potapovitch AI. Chelating and free radical scavenging mechanisms of inhibitory action of rutin and quercetin in lipid peroxidation. *Biochemical pharmacology*. 1989 Jun 1;38(11):1763-9. Available: [https://doi.org/10.1016/0006-2952\(89\)90410-3](https://doi.org/10.1016/0006-2952(89)90410-3)
- latifi Y, Chaharlang M, Daneshniya M, khaki Arani S, Barzanooni M, Boghori P. Antioxidant and antimicrobial properties of methanolic extract of green walnut skin. *FSCT*. 2021;17(108):127-136. Available: <https://doi.org/10.52547/fsct.17.108.127>
- Lourenço SC, Moldão-Martins M, Alves VD. Antioxidants of natural plant origins: From sources to food industry applications. *Molecules*. 2019;24(22):4132. Available: <https://doi.org/10.3390/molecules24224132>
- Latifi Z, Daneshniya M, Khademi F. Effect of preservative of Anarijeh (*Froriepia subpinnata*) extract on shelf life of silver carp fish (*Hypophthalmichthys molitrix*) at superchilling temperature (-3 °C). In Proceedings of the 4th International Conference on Applied Researches in Science and Engineering; 2019. Available: <https://en.civilica.com/doc/1000639/>
- Ribeiro JS, Santos MJ, Silva LK, Pereira LC, Santos IA, da Silva Lannes SC, da Silva MV. Natural antioxidants used in meat products: A brief review. *Meat science*. 2019;1:148:181-8.
- Wojcik M, Burzynska-Pedziwiatr I, Wozniak LA. A review of natural and synthetic antioxidants important for health and longevity. *Current Medicinal Chemistry*. 2010;1;17(28):3262-88.

16. Dorman HD, Bachmayer O, Kosar M, Hiltunen R. Antioxidant properties of aqueous extracts from selected Lamiaceae species grown in Turkey. *Journal of Agricultural and Food Chemistry*. 2004;52(4):762-70.
Available:<https://doi.org/10.1021/jf034908v>
17. Ardabili AG, Farhoosh R, Khodaparast MH. Frying stability of canola oil in presence of pumpkin seed and olive oils. *European journal of lipid science and technology*. 2010;112(8):871-7.
Available:<https://doi.org/10.1002/ejlt.200900257>
18. Formanek Z, Kerry JP, Higgins FM, Buckley DJ, Morrissey PA, Farkas J. Addition of synthetic and natural antioxidants to α -tocopheryl acetate supplemented beef patties: effects of antioxidants and packaging on lipid oxidation. *Meat science*. 2001;58(4):337-41.
Available:[https://doi.org/10.1016/S0309-1740\(00\)00149-2](https://doi.org/10.1016/S0309-1740(00)00149-2)
19. Biswas AK, Keshri RC, Bisht G.S. Effect of enrobing and antioxidants on quality characteristics of precooked pork patties under chilled and frozen storage conditions. *Meat science*. 2004;66(3):733-41.
Available:<https://doi.org/10.1016/j.meatsci.2003.07.006>
20. Jayathilakan K, Sharma GK, Radhakrishna K, Bawa AS. Antioxidant potential of synthetic and natural antioxidants and its effect on warmed-over-flavour in different species of meat. *Food Chemistry*. 2007;105(3):908-16.
Available:<https://doi.org/10.1016/j.foodchem.2007.04.068>
21. Puri M, Sharma D, Barrow CJ. Enzyme-assisted extraction of bioactives from plants. *Trends in biotechnology*. 2012;30(1):37-44.
Available:<https://doi.org/10.1016/j.tibtech.2011.06.014>
22. Yadav RN, Agarwala M. Phytochemical analysis of some medicinal plants. *Journal of phytology*. 2011;3(12).
Available:<https://updatepublishing.com/journal/index.php/jp/article/view/2737>
23. Eliassen AH, Hendrickson SJ, Brinton LA, Buring JE, Campos H, Dai Q, Dorgan JF, Franke AA, Gao YT, Goodman MT, Hallmans G. Circulating carotenoids and risk of breast cancer: pooled analysis of eight prospective studies. *Journal of the National Cancer Institute*. 2012;104(24):1905-16.
Available:<https://doi.org/10.1093/jnci/djs461>
24. Pojer E, Mattivi F, Johnson D, Stockley CS. The case for anthocyanin consumption to promote human health: a review. *Comprehensive Reviews in Food Science and Food Safety*. 2013;12(5):483-508.
Available:<https://doi.org/10.1111/1541-4337.12024>
25. Tanaka T, Shnimizu M, Moriwaki H. Cancer chemoprevention by carotenoids. *Molecules*. 2012;17(3):3202-42.
Available:<https://doi.org/10.3390/molecules17033202>
26. Tokusoglu Ö, Hall III CA, editors. *Fruit and cereal bioactives: sources, chemistry, and applications*. 1st ed. CRC press; 2011.
Available:<https://doi.org/10.1201/b10786>
27. Latifi Z, Sadeghi F, Azizi J, Daneshniya M. Measuring of phytochemical and antioxidant compounds and identification of main phenolic compound in Anarijeh plant (*Froriepiasubpinnata*) extract by RPHPLC method. 13th International Conference on Engineering & Technology, Norway; 2019.
28. Bravo L. Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. *Nutrition reviews*. 1998;56(11):317-33.
Available:<https://doi.org/10.1111/j.1753-4887.1998.tb01670.x>
29. Daneshniya M, Maleki MH, Mohammadi MA, Ahangarian K, Kondeskalaei VJ, Alavi H. Antioxidant and Antimicrobial Activity of *Ferula* Species' Essential Oils and Plant Extracts and their Application as the Natural Food Preservatives. *South Asian Research Journal of Natural Products*. 2021;4(3):1-23.
Available:
<https://www.journalsarjnp.com/index.php/SARJNP/article/view/30133>
30. Heim KE, Tagliaferro AR, Bobilya DJ. Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *The Journal of nutritional biochemistry*. 2002;13(10):572-84.
Available:[https://doi.org/10.1016/S0955-2863\(02\)00208-5](https://doi.org/10.1016/S0955-2863(02)00208-5)
31. Amorati R, Valgimigli L. Modulation of the antioxidant activity of phenols by non-covalent interactions. *Organic & biomolecular chemistry*. 2012;10(21):4147-58.

- Available:<https://doi.org/10.1039/C2OB25174D>
32. Balasundram N, Sundram K, Samman S. Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence, and potential uses. *Food chemistry*. 2006;99(1):191-203. Available:<https://doi.org/10.1016/j.foodchem.2005.07.042>
 33. Harborne JB, Williams CA. Advances in flavonoid research since 1992. *Phytochemistry*. 2000; 55(6):481-504. Available:[https://doi.org/10.1016/S0031-9422\(00\)00235-1](https://doi.org/10.1016/S0031-9422(00)00235-1)
 34. Holtekjølen AK, Bævre AB, Rødbotten M, Berg H, Knutsen SH. Antioxidant properties and sensory profiles of breads containing barley flour. *Food chemistry*. 2008;110(2):414-21. Available:<https://doi.org/10.1016/j.foodchem.2008.02.054>
 35. Lam RY, Woo AY, Leung PS, Cheng CH. Antioxidant actions of phenolic compounds found in dietary plants on low-density lipoprotein and erythrocytes in vitro. *Journal of the American College of Nutrition*. 2007;26(3):233-42. Available:<https://doi.org/10.1080/07315724.2007.10719606>
 36. Tayel AA, EL-TRAS WF. Plant extracts as potent biopreservatives for *Salmonella typhimurium* control and quality enhancement in ground beef. *Journal of Food Safety*. 2012;32(1):115-21. Available:<https://doi.org/10.1111/j.1745-4565.2011.00357.x>
 37. Devatkal SK, Narsaiah K, Borah A. Antioxidant effect of extracts of kinnow rind, pomegranate rind and seed powders in cooked goat meat patties. *Meat Science*. 2010;85(1):155-9. Available:<https://doi.org/10.1016/j.meatsci.2009.12.019>
 38. Huang B, He J, Ban X, Zeng H, Yao X, Wang Y. Antioxidant activity of bovine and porcine meat treated with extracts from edible lotus (*Nelumbo nucifera*) rhizome knot and leaf. *Meat science*. 2011;87(1):46-53. Available:<https://doi.org/10.1016/j.meatsci.2010.09.001>
 39. Wójciak KM, Dolatowski ZJ, Okoń A. The effect of water plant extracts addition on the oxidative stability of meat products. *Acta Scientiarum Polonorum Technologia Alimentaria*. 2011;10(2):175-88. Available:https://www.food.actapol.net/pub/4_2_2011.pdf
 40. Das AK, Rajkumar V, Verma AK, Swarup D. Moringa oleifera leaves extract: a natural antioxidant for retarding lipid peroxidation in cooked goat meat patties. *International journal of food science & technology*. 2012;47(3):585-91. Available:<https://doi.org/10.1111/j.1365-2621.2011.02881.x>
 41. Latifi Z, Daneshniya M, Khademi F. Effect of preservative of Anarijeh (*Froriepia subpinnata*) extract on shelf life of silver carp fish (*Hypophthalmichthys molitrix*) at superchilling temperature (-3 C). In *Proceedings of the 4th International Conference on Applied Researches in Science and Engineering*;2019. Available:<https://en.civilica.com/doc/1000639/>
 42. Ebadi M, Latifi Z, Daneshniya M. Optimization of the Antioxidant Effect of Ethanolic Extract of Thistle (*Carduus pycnocephalus* L.) by Response Surface Method and Comparison of the Antioxidant Effect of Extract and Essential Oil on Oxidative Soybean Oil Resistance. *Journal of food science and technology (Iran)*. 2021;17(108):151-64. Available:<https://doi.org/10.52547/fsct.17.108.151>
 43. Cheynier V. Phenolic compounds: from plants to foods. *Phytochemistry reviews*. 2012;11(2):153-77. Available:<https://doi.org/10.1007/s11101-012-9242-8>
 44. Singleton VL. Naturally occurring food toxicants: phenolic substances of plant origin common in foods. *Advances in food research*. 1981;27:149-242. Available:[https://doi.org/10.1016/S0065-2628\(08\)60299-2](https://doi.org/10.1016/S0065-2628(08)60299-2)
 45. Su TR, Lin JJ, Tsai CC, Huang TK, Yang ZY, Wu MO, Zheng YQ, Su CC, Wu YJ. Inhibition of melanogenesis by gallic acid: Possible involvement of the PI3K/Akt, MEK/ERK and Wnt/ β -catenin signaling pathways in B16F10 cells. *International journal of molecular sciences*. 2013;14(10):20443-58. Available:<https://doi.org/10.3390/ijms141020443>
 46. Boz H. p-Coumaric acid in cereals: presence, antioxidant and antimicrobial effects. *International Journal of Food Science & Technology*. 2015;50(11): 2323-8.

- Available:<https://doi.org/10.1111/ijfs.12898>
47. Marchev AS, Vasileva LV, Amirova KM, Savova MS, Koycheva IK, Balcheva-Sivenova ZP, Vasileva SM, Georgiev MI. Rosmarinic acid-From bench to valuable applications in food industry. *Trends in Food Science & Technology*. 2021;117:182-93. Available:<https://doi.org/10.1016/j.tifs.2021.03.015>
 48. Magnani C, Isaac VL, Correa MA, Salgado HR. Caffeic acid: a review of its potential use in medications and cosmetics. *Analytical Methods*. 2014;6(10):3203-10. Available:<https://doi.org/10.1039/C3AY41807C>
 49. Kakkar S, Bais S. A review on protocatechuic acid and its pharmacological potential. *International Scholarly Research Notices*. 2014;2014. Available:<https://doi.org/10.1155/2014/952943>
 50. Andrade JM, Faustino C, Garcia C, Ladeiras D, Reis CP, Rijo P. *Rosmarinus officinalis* L.: an update review of its phytochemistry and biological activity. *Future science OA*. 2018;4(4):FSO283. Available:<https://doi.org/10.4155/fsoa-2017-0124>
 51. Sugihara N, Ohnishi M, Imamura M, Furuno K. Differences in antioxidative efficiency of catechins in various metal-induced lipid peroxidations in cultured hepatocytes. *Journal of health science*. 2001;47(2):99-106. Available:<https://doi.org/10.1248/jhs.47.99>
 52. Sivakanthan S, Rajendran S, Gamage A, Madhujith T, Mani S. Antioxidant and antimicrobial applications of biopolymers: A review. *Food Research International*. 2020;136:109327. Available:<https://doi.org/10.1016/j.foodres.2020.109327>
 53. Cavia-Saiz M, Busto MD, Pilar-Izquierdo MC, Ortega N, Perez-Mateos M, Muñoz P. Antioxidant properties, radical scavenging activity and biomolecule protection capacity of flavonoid naringenin and its glycoside naringin: A comparative study. *Journal of the Science of Food and Agriculture*. 2010 May;90(7):1238-44. Available:<https://doi.org/10.1002/jsfa.3959>
 54. Baghel SS, Shrivastava N, Baghel RS, Agrawal P, Rajput S. A review of quercetin: antioxidant and anticancer properties. *World J Pharm Pharmaceutical Sci*. 2012;1(1):146-60. Available:https://www.researchgate.net/publication/267333748_A_review_of_quercetin_Antioxidant_and_anticancer_properties
 55. Kashyap P, Shikha D, Thakur M, Aneja A. Functionality of apigenin as a potent antioxidant with emphasis on bioavailability, metabolism, action mechanism and in vitro and in vivo studies: A review. *Journal of food biochemistry*. 2021 Sep 26:e13950. Available:<https://doi.org/10.1111/jfbc.13950>
 56. Miguel MG. Anthocyanins: Antioxidant and/or anti-inflammatory activities. *Journal of Applied Pharmaceutical Science*. 2011;1(6):7-15. Available:<http://hdl.handle.net/10400.1/6317>
 57. Weng L, Zhang F, Wang R, Ma W, Song Y. A review on protective role of genistein against oxidative stress in diabetes and related complications. *Chemico-biological interactions*. 2019;310:108665. Available:<https://doi.org/10.1016/j.cbi.2019.05.031>
 58. Nieto G, Ros G, Castillo J. Antioxidant and antimicrobial properties of rosemary (*Rosmarinus officinalis*, L.): A review. *Medicines*. 2018;5(3):98. Available:<https://doi.org/10.3390/medicines5030098>
 59. Yilmaz Y. Novel uses of catechins in foods. *Trends in Food Science & Technology*. 2008;17(2):64-71. Available:<https://doi.org/10.1016/j.tifs.2005.10.005>
 60. He J, Xu L, Le Yang XW. Epigallocatechin gallate is the most effective catechin against antioxidant stress via hydrogen peroxide and radical scavenging activity. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research*. 2018;24:8198. Available:<https://doi.org/10.12659/MSM.911175>
 61. Hu C, Kitts DD. Evaluation of antioxidant activity of epigallocatechin gallate in biphasic model systems in vitro. *Molecular and Cellular Biochemistry*. 2001;218(1):147-55. Available:<https://doi.org/10.1023/A:1007220928446>
 62. Yang J, Guo J, Yuan J. In vitro antioxidant properties of rutin. *LWT-Food Science and Technology*. 2008;41(6):1060-6. Available:<https://doi.org/10.1016/j.lwt.2007.06.010>

63. Gülçin İ. Antioxidant activity of eugenol: A structure–activity relationship study. *Journal of medicinal food*. 2011;14(9):975-85.
Available:<https://doi.org/10.1089/jmf.2010.0197>
64. Aristatile B, Al-Numair KS, Al-Assaf AH, Veeramani C, Pugalendi KV. Protective effect of carvacrol on oxidative stress and cellular DNA damage induced by UVB irradiation in human peripheral lymphocytes. *Journal of biochemical and molecular toxicology*. 2015;29(11):497-507.
Available:<https://doi.org/10.1002/jbt.20355>
65. Viuda-Martos M, Ruiz Navajas Y, Sánchez Zapata E, Fernández-López J, Pérez-Álvarez JA. Antioxidant activity of essential oils of five spice plants widely used in a Mediterranean diet. *Flavour and Fragrance Journal*. 2010 Jan;25(1):13-9.
Available:<https://doi.org/10.1002/ffj.1951>
66. Eid AM, Hawash M. Biological evaluation of Safrole oil and Safrole oil Nanoemulgel as antioxidant, antidiabetic, antibacterial, antifungal and anticancer. *BMC Complementary Medicine and Therapies*. 2021 Dec;21(1):1-2.
Available:<https://doi.org/10.1186/s12906-021-03324-z>
67. Wu Z, Tan B, Liu Y, Dunn J, Martorell Guerola P, Tortajada M, Cao Z, Ji P. Chemical composition and antioxidant properties of essential oils from peppermint, native spearmint and scotch spearmint. *Molecules*. 2019;24(15):2825.
Available:<https://doi.org/10.3390/molecules24152825>
68. Sun Z, Wang H, Wang J, Zhou L, Yang P. Chemical composition and anti-inflammatory, cytotoxic and antioxidant activities of essential oil from leaves of *Mentha piperita* grown in China. *PloS one*. 2014 Dec 10;9(12):e114767.
Available:<https://doi.org/10.1371/journal.pone.0114767>
69. Lim H, Shin S. Antibacterial and Antioxidant Activities of the Essential Oil from the Roots of *Anthriscus sylvestris*. *Yakhak Hoeji*. 2012;56(5):320-5.
Available:<https://www.koreascience.or.kr/article/JAKO201217642691877.page>
70. Mimica-Dukić N, Božin B, Soković M, Mihajlović B, Matavulj M. Antimicrobial and antioxidant activities of three *Mentha* species essential oils. *Planta medica*. 2003;69(05):413-9.
Available:<https://doi.org/10.1055/s-2003-39704>
71. Larayetan RA, Okoh OO, Sadimenko A, Okoh AI. Terpene constituents of the aerial parts, phenolic content, antibacterial potential, free radical scavenging and antioxidant activity of *Callistemon citrinus* (Curtis) Skeels (Myrtaceae) from Eastern Cape Province of South Africa. *BMC complementary and alternative medicine*. 2017;17(1):1-9.
Available:<https://doi.org/10.1186/s12906-017-1804-2>
72. Chaudhari AK, Singh A, Singh VK, Dwivedy AK, Das S, Ramsdam MG, Dkhar MS, Kayang H, Dubey NK. Assessment of chitosan biopolymer encapsulated α -Terpineol against fungal, aflatoxin B1 (AFB1) and free radicals mediated deterioration of stored maize and possible mode of action. *Food chemistry*. 2020;311:126010.
Available:<https://doi.org/10.1016/j.foodchem.2019.126010>
73. Reza M, Nejad S. Antibacterial and Antioxidant Activity and Essential Oil Composition of *Grammosciadium scabridum* Boiss. from Iran. *Zeitschrift für Naturforschung C*;2014.
74. Subash-Babu P, Alshatwi AA, Ignacimuthu S. Beneficial antioxidative and antiperoxidative effect of cinnamaldehyde protect streptozotocin-induced pancreatic β -cells damage in wistar rats. *Biomolecules & therapeutics*. 2014; 22(1):47.
Available:<https://doi.org/10.4062/biomolther.2013.100>
75. Carp OE, Moraru A, Pinteala M, Arvinte A. Electrochemical behaviour of piperine. Comparison with control antioxidants. *Food Chemistry*. 2021;339:128110.
Available:<https://doi.org/10.1016/j.foodchem.2020.128110>
76. Alamed J, Chaiyasit W, McClements DJ, Decker EA. Relationships between free radical scavenging and antioxidant activity in foods. *Journal of agricultural and food chemistry*. 2009;57(7):2969-76.
Available:<https://doi.org/10.1021/jf803436c>
77. Hussein MA. A convenient mechanism for the free radical scavenging activity of resveratrol. *International Journal of Phytomedicine*. 2011;3(4):459.
Available:<https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.868.4033&rep=rep1&type=pdf>

78. Queenthy SS, John B. Diosmin exhibits anti-hyperlipidemic effects in isoproterenol induced myocardial infarcted rats. *European journal of pharmacology*. 2013;718(1-3):213-8. Available:<https://doi.org/10.1016/j.ejphar.2013.08.031>
79. Chu YH, Chang CL, Hsu HF. Flavonoid content of several vegetables and their antioxidant activity. *Journal of the Science of Food and Agriculture*. 2000;80(5):561-6. Available:[https://doi.org/10.1002/\(SICI\)1097-0010\(200004\)80:5<561::AID-JSFA574>3.0.CO;2-%23](https://doi.org/10.1002/(SICI)1097-0010(200004)80:5<561::AID-JSFA574>3.0.CO;2-%23)
80. Pietta PG. Flavonoids as antioxidants. *Journal of natural products*. 2000;63(7):1035-42. Available:<https://doi.org/10.1021/np9904509>
81. Ververidis F, Trantas E, Douglas C, Vollmer G, Kretzschmar G, Panopoulos N. Biotechnology of flavonoids and other phenylpropanoid-derived natural products. Part I: Chemical diversity, impacts on plant biology and human health. *Biotechnology Journal: Healthcare Nutrition Technology*. 2007;2(10):1214-34. Available:<https://doi.org/10.1002/biot.200700084>
82. Laura A, Moreno-Escamilla JO, Rodrigo-García J, Alvarez-Parrilla E. Phenolic compounds. In *Postharvest physiology and biochemistry of fruits and vegetables*. Woodhead Publishing. 2019;253-271. Available:<https://doi.org/10.1016/B978-0-12-813278-4.00012-9>
83. Borges Bubols G, da Rocha Vianna D, Medina-Reimon A, von Poser G, Maria Lamuela-Raventos R, Lucia Eiffler-Lima V, Cristina Garcia S. The antioxidant activity of coumarins and flavonoids. *Mini Reviews in Medicinal Chemistry*. 2013;13(3):318-34.
84. Banjarnahor SD, Artanti N. Antioxidant properties of flavonoids. *Medical Journal of Indonesia*. 2014;23(4):239-44.
85. Amić D, Davidović-Amić D, Beslo D, Rastija V, Lucić B, Trinajstić N. SAR and QSAR of the antioxidant activity of flavonoids. *Current Medicinal Chemistry*. 2007;14(7):827-45.10.
86. Dugas Jr AJ, Castañeda-Acosta J, Bonin GC, Price KL, Fischer NH, Winston GW. Evaluation of the total peroxy radical-scavenging capacity of flavonoids: structure– activity relationships. *Journal of Natural Products*. 2000;26;63(3):327-31.
87. Santos MR, Mira L. Protection by flavonoids against the peroxy-nitrite-mediated oxidation of dihydrorhodamine. *Free Radical Research*. 2004;1;38(9):1011-8.
88. Moalin M, Van Strijdonck GP, Beckers M, Hagemen GJ, Borm PJ, Bast A, Haenen GR. A planar conformation and the hydroxyl groups in the B and C rings play a pivotal role in the antioxidant capacity of quercetin and quercetin derivatives. *Molecules*. 2011;16(11):9636-50.
89. Gutiérrez-del-Río I, López-Ibáñez S, Magadán-Corpas P, Fernández-Calleja L, Pérez-Valero Á, Tuñón-Granda M, Miguélez EM, Villar CJ, Lombó F. Terpenoids and polyphenols as natural antioxidant agents in food preservation. *Antioxidants*. 2021;10(8):1264. Available:<https://doi.org/10.3390/antiox10081264>
90. Manassis G, Kalogianni AI, Lazou T, Moschovas M, Bossis I, Gelasakis AI. Plant-derived natural antioxidants in meat and meat products. *Antioxidants*. 2020;9(12):1215. Available:<https://doi.org/10.3390/antiox9121215>
91. Carocho M, Morales P, Ferreira IC. Antioxidants: Reviewing the chemistry, food applications, legislation and role as preservatives. *Trends in Food Science & Technology*. 2018;71:107-20. Available:<https://doi.org/10.1016/j.tifs.2017.11.008>
92. European Parliament and Council Commission Regulation (E.U.) No 1129/2011 of 11 November 2011 Amending Annex II to Regulation (E.C.) No 1333/2008 of the European Parliament and of the Council by Establishing a Union List of Food Additives (Text with EEA Relevance). Available: <https://op.europa.eu/en/publication-detail/-/publication/28cb4a37-b40e-11e3-86f9-01aa75ed71a1>
93. Oswell NJ, Thippareddi H, Pegg RB. Practical use of natural antioxidants in meat products in the U.S.: A review. *Meat science*. 2018;145:469-79. Available:<https://doi.org/10.1016/j.meatsci.2018.07.020>
94. FDA US. (Food and Drug Administration). CFR—Code of Federal Regulations Title 21 Part 172. Food Aditives Permitted for

- Direct Addition to Food for Human Consumption.
Available:<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?CFRPart=172>
95. FDA US. (Food and Drug Administration). CFR—Code of Federal Regulations Title 21 Part 182. Substances Generally Recognized as Safe.
Available:<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfCFR/CFRSearch.cfm?CFRPart=182> (accessed on 15 July 2021).
 96. FDA U.S. (Food and Drug Administration). Substances Added to Food (Formerly EAFUS). Available online: <https://www.cfsanappsexternal.fda.gov/scripts/fdcc/?set=FoodSubstances> (accessed on 10 January 2022).
 97. European Commission Food Additives Database.
Available:https://www.cfsanappsexternal.fda.gov/scripts/fdcc/?set=FoodSubstances&sort=Sortterm_ID&order=ASC&startrow=1&type=basic&search=ASCORBIC (accessed on 17 January 2022).
 98. Barden L, Decker EA. Lipid oxidation in low-moisture food: A review. *Critical reviews in food science and nutrition*. 2016;56(15):2467-82.
Available:<https://doi.org/10.1080/10408398.2013.848833>
 99. Yang X, Sun Z, Wang W, Zhou Q, Shi G, Wei F, Jiang G. Developmental toxicity of synthetic phenolic antioxidants to the early life stage of zebrafish. *Science of the Total Environment*. 2018;643:559-68.
 100. Lin D, Xiao M, Zhao J, Li Z, Xing B, Li X, Kong M, Li L, Zhang Q, Liu Y, Chen H. An overview of plant phenolic compounds and their importance in human nutrition and management of type 2 diabetes. *Molecules*. 2016 Oct;21(10):1374.
Available:<https://doi.org/10.3390/molecules21101374>
 101. Kwan YH, Tung YK, Kochhar JS, Li H, Poh AL, Kang L. 2-Regulation of cosmetics. *Handbook of Cosmeceutical Excipients and their Safeties*, Yu Heng Kwan, Yee Kei Tung, Jaspreet Singh Kochhar, Hairui Li, Ai-Ling Poh, and Lifeng Kang (Eds.). Woodhead Publishing. 2014:7-22.
Available:<https://www.elsevier.com/books/handbook-of-cosmeceutical-excipients-and-their-safeties/kwan/978-1-907568-53-4>
 102. EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS). Scientific Opinion on the re-evaluation of butylated hydroxyanisole–BHA (E 320) as a food additive. *EFSA Journal*. 2011;9(10):2392.
Available:<https://doi.org/10.2903/j.efsa.2011.2392>
 103. EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS). Scientific Opinion on the re-evaluation of butylated hydroxytoluene BHT (E 321) as a food additive. *EFSA Journal*. 2012;10(3):2588.
Available:<https://doi.org/10.2903/j.efsa.2012.2588>
 104. EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS). Statement on the refined exposure assessment of tertiary-butyl hydroquinone (E 319). *EFSA Journal*. 2016;14(1):4363.
Available:<https://doi.org/10.2903/j.efsa.2016.4363>
 105. EFSA Panel on Food Additives and Nutrient Sources Added to Food (ANS). Scientific Opinion on the re-evaluation of propyl gallate (E 310) as a food additive. *EFSA Journal*. 2014;12(4):3642.
Available:<https://doi.org/10.2903/j.efsa.2014.3642>
 106. EFSA Panel on Food additives and Nutrient Sources added to Food (ANS). Scientific Opinion on the re-evaluation of ascorbic acid (E 300), sodium ascorbate (E 301) and calcium ascorbate (E 302) as food additives. *Efsa Journal*. 2015;13(5):4087.
Available:<https://doi.org/10.2903/j.efsa.2015.4087>
 107. EFSA Panel on Food Additives and Nutrient Sources added to food (ANS). Scientific Opinion on the re-evaluation of erythorbic acid (E 315) and sodium erythorbate (E 316) as food additives. *EFSA Journal*. 2016;14(1):4360.
Available:<https://doi.org/10.2903/j.efsa.2016.4360>
 108. EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS). Scientific Opinion on the re-evaluation of ascorbyl palmitate (E 304 (i)) and ascorbyl stearate (E 304 (ii)) as food additives. *EFSA Journal*. 2015;13(11):4289.
Available:<https://doi.org/10.2903/j.efsa.2015.4289>
 109. EFSA Panel on Food additives and Nutrient Sources added to Food (ANS).

- Scientific Opinion on the re-evaluation of sulfur dioxide (E 220), sodium sulfite (E 221), sodium bisulfite (E 222), sodium metabisulfite (E 223), potassium metabisulfite (E 224), calcium sulfite (E 226), calcium bisulfite (E 227) and potassium bisulfite (E 228) as food additives. *Efsa Journal*. 2016;14(4): 4438.
Available:<https://doi.org/10.2903/j.efsa.2016.4438>
110. EFSA Panel on Dietetic Products, Nutrition, and Allergies (NDA). Scientific Opinion on Dietary Reference Values for vitamin E as α -tocopherol. *EFSA Journal*. 2015;13(7):4149.
Available:<https://doi.org/10.2903/j.efsa.2015.4149>
 111. Lourenço SC, Moldão-Martins M, Alves VD. Antioxidants of natural plant origins: From sources to food industry applications. *Molecules*. 2019;24(22):4132.
Available:<https://doi.org/10.3390/molecules24224132>
 112. Xu X, Liu A, Hu S, Ares I, Martínez-Larrañaga MR, Wang X, Martínez M, Anadón A, Martínez MA. Synthetic phenolic antioxidants: Metabolism, hazards and mechanism of action. *Food Chemistry*. 2021;353:129488.
Available:<https://doi.org/10.1016/j.foodchem.2021.129488>
 113. Liu R, Mabury SA. Synthetic phenolic antioxidants: A review of environmental occurrence, fate, human exposure, and toxicity. *Environmental science & technology*. 2020;54(19):11706-19.
Available:<https://doi.org/10.1021/acs.est.0c05077>
 114. Baran A, Yildirim S, Ghosigharehaghaji A, Bolat İ, Sulukan E, Ceyhun SB. An approach to evaluating the potential teratogenic and neurotoxic mechanism of BHA based on apoptosis induced by oxidative stress in zebrafish embryo (*Danio rerio*). *Human & experimental toxicology*. 2021;40(3):425-38.
Available:<https://doi.org/10.1177/0960327120952140>
 115. Gülçin I. Antioxidant activity of food constituents: an overview. *Archives of toxicology*. 2012;86(3):345-91.
Available:<https://doi.org/10.1007/s00204-011-0774-2>
 116. Procházková D, Boušová I, Wilhelmová N. Antioxidant and prooxidant properties of flavonoids. *Fitoterapia*. 2011;82(4):513-23.
 117. Pietta PG. Flavonoids as antioxidants. *Journal of natural products*. 2000 Jul 28;63(7):1035-42.
 118. Nerland DE. The antioxidant/electrophile response element motif. *Drug metabolism reviews*. 2007;39(1):235-48.
 119. Hirano R, Sasamoto W, Matsumoto A, Itakura H, Igarashi O, Kondo K. Antioxidant ability of various flavonoids against DPPH radicals and LDL oxidation. *Journal of nutritional science and vitaminology*. 2001;47(5):357-62.
 120. Masuoka N, Matsuda M, Kubo I. Characterisation of the antioxidant activity of flavonoids. *Food chemistry*. 2012;131(2):541-5.
 121. Olszanecki R, Gebaska A, Kozlovski VI, Gryglewski RJ. Key words: flavonoids, nitric oxide synthase, macrophages. *Journal of Physiology and Pharmacology*. 2002;53(4):571-84.
 122. Lotito SB, Frei B. Consumption of flavonoid-rich foods and increased plasma antioxidant capacity in humans: cause, consequence, or epiphenomenon?. *Free Radical Biology and Medicine*. 2006;41(12):1727-46.
 123. Yeh SL, Wang WY, Huang CH, Hu ML. Pro-oxidative effect of β -carotene and the interaction with flavonoids on UVA-induced DNA strand breaks in mouse fibroblast C3H10T1/2 cells. *The Journal of nutritional biochemistry*. 2005;16(12):729-35.
 124. Heijnen CG, Haenen GR, Van Acker FA, Van der Vijgh WJ, Bast A. Flavonoids as peroxynitrite scavengers: the role of the hydroxyl groups. *Toxicology In vitro*. 2001;15(1):3-6.
Available:[https://doi.org/10.1016/S0887-2333\(00\)00053-9](https://doi.org/10.1016/S0887-2333(00)00053-9)
 125. Terahara N. Flavonoids in foods: a review. *Natural product communications*. 2015;10(3):1934578X1501000334.
Available:<https://doi.org/10.1177/1934578X1501000334>
 126. D'Amelia V, Aversano R, Chiaiese P, Carputo D. The antioxidant properties of plant flavonoids: their exploitation by molecular plant breeding. *Phytochemistry Reviews*. 2018;17(3):611-25.
Available:<https://doi.org/10.1007/s11101-018-9568-y>
 127. Petersen B, Snapp S. What is sustainable intensification? Views from experts. *Land use policy*. 2015;46:1-0.
Available:<https://doi.org/10.1016/j.landusepol.2015.02.002>

128. Shahidi F, Ambigaipalan P. Phenolics and polyphenolics in foods, beverages and spices: Antioxidant activity and health effects—A review. *Journal of functional foods*. 2015;18:820-97. Available:<https://doi.org/10.1016/j.jff.2015.06.018>
129. Kumar S, Pandey AK. Chemistry and biological activities of flavonoids: an overview. *The scientific world journal*. 2013 Oct;2013. <https://doi.org/10.1155/2013/162750>
130. Agrawal AD. Pharmacological activities of flavonoids: A review. *International Journal of Pharmaceutical Sciences and Nanotechnology*. 2011;4(2):1394-8. Available:<https://doi.org/10.37285/ijpsn.2011.4.2.3>
131. Kaleem M, Ahmad A. Flavonoids as nutraceuticals. In *Therapeutic, probiotic, and unconventional foods 2018*;137-155. Academic Press. Available:<https://doi.org/10.1016/B978-0-12-814625-5.00008-X>
132. Neha P, Pandey-Rai S. Biochemical activity and therapeutic role of antioxidants in plants and humans. *Plants as a Source of Natural Antioxidants*; CABI Publishing: Wallingford, UK. 2014;191-224. Available:https://www.researchgate.net/publication/267395391_Biochemical_Activity_and_Therapeutic_Role_of_Antioxidants_in_Plants_and_Humans
133. Zhang Y, Butelli E, De Stefano R, Schoonbeek HJ, Magusin A, Pagliarani C, et al. Anthocyanins double the shelf life of tomatoes by delaying overripening and reducing susceptibility to gray mold. *Current Biology*. 2013;23(12):1094-100. Available:<https://doi.org/10.1016/j.cub.2013.04.072>
134. Zhang Y, De Stefano R, Robine M, Butelli E, Bulling K, Hill L, Rejzek M, Martin C, Schoonbeek HJ. Different reactive oxygen species scavenging properties of flavonoids determine their abilities to extend the shelf life of tomato. *Plant Physiology*. 2015;169(3):1568-83. Available:<https://doi.org/10.1104/pp.15.00346>
135. Hintz T, Matthews KK, Di R. The use of plant antimicrobial compounds for food preservation. *BioMed research international*. 2015;2015. Available:<https://doi.org/10.1155/2015/246264>
136. López de Dicastillo C, Nerín C, Alfaro P, Catalá R, Gavara R, Hernández-Muñoz P. Development of new antioxidant active packaging films based on ethylene vinyl alcohol copolymer (EVOH) and green tea extract. *Journal of agricultural and food chemistry*. 2011;59(14):7832-40. Available:<https://doi.org/10.1021/jf201246g>
137. Viji P, Binsi PK, Visnuvinayagam S, Bindu J, Ravishankar CN, Srinivasa Gopal TK. Efficacy of mint (*Mentha arvensis*) leaf and citrus (*Citrus aurantium*) peel extracts as natural preservatives for shelf life extension of chill stored Indian mackerel. *Journal of food science and technology*. 2015;52(10):6278-89. Available:<https://doi.org/10.1007/s13197-015-1788-1>

© 2022 Maleki et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/84048>