



Influence of Spacing and Zinc Application on Growth and Productivity of Baby Corn (*Zea mays* L.)

**Gopathi Rohini^{a*}, Vikram Singh^{a#}, Shruti G. George^{a‡}
and Shrish Kumar Singh^{b†}**

^a Department of Agronomy, Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh, India.

^b Department of Agronomy, TDPG College, Jaunpur, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i1731044

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86231>

Received 12 February 2022

Accepted 20 April 2022

Published 30 April 2022

Original Research Article

ABSTRACT

The present trial was undertaken at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, Uttar Pradesh during the 2021 summer season (*Zaid*). The test was a Randomized Block Design having three replicates. A total of nine treatments were designed with different levels of spacing and quantity of zinc application. The cultivation land was uniform with sandy loam soil with pH neutral. The soil had low level of organic carbon (0.72%), medium level of available nitrogen (278.28 kg ha⁻¹) and potassium (233.24 kg ha⁻¹) and higher level of phosphorus (27.80 kg ha⁻¹). Amongst all the treatments, T₉ with 60 X 20 cm spacing and 25 kgha⁻¹ZnSO₄ had the maximum plant height (168.13 cm), number of leaves per plant (13.25), dry plant mass (90.96 g/plant), number of cobs per plant (2.37), length of the cob per plant (18.77 cm), cob weight with husk (47.92 g), cob weight without husk (22.70 g). Further, the maximum crop growth rate (30.45 g/m²/day), cob yield with husk (14.63 t/ha), cob yield without husk (5.09 t/ha) and green fodder yield (28.83 t/ha) was observed for T₃ with 40 X 20 cm spacing and 20 kg ha⁻¹ZnSO₄.

Keywords: *Baby corn; zinc application; productivity; spacing.*

[‡]M. Sc. Scholar;

[#]Associate Professor;

[‡]Ph.D. Scholar;

[†]Associate Professor and Head;

*Corresponding author: E-mail: gopathirohini445@gmail.com;

1. INTRODUCTION

Baby corn has become one of the most sought-after crops at the global level with great processing and export possibilities. Known by different names such as young corn, mini corn or candle corn, baby corn is maize (*Zea mays* L.) cob or ears harvested early from the female flower without any fertilization while the stalks are undeveloped. At present, China and Thailand are the leading producers and its widespread use has increased in India [1] as well since it can be grown at any point in the year and marketable as a vegetable crop [2]. Different regions of Meghalaya, Uttar Pradesh, Maharashtra, Karnataka, Andhra Pradesh and Haryana are the major players in the production of baby corn [3].

Based on the endosperm of grain, maize is defined into eight groups of which baby corn is utilized for vegetable purposes. It is the immature maize cob or ear harvested before the pollination of the flower or when there is surfacing of the baby corn silk. For better marketability, yellow-coloured cobs with row array being regular having 10-12 cm spacing and 1-1.5 cm broadness is preferential. Globally, this crop is harvested in Thailand, Taiwan, Sri Lanka, Myanmar, Guatemala and South Africa. In India, for a growth area of 8.49 m per hectare, the production and yield rate are 21.28 m per tonnes and 2507 kg per hectare respectively. Diversification to this high-value crop is beneficial to the Indian farmers due to low financial risk and economic wellbeing [4].

The metabolic reactions in plants are driven by an important mineral, Zinc. Zinc deficiency results in poor production of chlorophyll, carbohydrates, proteins, auxins and hindrance in growth and development of maize [5]. It has an indispensable part in regulation of RNA and DNA structure and certain enzymes such as hydrogenase and carbonic anhydrase [6].

The plant yield is affected based on the density population since optimum spacing promotes resourceful usage of resources such as water, soil, nutrients and sunlight [7]. Optimal growth is obtained through ideal spacing though the productivity results vary depending on cultivar and environment [8]. Plants in closed spacing results in overcrowding and poor yield and plant growth [9].

1.1 Objectives

1. To study the influence of spacing and different levels of zinc on growth and productivity of baby corn.
2. To work out the economics of different treatment combinations.

2. MATERIALS AND METHODS

The present trial was undertaken using Super Goldy variety at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, Uttar Pradesh (25.57°N, 87.19°E, 98 m) during the 2021 summer season (*Zaid*). The test was a Randomized Block Design having three replicates. Different levels of spacing and quantity of zinc application were set and nine treatments were obtained- T₁: 15 kg ha⁻¹ ZnSO₄ + 40 cm x 20 cm, T₂: 20 kg ha⁻¹ ZnSO₄ + 40 cm x 20 cm, T₃: 25 kg ha⁻¹ ZnSO₄ + 40 cm x 20 cm, T₄: 15 kg ha⁻¹ ZnSO₄ + 50 cm x 20 cm, T₅: 20 kg ha⁻¹ ZnSO₄ + 50 cm x 20 cm, T₆: 25 kg ha⁻¹ ZnSO₄ + 50 cm x 20 cm, T₇: 15 kg ha⁻¹ ZnSO₄ + 60 cm x 20 cm, T₈: 20 kg ha⁻¹ ZnSO₄ + 60 cm x 20 cm, T₉: 25 kg ha⁻¹ ZnSO₄ + 60 cm x 20 cm.

The cultivation land was uniform with sandy loam soil with pH neutral. The soil had low level of organic carbon (0.72%), medium level of available nitrogen (278.28 kg ha⁻¹) and potassium (233.24 kg ha⁻¹) and higher level of phosphorus (27.80 kg ha⁻¹). The fertilizers used included urea, DAP and MOP at basal level during sowing. The data was recorded for the following growth parameters during harvesting: plant height, number of leaves and dry plant mass. The yield parameters included: number of cobs per plant, cob length (cm), cob mass, green cob productivity (t/ha), green fodder yield (t/ha). The collected data was subjected to analysis of variance (ANOVA) using --- (Gomez K.A. and Gomez A.A. 1984).

3. RESULTS AND DISCUSSION

3.1 Growth Attributes

3.1.1 Plant height

Amongst all the treatments, the maximum height was observed for T₉ (168.13 cm) with 25 kg ha⁻¹ zinc application and 60 X 20 cm spacing. Two treatments, T₈ (167.57 cm) and T₆, (167.13 cm) showed comparable results to T₉. Wider spacing resulted in better plant height as there was less competition for nutrients, water and sunlight whereas closed spacing leads to decrease in stem circumference. Overcrowding

brings out less light to the plant population and the lower internodes have restricted extension. The present results are in line with that of Neupane et al. [10]. Along with spacing, application of zinc was directly proportional to plant's height. Increase in zinc promotes photosynthesis and formation of chlorophyll and carbohydrates; regulation of stomata and enzyme activities. These results are akin to Arab et al. [11]. This was also confirmed by Rakesh Kumar and Bohra [12]. The least plant height was reported in the treatment (T1).

3.1.2 Number of leaves per plant

Amongst all the treatments, the maximum number of leaves per plant was observed from T₉ (13.25) with 25 kg ha⁻¹ zinc application and 60 X 20 cm spacing. Two treatments, T₈ (13.15) and T₆, (13.07) showed comparable results to T₉. The amount of zinc was directly proportional to the number of leaves per plant. With increment in applied zinc, there is promotion in growth and development hormone auxin, cell division and plant metabolism. The current results are in accordance to that of Tariq et al. [13]. The results were also in confirmity with Azab [14]. The lowest number of leaves was obtained in the treatment (T1).

3.1.3 Dry plant mass (g/plant)

Amongst all the treatments, the highest dry plant mass was observed for T₉ (90.96 g/plant) with 25 kg ha⁻¹ zinc application and with 60 X 20 cm spacing. Two treatments, T₈ (90.81 g/plant) and T₆, (90.46 g/plant) showed comparable results to T₉. Smaller spacing results in reduced photosynthesis due to poor sunlight and CO₂ level whereas wider spacing improves the dry matter accumulation from 20 to 80 DAS. These results are related to that of Sumeria et al. [15]. The quantity of zinc applied also had positive effect on the biomass of the plant. The higher quantity of zinc applied inflated the dry plant mass as well. In addition to this, other growth attributes such as plant height, stem girth and weight of roots affected the dry plant mass [16]. This is also confirmed by Amutham et al. [17]. The lowest dry weight was obtained in the treatment (T1).

3.2 Yield Attributes

3.2.1 Number of cobs per plant

Amongst all the treatments, the maximum number of cobs per plant was observed for T₉ (2.37) with 25 kg ha⁻¹ zinc application and with 60

X 20 cm spacing. Two treatments, T₈ (2.34) and T₆, (2.17) showed comparable results to T₉. These findings are in accordance to Anjum et al. [18] where it was stated that zinc plays a vital role in regulation of growth hormones such as auxin, promotes synthesis of carbohydrates, protein as well as pollen which in turn results in higher number of cobs. The results were also in confirmity with Meena et al. (2013).

3.2.2 Length of cob per plant (cm)

Amongst all the treatments, the maximum cob's length was observed for T₉ with 25 kg ha⁻¹ zinc application and with 60 X 20 cm spacing. Two treatments, T₈ and T₆, showed comparable results to T₉. This was also confirmed by Arab et al. [11].

3.2.3 Cob weight (g)

3.2.3.1 With husk

Amongst all the treatments, the highest weigh of cob (with husk) was observed for T₉ (47.92 g) with 25 kg ha⁻¹ zinc application and with 60 X 20 cm spacing. Two treatments, T₈ (47.43 g) and T₆, (46.87 g) showed comparable results to T₉.

3.2.3.2 Without husk

Amongst all the treatments, the highest weigh of cob (without husk) was observed for T₉ (22.70 g) with 25 kg ha⁻¹ zinc application and with 60 X 20 cm spacing. Two treatments, T₈ (22.20 g) and T₆, (21.72 g) showed comparable results to T₉.

The yield and its attributes are the sum of both the photosynthates as well as their translocation from source to their sink and this is affected by the presence of minerals. Minerals aid in production of enzymes and co-enzymes which in turn have an influence on photosynthetic pathways. One such mineral, Zinc, promotes synthesis of chlorophyll, metabolites, growth-related hormones such as auxin. Zinc is having a productive role over photosynthesis and other metabolic activities elevating growth and development of plant and thus, more yield. The results of our present findings are in accordance to those of Arab et al. [11] and Naik et al. [19].

3.2.4 Cob yield (t/ha)

3.2.4.1 With husk

Amongst all the treatments, the maximum cob yield (with husk) was observed for T₃ (14.63 t/ha) with 25 kg ha⁻¹ zinc application and with 60 X 20

cm spacing. Two treatments, T₈ (14.24 t/ha) and T₆, (13.67 t/ha) showed comparable results to T₃. This is also confirmed by Mona [20]. Treatment (T₇) obtained lowest cob yield with husk

3.2.4.2 Without husk

Amongst all the treatments, the maximum cob yield (without husk) was observed for T₃ (5.09 t/ha) with 25 kg ha⁻¹ zinc application and with 40 X 20 cm spacing. Another treatment, T₂ (4.77 t/ha) showed comparable results to T₃.

The amount of spacing contributes towards competition of nutrients, light and moisture

amongst the plant crops. Lowering the plant density has a positive effect on the yield of seeds. With more competitiveness, reduced sunlight hinders the growth at vegetative phase and ultimately the reproductive stage is not achieved which results in low yield. As concluded by Ariraman et al. [21] the reduced yield could be because lesser plants achieved reproductive phase.

Along with spacing, zinc application influenced the yield of seed as well. Zinc directly effects the synthesis of tryptophan and auxin which sequentially impacts the seed yield and its attributes such as cobs per plant, length and

Table 1. Effect of levels of zinc and spacing on growth attributes of baby corn

Treatments	Plant height (cm)	No. of leaves/plant	Dry weight (g/plant)
1. 15 kg ha ⁻¹ ZnSO ₄ + 40cm x 20 cm	161.40	12.13	86.02
2. 20 kg ha ⁻¹ ZnSO ₄ + 40cm x 20 cm	162.77	12.25	87.18
3. 25 kg ha ⁻¹ ZnSO ₄ + 40cm x 20 cm	164.27	12.70	88.20
4. 15 kg ha ⁻¹ ZnSO ₄ + 50cm x 20 cm	163.70	12.52	87.84
5. 20 kg ha ⁻¹ ZnSO ₄ + 50cm x 20 cm	166.30	12.93	89.83
6. 25 kg ha ⁻¹ ZnSO ₄ + 50cm x 20 cm	167.13	13.07	90.46
7. 15 kg ha ⁻¹ ZnSO ₄ + 60cm x 20 cm	165.80	12.75	88.67
8. 20 kg ha ⁻¹ ZnSO ₄ + 60cm x 20 cm	167.57	13.15	90.81
9. 25 kg ha ⁻¹ ZnSO ₄ + 60cm x 20 cm	168.13	13.25	90.96
F- test	S	S	S
S. Em (±)	0.34	0.09	0.09
C. D. (P = 0.05)	1.03	0.27	0.27

Table 2. Effect of levels of zinc and spacing on yield attributes and yield of baby corn

Treatments	No. of cobs/plant	Length of the cob (cm)		Cob weight (g)		Cob Yield(t/ha)		Green fodder Yield (t/ha)
		With husk	Without husk	With husk	Without husk	With husk	Without husk	
1. 15 kg ha ⁻¹ ZnSO ₄ +40cmx20 cm	1.24	13.20	7.30	41.02	17.64	12.39	3.77	26.15
2. 20 kg ha ⁻¹ ZnSO ₄ +40cmx20 cm	1.43	14.47	7.60	42.05	18.14	14.24	4.77	27.87
3. 25 kg ha ⁻¹ ZnSO ₄ +40cmx20 cm	1.72	15.93	7.83	44.15	19.35	14.63	5.09	28.83
4. 15 kg ha ⁻¹ ZnSO ₄ +50cmx20 cm	1.61	15.20	7.67	43.47	18.38	10.98	3.25	24.97
5. 20 kg ha ⁻¹ ZnSO ₄ +50cmx20 cm	1.95	17.20	8.20	45.57	20.85	12.89	4.09	26.94
6. 25 kg ha ⁻¹ ZnSO ₄ +50cmx20 cm	2.17	17.77	8.27	46.87	21.72	13.67	4.53	27.50
7. 15 kg ha ⁻¹ ZnSO ₄ +60cmx20 cm	1.83	16.63	7.90	44.69	20.46	9.60	2.83	23.78
8. 20 kg ha ⁻¹ ZnSO ₄ +60cmx20 cm	2.34	18.07	8.50	47.43	22.20	10.01	3.02	24.12
9. 25 kg ha ⁻¹ ZnSO ₄ +60cmx20cm	2.37	18.77	8.58	47.92	22.70	11.82	3.43	25.71
F test	S	S	S	S	S	S	S	S
S. Em (±)	0.07	0.31	0.07	0.39	0.34	0.32	0.20	0.31
CD (P = 0.05)	0.20	0.94	0.21	1.17	1.01	0.97	0.61	0.94

weight of seeds. Additionally, zinc contributes towards nutrient metabolism, biological activity and growth and development since enzyme activity is improved as well. This results in more cob yield. This was also similar to findings noticed by Naik et al. [19]. The least cob yield without husk was obtained in the treatment (T7).

3.2.5 Green fodder yield (t/ha)

Amongst all the treatments, the highest green fodder yield was observed for T₃ (28.33 t/ha) with 25 kg ha⁻¹ zinc application and with 40 X 20 cm spacing. Two treatments, T₂ (27.87 t/ha) and T₆, (27.50 t/ha) showed comparable results to T₃.

Zinc has an important role in plant growth and metabolism. It has a positive effect on physiological processes such as synthesis of chlorophyll, carbohydrates and protein as well as forms a vital part in gas exchange through stomata, plant biomass accumulation and starch utilization. The conversion of ammonia to nitrate is regulated via zinc as well. Parellel findings have been noted by Tamil Amutham et al. [17]. This was also confirmed by Mahid et al. [22]. Treatment (T7) obtained lowest green fodder yield.

4. CONCLUSION

Amongst all the treatments, T₉ having 60 X 20 cm spacing and 25 kg ha⁻¹ ZnSO₄ had the maximum plant productivity and its attributes such as plant height, number of leaves per plant, dry plant mass and cob's properties. Further, the maximum crop growth rate, yield and yield attributes such as cob yield and green fodder yield. It can be concluded that wider spacing and increased zinc application has positive effect on plant productivity and growth attributes as well.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Singh G, Kumar S, Singh R, Singh SS. Growth and yield of Baby Corn as influenced by varieties, spacings and dates of sowing. Indian Journal of Agriculture Research 2015;49(4):353-357.
- Dass S, Yadav VK, Kwatra A, Jat ML, Rakshit S, Kaul J, et al. Baby corn in India. Technical Bulletin, Directorate of Maize Research, Pusa Campus, New Delhi. 2008;6:1-45.
- Reena Rani, Sheoran RK, Pooja Gupta Soni, Sakshi Kaith, Arpita Sharma. Baby corn: A wonderful vegetable. International Journal of Science, Environment and Technology. 2017;6(2):1407-1412.
- Pandey AK, Mani VP, Ved, Prakash, Singh RD, Gupta HS. Effect of varieties and plant densities on yield, yield attributes and economics of baby corn (*Zea mays*). Indian Journal of Agronomy. 2002;47(2): 221-226.
- Marschner M. Mineral Nutrition of Higher Plants. 2nd Edition Academic Press, London, New York. 1995;200-255. ISBN-10:0124735436.
- Tisdale SL, Nelson WL, Beaten JD. Zinc In soil Fertility and Fertilizers. Fourth edition, New York: Macmillan Publishing Company; 1984.
- Monneveux. P, Zaidi PH, Sanchez C. Population density and low nitrogen affects yield. Associated Traits in Tropical Maize. Crop Science. 2005;45(2):103-106.
- Bruns HA, Abbas HK. Ultra-high plant populations and nitrogen fertility effects on corn in the Mississippi Valley. Agronomy Journal. 2005;97(4):1136.
- Gozobenli H, Kilinc M, Sener O, Konuskan O. Effects of single and twin row planting on yield and yield components in maize. Asian Journal of Plant Science. 2004;3:203-206.
- Neupane MP, Singh RK, Rakesh K, Anupma K. Response of baby corn (*Zea mays* L.) to nitrogen sources and row spacing. Environment and Ecology. 2011;29(3):1176-1179.
- Arab GM, Dina A Ghazi, El-Ghamry AM. Effect of zinc application on maize grown on alluvial soils. Journal of Soil Science and Agricultural Engineering, Mansoura University. 2018;9(9):419-426.
- Rakesh kumar, Bohra JS. Effect of NPKS and Zn application on growth, yield, economics and quality of baby corn. Archives Agron. Soil Sci. 2014;60(9):1193-1206.
- Tariq A, Shakeel A, Anjum, Mahmood A, Randhawa Ullah E, Naeem M, Qamar R, Ashraf U, Nadeem M. Influence of zinc nutrition on growth and yield behaviour of maize (*Zea mays* L.) hybrids. American Journal of Plant Sciences. 2014;5:2646-2654.

14. Azab MEE. Increasing Zn ratio in a compound foliar NPK fertilizer in relation to growth, yield and quality of corn plant. *Journal of Innovation of Pharma Biological Science*. 2015;2:451-468.
15. Sumeriya HK, Singh P, Nepalia V, Sharma V, Upadhyay B. Response of elite sorghum genotypes to planting geometry and fertility levels. *Research on Crops*. 2007;8(2):312-315.
16. Palai JB, Sarkar NC, Jena J. Effect of zinc on growth, yields, zinc use efficiency and economics in baby corn. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(2):1641-1645.
17. Tamil Amutham G, Karthikeyan R, Thavaprakash N, Bharathi C. Agronomic bio-fortification with zinc on growth and yield of baby corn under irrigated condition. *Journal of Pharmacognosy and Phytochemistry*. 2019;8(3):434-437.
18. Anjum SA, Saleem MF, Shahid M, Shakoor A, Safeer M, Khan I, Farooq A, Ali I, Nazir U. Dynamics of soil and foliar applied boron and zinc to improve maize productivity and profitability. *Pakistan Journal of Agricultural Research*. 2017;30(3):294-302.
19. Naik C, Meena MK, Ramesha YM, Amaregouda A, Ravi M, Dhanoji MM. Morpho-physiological impact of growth indices to Biofortification on growth and yield of sweet corn (*Zea mays* L. Saccharata). *Bulletin of Environment, Pharmacology and Life Sciences*. 2020;9(3):37-43.
20. Mona EA. Increasing Zn ratio in a compound foliar NPK fertilizer in relation to growth, yield and quality of corn plant. *Journal of innovations in Pharmaceuticals and biological Sciences*. 2015;2(4):451-4.
21. Ariraman R, Selvakumar S, Mansingh MDI, Karthikeyan M, Vasline YA. Effect of zinc application on growth, yield parameters, nutrient uptake, yield and economics of maize. *Agricultural Reviews*. 2021;10:1-6.
22. Mahdhi SS, Hysain B, Singh L. Influence of seed rate, nitrogen and Zinc on fodder maize (*Zea mays*) in temperate condition of western Himalayas. *Indian journal of Agronomy*. 2012;57(1):85-88.

© 2022 Rohini et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/86231>