



## **Ameliorative effect of Ethanol Extract of *Annona Muricata* Leaves in Sodium Arsenite Induced-Toxicity in Male Wistar Rats**

**Adegboyega A. M.<sup>1,2\*</sup>, Odunola O. A.<sup>1</sup>, Akinwumi K. A.<sup>1</sup>, Babalola O.<sup>2</sup>  
and Akinwande A.<sup>2</sup>**

<sup>1</sup>*Cancer Research and Molecular Biology Laboratories Department of Biochemistry,  
University of Ibadan, Ibadan, Oyo State, Nigeria.*

<sup>2</sup>*The Polytechnic, Ibadan, Chemistry Department, Ibadan, Oyo State, Nigeria.*

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Author AAM designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors AKA and OOA managed the analyses of the study. Authors BO and AA managed the literature searches. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJBCRR/2021/v30i230249

#### Editor(s):

(1) Dr. Varun Gupta, Albert Einstein College of Medicine, USA.

#### Reviewers:

(1) Noha M. Shawky, University of Maryland Medical Center (UMMC), USA.

(2) Ana Rosa Rincón Sánchez, University of Guadalajara, Mexico.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/66966>

**Original Research Article**

**Received 14 March 2021**

**Accepted 19 May 2021**

**Published 25 May 2021**

### **ABSTRACT**

Ingestion of arsenic, a known contaminant in drinking water causes cancer at multiple tissues and there is no cure. Consumption of arsenic contaminated water has been implicated metalloinduced carcinogenesis. Research is therefore directed at chemoprevention using medicinal herbs for the management of arsenicosis. In this study hepatoprotective activity of ethanolic extract of *Annona muricata* (AM) leaves was assessed against sodium arsenite (SA) induced hepatic injury in albino rats. The animals were pre-treated with either 250 or 500mg/kg body weight of rat before exposure to SA. SA was dissolved in distilled water and administered at a dose of 5 mg/kg body weight on the 7<sup>th</sup>, 14<sup>th</sup> and 21<sup>st</sup> day of the experiment. SA was observed to induce a significant increase ( $p < 0.05$ ) in serum aspartate transaminase (AST), alanine transaminase (ALT) and alkaline phosphatase activities (ALP). However, pretreatments of rats with various doses of AM significantly ( $P < 0.005$ ) reduced serum enzyme levels to near normal against SA treated rats. Furthermore,

\*Corresponding author: E-mail: [adeniketijaniade@gmail.com](mailto:adeniketijaniade@gmail.com);

histopathological observations revealed that treatment with AM extract protected the animals from SA induced liver damage. The results indicated that the leaves of *Annona muricata* possess hepatoprotective activity on SA induced hepatic injury in rats.

**Keywords:** Hepatoprotective; *Annona muricata*; sodium arsenite; transaminase; Oxidative stress.

## 1. INTRODUCTION

Continuous exposure of humans to arsenic through long term ingestion of contaminated water and its attendant health problem has been reported [1]. Epidemiological studies conducted in Taiwa [2], Chile [3] and Japan [4] indicated a connection between arsenic exposures from contaminated drinking water among the inhabitants. Arsenic is a well-known human carcinogen, which potentially affects ~160 million people worldwide via exposure to unsafe levels in drinking water [5]. It is an element present in food, soil, water and air, and it is released into the environment from both natural and man-made sources [6]. Humans may be exposed to arsenic via ingestion through drinking water (major), inhalation and skin absorption [7]. Arsenic in drinking water is typically inorganic, and can be present either as  $As^{+3}$  (arsenite) or  $As^{+5}$  (arsenate). However, the ingestion of inorganic arsenic is a significant public health hazard in the world. Arsenic toxicity has been reported to be associated with a variety of cancers, dermatitis, cardiovascular diseases, peripheral neuropathy, diabetes mellitus, renal failure and liver dysfunction. Also, arsenic toxicity also induces generation of reactive oxygen species (ROS), which may lead to membrane damage, oxidative stress and carcinogenesis of various organs and subsequent cellular damage and organ disorders [8]. The liver is the primary target organ for the metabolism of arsenicals. The major metabolic pathway of inorganic arsenic in humans is its methylation in the liver. The methylation of arsenic has been demonstrated by the presence of monomethylarsonic acid (MMA) and dimethylarsinic acid (DMA) in the urine and bile [9,10]. Liver function test is among the most commonly used and primary clinical investigation for the assessment of liver function. Arsenic intoxication in experimental animals has been linked with micronucleus formation and hepatic tumors [11,12]. In spite of the tremendous advances made in allopathic medicine, no effective hepatoprotective medicine is available.

*Annona muricata* L. (soursop), a member of the Annonaceae family, is a widely distributed plant in Central and South America and tropical

countries like Nigeria where it is used in the treatment of asthma, cough, fever, headache, hypertension, and toothache. Previous studies have shown that *A. muricata* is active against several cancer cell lines (Moghadamtousi et al, 2014). The anticancer activity of *A. muricata* has been attributed to antioxidant and apoptosis inducing potential in cells (Moghadamtousi et al, 2014). However, there are no scientific evidences regarding the hepatoprotective activity of this plant against sodium arsenite.

Therefore, the aim of this study was to investigate the mitigating effect of *Annona muricata* leaves against arsenic-induced liver oxidative damage in rats.

## 2. MATERIALS AND METHODS

### 2.1 Reagents and Kits

Sodium arsenite ( $NaAsO_2$ ; BDH chemicals Ltd poole England) was dissolved in distilled water and administered at a dose of 2.5 mg/kg body weight corresponding to 1/10th of the oral LD50 of the salt [13]. Aspartate aminotransferase (AST), alanine aminotransferase (ALT), and alkaline phosphate kits were obtained from Randox Laboratories, UK. All other chemicals and reagents were of analytical grade and were products of Sigma Chemical Co. St. Louis, MO., USA or BDH Chemical Ltd, Poole, England.

### 2.2 Plant Extraction

Fresh leaves of the plant were harvested during the month of August, 2018, identified and Voucher specimen deposited at the herbarium of the Department of Botany, University of Ibadan. Extraction of air-dried leaves of *Annona muricata* was carried out in ethanol for 120 hours at room temperature. The extract was filtered, concentrated, freeze-dried and stored at 4°C.

### 2.3 Phytochemical Screening

A preliminary phytochemical screening of *Annona muricata* leaves was carried out. The phytochemical profile was performed as described by Kokate, [14] and Harborne, [15]. The presence of alkaloids, flavonoids, glycosides, tannins and saponins were detected.

## 2.4 Experimental Animals

Thirty male Wistar albino rats (180–200 g) were obtained from the animal house of the Physiology Department, University of Ibadan, Ibadan. The animals were grouped and housed in cages with not more than six animals per cage and maintained under standard laboratory conditions (temperature  $25 \pm 2$  °C) with light and dark cycles of 12 and 12 h, respectively. They were allowed free access to the standard dry pellet diet and water *ad libitum*.

## 2.5 Hepatoprotective Study

The rats were divided into six groups of five animals each. Group I, Negative control, was given 1 mL normal saline orally once daily, Group II was treated with 5.0 mg/kg BW of NaAsO<sub>2</sub>, Group III was administered with 250 mg/kg BW AM only, Group IV was given 500 mg/kg BW AM only, Group V was given 250 mg/kg of AM and NaAsO<sub>2</sub>, Group VI was administered with 500 mg/kg AM and NaAsO<sub>2</sub>. The NaAsO<sub>2</sub> was given orally once on days 7, 14 and 21, while AM was administered orally daily for 21 days. The dose of SA used corresponds to 1/5<sup>th</sup> of the oral LD<sub>50</sub> of the salt [13]. The activity of serum aspartate transaminase (AST) and alanine transaminase (ALT) was estimated according to Reitman and Frankel [16] using commercial diagnostic kits. This method involves the reaction of pyruvate, the product of transamination reaction catalyzed by ALT or AST, with 2, 4 -dinitrophenyl hydrazine to produce intensely colored hydrazone read at 546 nm using a spectrophotometer (Spectronic-20). and serum alkaline phosphatase activity (ALP) was estimated according to Kind and King [17].

## 2.6 Histopathological Analysis of Liver

For histological studies, liver tissues were fixed with 10 % phosphate-buffered neutral formalin, dehydrated in graded (50–100 %) alcohol and embedded in paraffin wax and sections were made of 4-6µm. After staining with hematoxylin and eosin, slides were examined under the microscope (Olympus, Japan) for histopathological changes and photographed.

## 2.7 Statistical Analysis

Experimental data were analyzed using one way analysis of variance (ANOVA). Duncan's multiple range test was used to determine significant differences between means. The statistical analyses were performed using computer program Statistical Packages for Social Science

(SPSS) (SPSS, 17.0). Differences between means were considered significant at  $P < 0.05$ .

## 3. RESULTS

Elevated levels of serum enzymes are indicative of cellular leakage and loss of functional integrity of cell membrane in liver. The intoxication of SA increased hepatic transaminases level in serum ( $P < 0.05$ ), and these changes was ameliorated by *Annona muricata* ( $P < 0.05$ ) coadministration which could be attributed to the phytochemicals present in *Annona muricata*. Furthermore, histological observations confirmed hepatoprotective role of *Annona muricata* leaves against sodium arsenite induced hepatotoxicity.

## 4. DISCUSSION

Cancer disease is increasing annually worldwide, creating some concerns regarding the efficacy of the present treatment study. Arsenic toxicity has been linked to diverse defects in both experimental animals and in humans [1] and exposure to arsenic via the intake of contaminated water has been linked with diverse health defects like certain forms of cancer, skin lesions, and non-cancer health effects such as neurological disorders and impaired cognitive development in children [18].

Arsenic intoxication in experimental animals has been linked with micronucleus formation and hepatic tumors [12] and the liver is an important target organ for arsenic toxicity [19]. Consumption of food products rich in antioxidants have been suggested and shown to hold promise in mitigating disease onset and chemical-induced carcinogenesis. *Annona muricata* is commonly known as soursop or Gaviola which produces an edible fruit. This study reports the effect of the ethanolic extract of *Annona muricata* leaves on SA-induced toxicity on albino Wistar rats. The result obtained from the phytochemical analysis of ethanol extract of *Annona muricata* leaves indicated the presence significant amount of flavonoids, tannins, saponins, alkaloids and glycosides as presented in Table 1. These chemicals have been documented with antitumor, antiparasitic and antimicrobial activities. *Annona muricata* leaves is very rich in flavonoids which have antioxidants effects associated with various diseases such as cancer, Alzheimer's disease, atherosclerosis [20]. Alkaloid is known to possess antioxidant properties that can mitigate cellular macromolecule damage from oxidative radical species [21]. Such as DNA strand damage,

resulting in genomic instability implicated in carcinogenesis. Saponins have been reported to have hemolytic properties, although they are not toxic when administered orally. Polyphenolic tannins have been reported to exhibit metal ions chelating properties that can help in mopping up trivalent arsenite upon exposure [22]. However, tannins can be poisonous by depriving the body of essential metal ions needed for proper functioning.

There was no significant increase in body weight throughout the course of the study (Table 2). Compared with the negative control, there was significant decrease in liver weight of SA-only-treated group. This resulted from arsenic toxicity in the liver and increased hepatic metabolism to eliminate it. These observations are consistent with reports that SA toxicity can compromise the integrity of the liver in mouse, rat, and goat [23], [24]. Liver weights and relative liver weights of groups treated with SA and AM and AM only were similar to those of the negative control, indicative of AM exhibiting a potent protective effect on hepatocytes on one hand and in response to SA-induced toxicity thereby ameliorating SA-induced toxicity on hepatocytes.

Exposure to SA had been shown to increase activity of liver transaminases in the blood [25], this is an index of hepatotoxicity which might have resulted from oxidative stress-related damages to hepatocytes membrane and leakage of hepatic transaminases into extracellular spaces ultimately finding their way into the blood from the liver [26]. Aspartate aminotransferase (AST) and Alanine aminotransferase (ALT) are members of transaminase family of enzymes. They are also known as aminotransferases, they catalyze the transfer of amino groups between L-alanine and glutamate for physiological purpose. ALT and AST are found in large amount in the liver and also small amount are found in the heart, kidney and muscles. When the liver is injured or inflamed as the case may be via its exposure to various forms of toxic substances, the level of ALT and AST in the blood is usually elevated. The level of these enzymes in the blood is directly related to the extent of the tissue

damage [27]. The induction of ALT and AST activity following sodium arsenite has been well documented [28].

From the results presented in (Table 3) there is a significant increase in the activities of AST, ALP, and ALT ( $P < 0.05$ ) in the serum of SA-treated rats when compared with the negative control. The observations made here are consistent with the findings from previous findings [28,29] However, there is a significant decrease ( $P < 0.05$ ) in serum activities of AST and ALP in the SA and AM-treated group compared with negative control. This is suggestive of hepatocytes protection from AM co-treatment against SA-induced damages. This can be attributed to the antioxidant present in AM. This is in accordance with some recent studies that recommend the use of antioxidants and antioxidant-rich foods and herbal medicinal plant for the management of arsenicosis [30]. The activities of the liver enzymes (AST, ALT, and ALP) were significantly low in the AM-only-treated group confirming protective property of AM. The reversal of increased serum enzymes in SA-induced liver damage by the extract may be due to the prevention of the leakage of intracellular enzymes by its membrane stabilizing activity. This is in agreement with the commonly accepted view that serum levels of transaminases return to normal with the healing of hepatic parenchyma and the regeneration of hepatocytes [31].

Histological examinations of liver sections of treated animals showed that SA was potentially hepatotoxic as reflected by severe vacuolation with moderate congestion of the sinusoid. Liver sections from SA and AM-treated group exhibited very mild vacuolation, while those of the AM-treated group showed no visible lesions confirming a modulatory effect of AM on SA-induced hepatocytes damage.

Our finding from liver histopathology suggests SA-induced toxicity and ameliorative potentials of AM on SA induced hepatocytes damage.

**Table 1. Phytochemical screening of *Annona muricata* leaves**

Constituents	Reaction to Tests/Reagents	Inference
Alkaloids	Positive to Mayer, Dragendoff Wagner and Picric reagent	X
Flavonoids	Acid-alcohol test	XXXX
Glycosides	Reddish-brown colour at the interference	XX
Saponins	Positive to Froth	XXX
Tannins	Turned blue-black precipitate on addition of FeCl <sub>3</sub>	X

Key: X = present, XX = Moderately present, XXXX = abundance

**Table 2. Influence of AM treatment on body weight and relative change in organ weight of rats treated with sodium arsenite.**


Treatments	Initial body weight (g)	Final body weight (g)	% Weight change	Liver weight (g)	% Liver weight
Negative control	180.00 ± 10.96	204.00 ± 15.00	13.3 3± 2.95	6.89 ± 1.00	3.38 ± 0.25
SA	164.00 ± 16.50	188.00 ± 6.96	14.63 ± 3.01	5.52* ± 0.66	2.94* ± 0.27
SA + 250 mg/kg extract	160.00 ± 18.90	186.00 ± 16.45	16.25 ± 3.22	6.70 ± 0.74	3.60 ± 0.13
SA + 500 mg/kg extract	156.00 ± 9.40	174.00 ± 15.46	11.54 ± 2.86	5.54 ± 0.68	3.18 ± 0.47
250 mg/kg extract	171.00 ± 5.83	195.00 ± 11.20	14.04 ± 3.40	5.80 ± 0.76	2.97 ± 0.01
500 mg/kg extract	186.00 ± 10.86	202.00 ± 12.80	8.60 ± 2.01	6.30 ± 0.54	3.12 ± 0.13

*Values are represented as mean ± SD of six animals in each group; \* = The mean difference is significant (p<0.05) when compared with the negative control*

**Table 3. Effects of AM administration on serum level of hepatic alanine transaminase (ALT), aspartate transaminase (AST) and alkaline phosphatase (ALP) in rats treated with sodium arsenite**

Treatments	ALT/IU	AST/IU	ALP/IU
Negative control	11.00 ± 2.0	13.00 ± 1.8	29.5 ± 2.5
SA	39.5 ± 2.6*	38.5 ± 2.0*	47.0 ± 2.3*
SA + 250 mg/kg extract	30.00 ± 1.8	29.00 ± 2.5	41.00 ± 1.8
SA + 500 mg/kg extract	26.8 ± 1.6	28.7 ± 3.1	39.00 ± 1.5
250 mg/kg extract	26.34 ± 1.2	25.00 ± 1.3	33.80 ± 1.8
500 mg/kg extract	24.50 ± 2.2	23.80 ± 1.4	31.00 ± 1.2

*Values represent mean ± SD of six animals in each group; \* = Significantly different from the negative control*



**Fig. 1.** The photomicrograph of liver sections of rats treated with *Annona muricata* and / or sodium arsenite (A) Negative control with no visible lesion (B) Animals administered sodium arsenite only showing severe vacuolation (arrow) with moderate congestion of the sinusoid. (C) Animals fed 250 and 500 mg/kg BW *Annona muricata* with no visible lesion. (D) Animals exposed to sodium arsenite and 250 and 500mg/kg BW *Annona muricata* with mild vacuolation (arrow). (Mx 400)

All these results indicated a hepatoprotective potential of the extract of *Annona muricata* and these biological activities may be due to the phytochemicals present in the extract. Further studies to isolate, purify and characterize the active compounds in the extract are significant and also a detailed study at molecular level is needed to know the exact mechanism of *Annona muricata* protective mechanism.

## 5. CONCLUSION

Based on the results obtained, pretreatment with the extract was found to protect the liver from sodium arsenite induced damage which may be attributed to the rich antioxidant nature of the leaf extract. It may be concluded that the extract of *annona muricata* has a significant effect on liver injuries resulting in improved serum biochemical parameters such as ALP, AST and ALT and hence may be use in mitigating arsenic induced toxicity.

## DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for

any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## ETHICAL APPROVAL

All procedures of this study described were reviewed and approved by the Institutional Animal Ethical Committee.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Waalkes MP, Ward JM, Diwan BA. Induction of tumors of the liver, lung, ovary and adrenal in adult mice after brief maternal gestational exposure to inorganic arsenic: Promotional effects of postnatal phorbol ester exposure on hepatic and pulmonary, but not dermal cancers. *Carcinogenesis*. 2004;25:133–41
2. Chiou HY, Hsueh YM, Liaw KF, Horng SF, Chiang MH. Incidence of internal cancers and ingested inorganic arsenic: a seven-year follow-up study in Taiwan. *Cancer Res*. 1995;55:1296-1300.

3. Smith AH, Goycolea M, Haque R, Biggs ML. Marked increase in bladder and lung cancer mortality in a region of Northern Chile due to arsenic in drinking water. *Am J Epidemiol.* 1998;147:660-669.
4. Tsuda T, Babazono A, Yamamoto E, Kurumatani N, Mino Y, et al. Ingested arsenic and internal cancer: a historical cohort study followed for 33 years. *Am J Epidemiol.* 1995;141:198-209.
5. IARC (International Agency for Research on Cancer): Monographs on evaluation of carcinogenic risk to humans. In: Some Drinking Water Disinfectants and Contaminants, including Arsenic. 2004;84:269-477.
6. Chakraborti D, Sengupta MK, Rahman MM, Ahmed S, Chowdhury UK, Hossain MA, Mukherjee SC, Pati S, Saha KC, Dutta RN, Quamruzzaman Q: Groundwater arsenic contamination and its health effects in the Ganga-Meghna-Brahmaputra plain. *J Environ Monit.* 2004;6:74-83.
7. Gupta R, Kannan GM, Sharma M, Flora SJS. Therapeutic effects of *Moringa oleifera* on arsenic induced toxicity in rats. *Environ. Toxicol. Pharmacol.*, 2005;20:456-464.
8. Tanju S, Madhuri D. Arsenic induced oxidative stress, hemato-biochemical and histological changes in liver and protective effect of Moringa leaf powder and ascorbic acid in broiler chicken. *J. Chem. Pharm. Res.* 2013;5:112-116.
9. Li X, Pi J, Li B, Xu Y, Jin Y, Sun G: Urinary arsenic speciation and its correlation with 8-OHdG in Chinese residents exposed to arsenic through coal burning. *Bull Environ Contam Toxicol.* 2008;81:406-411.
10. Cui X, Kobayashi Y, Hayakawa T, Hirano S: Arsenic speciation in bile and urine following oral and intravenous exposure to inorganic and organic arsenics in rats. *Toxicol Sci.* 2004;82:478-487.
11. Moore LE, Smith AH, Hopenhayn-Rich C, Biggs ML, Kalman DA, Smith MT. Decrease in bladder cell micronucleus prevalence after intervention to lower the concentration of arsenic in drinking water. *Cancer Epidemiol Biomarkers Prev.* 1997;6(12):1051-1056.
12. Mazumder DN. Effect of chronic intake of arsenic-contaminated water on liver. *Toxicol Appl Pharmacol.* 2005;206(2): 169-175.
13. Preston RJ, Dean BJ, Galloway S, Holden H, McFee AF, Shelby M. Mammalian in vivo cytogenetic assays. Analysis of chromosome aberrations in bone marrow cells. *Mutat Res.* 1987;189(2):157-165.
14. Kokate CK. Practical Pharmacognosy Vallabh Prakashan, New Delhi. 1994;107-113
15. Harborne JB. Phytochemical methods Chapman Hall, London. 1998;60-66
16. Reitman S, Frankel S. A calorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *Am. J. clin. Pathol.* 1957;28:56.
17. Kind P, King E. Estimation of plasma phosphatase by determination of hydrolysed phenol with aminoantipyrine. *J. clin. Pathol.*, 1954;7:322.
18. Abernathy CO, Liu YP, Longfellow D, Aposhian HV, Beck B, Fowler B, Goyer R, Menzer R, Rossman T, Thompson C, Waalkes M. Arsenic: health effects, mechanisms of actions, and research issues. *Environ Health Perspect.* 1999;107(7):593-597.
19. Parvez F, Chen Y, Argos M, Hussain AZ, Momotaj H, Dhar R, van Geen A, Graziano JH, Ahsan H. Prevalence of arsenic exposure from drinking water and awareness of its health risks in a Bangladeshi population: results from a large population-based study. *Environ Health Perspect.* 2006;114(3):355-359.
20. Burak M, Imen Y. Flavonoids and their antioxidants properties. *Turkiye Klin Bil Derg.* 1999;19:296-304.
21. Lambert JD, Hong J, Yang GY, Liao J, Yang CS. Inhibition of carcinogenesis by polyphenols: Evidence from laboratory investigations. *Am J Clin Nutr.* 2005;81(Suppl 1):284S-291S.
22. Okonkwo TJ, Okorie O, Okonta JM, Okonkwo CJ. Sub-chronic hepatotoxicity of *Anacardium occidentale* (Anacardiaceae) inner stem bark extract in rats. *Indian J Pharm Sci.* 2010;72:353-7.
23. Sharma A, Sharma MK, Kumar M. Modulatory role of *Emblica officinalis* fruit extract against arsenic induced oxidative stress in Swiss albino mice. *Chem Biol Interact.* 2009;180:20-30.
24. Roy S, Roy M, Pandey PK, Tiwari SP. Effects of tissue trace minerals status and histopathological changes in chronic arsenicosis in goats. *Vet World.* 2009;2(1):8-9.
25. Chattopadhyay S, Ghosh S, Debnath J, Ghosh D. Protection of sodium arsenite -

- induced ovarian toxicity by coadministration of L-ascorbate (Vitamin C) in mature strain rat. Arch Environ Contain Toxicol. 2001;41:83-9
26. Owumi SE, Odunola OA, Gbadegesin MA, Nulah KL. Protective effect of Juglans nigra on sodium arsenite-induced toxicity in rats. Pharmacognosy Research. 2013;5(3):183.
  27. Lum G, Gambino SR. Serum gammaglutamyl transpeptidase activity as an indicator of disease of liver, pancreas, or bone. Clin Chem. 1972;18(4):358-362.
  28. Meliker JR, Wahl RL, Cameron LL, Nriagu JO. Arsenic in drinking water and cerebrovascular disease, diabetes mellitus, and kidney disease in Michigan: a standardized mortality ratio analysis. Environ Health. 2007;2:4-6.
  29. Odunola OA. Comparative effects of some local food condiments on sodium arsenite-induced clastogenicity. Afr J Med Med Sci. 2003;32(1):75-80.
  30. Das NK, Sengupta SR. Arsenicosis: Diagnosis and treatment. India J Dermatol Venereol Leprol. 2008;74:571-81
  31. Thabrew M, Joice P. A comparative study of the efficacy of Pavetta indica and Osbeckia octanda in the treatment of liver dysfunction. Planta Med. 1987;53:239-241.

© 2021 Adegboyega et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle4.com/review-history/66966>