



## Ten Years Post Bilateral TKR with LBA – Physiotherapy Can Help?

**S. S. Subramanian** <sup>a,b#\*</sup>

<sup>a</sup> Sree Balaji College of Physiotherapy, Chennai - 100, India.  
<sup>b</sup> Constituent to (Bharath) University, BIHER, Chennai -73, India.

### **Author's contribution**

*The sole author designed, analysed, interpreted and prepared the manuscript.*

### **Article Information**

#### Editor(s):

(1) Dr. Ikem, Innocent Chiedu, Obafemi Awolowo University, Nigeria.

#### Reviewers:

(1) Narendra Singh Kushwaha, King George Medical University, India.

(2) Nandkishor Goyal, ACPM Medical College, India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:  
<https://www.sdiarticle5.com/review-history/80505>

**Received 05 November 2021**

**Accepted 11 January 2022**

**Published 28 January 2022**

**Original Research Article**

### **ABSTRACT**

Bilateral knee arthroplasty (TKR) remains a common orthopaedic procedure. Geriatric subject after 10 years of bilateral TKR developing chronic Low back ache (CLBA) was treated for two years (Jan 2019 to Dec 2020) with specific exercises, clinical prognosis, and results were discussed with due evidence. Pre-operative existence of lumbar spinal conditions, obesity, and demineralisation were identified to influence on post bilateral TKR to have CLBA. A knowledge of these factors were, to known for clinician to be discussed with the subjects as well due therapeutic measures to be taken by therapist.

**Keywords:** TKA: Total Knee Arthroplasty; LBA: Low Back Ache; OA: Osteoarthritis; QOL: Quality of Life; BMI: Body Mass Index; WC: Waist Circumference.

<sup>#</sup>The Principal,

<sup>\*</sup>Corresponding author: Email: [subramanian.sbcp@bharathuniv.ac.in](mailto:subramanian.sbcp@bharathuniv.ac.in), [subramanian.podhigai1968@gmail.com](mailto:subramanian.podhigai1968@gmail.com);

## 1. INTRODUCTION

An increasing number of Total knee arthroplasty (TKA) globally, there arises a need for improving quality of subjects rehabilitation, follow up, and ensuring patient satisfaction. Outcome of TKA as influenced by literature evidences including preoperative obesity, falls, lumbar pain disorders, chronic OA knee, decrease in proprioception. Whenever possible subjects should pre operatively be treated with known musculoskeletal disorders, along with lowering of obesity to maximise post TKA patients recovery. There were few research follow up post TKA on low back ache, hence this research subject ten years post TKA having chronic Low back ache (LBA) with specific physiotherapy, using lumbar stabilisation exercises, yoga postures, in a year has shown moderate reduction in obesity and an improved functional activities were discussed with evidence.

Boyle et al. [1] have among 40 subjects with pre-operative LBA, have shown no improvement in mental health of SF 36 post TKR. Ayers et al. [2] have among 180 patients with TKA; those with pre-operative musculoskeletal LBA have poor physical function after six months in SF 36, as outcome means on their Quality of life. Stail Bano et al 2014 have insisted on pre-operative lumbar spine disease to be counselled as TKA outcome may be improved. Smith et al. [3] have among 772 TKA with two years follow up have recorded modest deterioration in the contralateral knee with 3.5 units / year. Medication, education, exercises, manual therapy, modalities were not evidenced in chronic LBA [4].

### 1.1 Aims & Objectives of this Research

- a. Where bilateral TKR subjects developing chronic LBA, factors analysed.
- b. To identify the role of specific physiotherapy among these subjects, and evaluate their efficacy.

## 3. CLINICAL PROGNOSIS AND RESULTS

Table of results on cadence, WC, Oswestry scale pre and post values

Values	Oswestry Scale	Cadence	Waist Circumference
PRE	68	25/mt	106 cm
POST	12	48	102 cm
'P'- value	SE – 6.35 P < 0.001	SE – 0.16 P < 0.05	

## 2. MATERIALS AND METHODOLOGY

### 2.1 Background Information

72 year old endomorph female post bilateral TKA complaints of chronic lowbackache Body mass index (BMI) : 29 Kg/ m<sup>2</sup> Waist circumference (WC) : 106 cm Non diabetic/ normotensive, mother of two adults has undergone Bilateral TKR: Left -2008, Right - 2009 was rehabilitated elsewhere for the last Two years & (Since 2018) complaining of chronic lowback ache with occasional radicular symptoms with specific exercises. Her clinical prognosis were discussed with her past medical history reveals hysterectomy in 1990 (She was 42 years) and bilateral OA knees from her age of 48 years. With recurrent lowback pain, which was treated with Non – steroidal Anti – Inflammatory drugs (NSAID) and electrical modalities by physiotherapist elsewhere.

- She was getting treated for her chronic lowback ache with hot pack and specific exercises alone by the author and her clinical course of Two years follow-up (from January 2019 - December 2020). With clinical reasoning on her clinical prognosis and evidenced discussions forms core of this research presentation.
- Aims and objectives of this research were
  1. To analyse chronic lowback ache with bilateral TKR
  2. Evaluate the role of physiotherapy in this clinical scenario

### 2.2 Treatment

She was treated with lumbar stabilisation exercises, Selective yoga postures, Exercises to improve balance, Home programme, Regular walking and avoid long standing with weekly twice frequency of regular physiotherapy during the period from January 2019 till December 2020 her level of prognosis were as below:

- With nature of exercises in supine, side, prone, high sitting and standing postures, exercises were progressed gradually.
- Lumbar stabilisation exercises have resulted in reasonable reduction in pain and an increased quality of life as seen in the above table.
- Yoga postures such as pawanmuktasan, setubandhasana, bshujangasana, paschimottanasana were used mainly to strengthen lower back and the spinal flexibility
- With regular walking herself confidence has increased as her self-care and social activities including driving two wheelers along with an increased duration of walking and improved cadence were recorded.
- She was advised to continue balance exercises and bimonthly follow up with physiotherapy to continued the achieved clinical progress

#### 4. DISCUSSION

Critical research questions arising from this study were answered with evidence as below:

##### 1. What was the significance of lowback ache among bilateral TKR?

Pre - operative evaluation of lumbar spinal disorders remains a major determinant among bilateral TKR, as this can influence outcome of the surgery. Bone demineralisation could again be a factor as this subject had hysterectomy at her age of 42 years, hence proper calcium and Vitamin D supplement should be considered. With being obese, weight reduction means can influence on both low back and the TKR procedure. Longstaff et al. [5] have recorded presence of LBA can lower functional outcome score after TKA. Absence of LBA after TKA was associated with Health related quality of life (HRQOL) Pre - intervention patient factors on outcome of TKA was low back pain [6]. An improved cadence by 48% along with Oswestry scale reflection on her functional activities by 82% improvement were significant.

##### 2. How to treat with physiotherapy in this clinical situation?

Yoga postures helps to improve flexibility among chronic back pain subjects. This research subject was treated with specific yoga postures such as pawanmuktasan, setubandhasana,

bshujangasana, paschimottanasana, veerasana to name a few, as these postures can be added later with her home exercises. Lumbar stabilisation exercises were quite effective as evidenced in reduction of pain, improving strength of spine, lumbar pelvic muscles and evidenced with an improved Quality of life (QOL) as displaced with Oswestry scale in the table of results. Again most of these exercises the subject was advised to continue at home regularly arthroplasty of knee have become a common orthopaedic surgery for the last decade. With an increasing trends in obesity, and prevalence of OA knee, total knee arthroplasty procedures shall be one of the most frequently performed orthopaedic surgeries. At the same time measures to improve quality based rehabilitative measures, due follow up are paramount ensuring greater outcome post TKR. With lesser research on post bilateral TKR subjects QOL, long-term follow up reports are reported. This original research where 72 year old female post bilateral TKR subject developing chronic lowback ache (Nearly after 10 years of knee arthroplasty), was rehabilitated using specific physiotherapy for a year and the results were analysed with evidence. Various causes for bilateral TKR subjects developing chronic LBA including demineralisation after hysterectomy, obesity, pre-operative existing spinal conditions, were discussed.

As these factors can influence on successful outcome of bilateral TKR, should be duly recorded and explained to subjects prior to TKR. Lumbar spinal Exercises (LSE) may be effective in reducing impairments, activity limitations and participation restrictions associated with chronic LBA [7]. LSE where specific exercises aimed at improving motor performances of spiral muscles like transverse abdominis, multifidus and internal oblique [8]. Muscle atrophy, fatigue, motor control deficits of transverse abdominis and multifidus are associated with LBP [9]. Macedo et al. [10] motor control to be an adjunct. Hayden et al. [11] have recorded no role for exercises in acute/ conservative, but chronic; reduce pain and dysfunction [12] found LSE effective among chronic LBA. May and Johnson et al. [13] in a (Systematic Review) SR more in a chronic LBA but not an alternative intervention Rackwitz et al. [14] in a SR, LSE to be better than physician medication but lesser than physio interventions. Haggard and person 2007 evidenced LSE to reduce pain and improve QOL. Standart et al 2008 with moderate evidence but should be individualised. Along with a marginal reduction of

obesity by 3.7% has added with lowering of her chronic LBA.

## 5. CONCLUSION

Ageing a normal process, where geriatric subjects to live with dignity, having maximal independence with self-care activities remains rehabilitative goals following major surgeries. Post bilateral total knee major surgeries. Post bilateral total knee arthroplasty subject developing chronic low back ache was treated for a year with specific exercises, results were analysed and discussed with due evidence in this research.

Post bilateral TKR, the need for continuous follow up including prevention balance exercises, evaluated obesity, keeping an active life style, treating comorbid conditions such as diabetes, special care for pre-operative lowback ache / lumbar disc lesions should be done on a regular basis of a minimal once in 3 months periods, duly recorded, discusses with orthopaedic surgeon as well subjects involved.

These long term regular follow up can maximise post TKR to obtain greater quality of life and maximise their aches and discomforts.

Further larger sample size studies long term retrospective studies of both sex can strengthen findings of this study.

Limitation of this study was the author has treated post TKR (Bilateral) subject only after 10 years and having developed chronic lowback ache. As he was not involved in rehabilitation of the subject post TKR, but TKR being an expensive procedure decreases pain but doesn't resolve many substantial functional limitations associated with OA knee as pointed out by Piva et al. [15], along with later stage intense exercises with individual based exercises to be safe post TKR as recorded by Thoffet et al 2004 and comprehensive exercise programme post TKR leading to functional recovery as supported by Minns et al. [16] were worthy that even after 10years Bilateral post TKR, subject can benefit with patient specific exercises as key findings of this research.

## CONSENT AND ETHICAL APPROVAL

It is not applicable.

## COMPETING INTERESTS

Author has declared that no competing interests exist.

## REFERENCES

1. Boyle KL. Managing a female patient with left low back pain and sacroiliac joint pain with therapeutic exercise: a case report. *Physiother Can.* 2011 Spring;63(2):154-63.
2. Ayers DC, Li W, Oatis C, Rosal MC, Franklin PD. Patient-reported outcomes after total knee replacement vary on the basis of preoperative coexisting disease in the lumbar spine and other nonoperatively treated joints: the need for a musculoskeletal comorbidity index. *J Bone Joint Surg Am.* 2013 Oct 16;95(20):1833-7.
3. Smith JW, Marcus RL, Peters CL, Pelt CE, Tracy BL, LaStayo PC. Muscle force steadiness in older adults before and after total knee arthroplasty. *J Arthroplasty.* 2014 Jun;29(6):1143-8.
4. Philadelphia Panel. Philadelphia Panel evidence-based clinical practice guidelines on selected rehabilitation interventions for low back pain. *Phys Ther.* 2001 Oct;81(10):1641-74.
5. Longstaff LM, Sloan K, Stamp N, Scaddan M, Beaver R. Good alignment after total knee arthroplasty leads to faster rehabilitation and better function. *J Arthroplasty.* 2009 Jun;24(4):570-8.
6. Escobar A, Quintana JM, Bilbao A, Azkárate J, Güenaga JI, Arenaza JC, Gutierrez LF. Effect of patient characteristics on reported outcomes after total knee replacement. *Rheumatology (Oxford).* 2007 Jan;46(1):112-9.
7. O'Sullivan PB. Lumbar segmental 'instability': clinical presentation and specific stabilizing exercise management. *Man Ther.* 2000 Feb;5(1):2-12.
8. Richardson CA, Snijders CJ, Hides JA, Damen L, Pas MS, Storm J. The relation between the transversus abdominis muscles, sacroiliac joint mechanics, and low back pain. *Spine (Phila Pa 1976).* 2002 Feb 15;27(4):399-405.
9. Hides JA, Jull GA, Richardson CA. Long-term effects of specific stabilizing exercises for first-episode low back pain. *Spine (Phila Pa 1976).* 2001 Jun 1;26(11):E243-8.

10. Macedo LG, Latimer J, Maher CG, Hodges PW, McAuley JH, Nicholas MK, Tonkin L, Stanton CJ, Stanton TR, Stafford R. Effect of motor control exercises versus graded activity in patients with chronic nonspecific low back pain: a randomized controlled trial. *Phys Ther.* 2012 Mar;92(3):363-77.
11. Hayden JA, van Tulder MW, Malmivaara A, Koes BW. Exercise therapy for treatment of non-specific low back pain. *Cochrane Database Syst Rev.* 2005 Jul 20;(3):CD000335..
12. Ferreira G, Stieven F, Araujo F, Wiebusch M, Rosa C, Plentz R, Silva M. Neurodynamic treatment did not improve pain and disability at two weeks in patients with chronic nerve-related leg pain: a randomised trial. *J Physiother.* 2016 Oct;62(4):197-202.
13. May S, Johnson R. Stabilisation exercises for low back pain: a systematic review. 2008. In: Database of Abstracts of Reviews of Effects (DARE): Quality-assessed Reviews [Internet]. York (UK): Centre for Reviews and Dissemination (UK); 1995.
14. Rackwitz B, de Bie R, Limm H, von Garnier K, Ewert T, Stucki G. Segmental stabilizing exercises and low back pain. What is the evidence? A systematic review of randomized controlled trials. *Clin Rehabil.* 2006 Jul;20(7):553-67.
15. Piva SR, Susko AM, Khoja SS, Josbeno DA, Fitzgerald GK, Toledo FG. Links between osteoarthritis and diabetes: implications for management from a physical activity perspective. *Clinics in Geriatric Medicine.* 2015;31(1):67–viii.
16. Minns Lowe CJ, Barker KL, Dewey M, Sackley CM. Effectiveness of physiotherapy exercise after knee arthroplasty for osteoarthritis: systematic review and meta-analysis of randomised controlled trials. *BMJ.* 2007;335(7624): 812.

© 2022 Subramanian; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/80505>*