



Impact of Weather on Yield Attributing Characters of Artificially and Naturally Inseminated Colonies of *Apis mellifera* L. and *Apis cerana indica* F.

Pooja Borah^{a*} and M. K. Deka^b

^a Sericultural Training Institute, Titabaor, Assam, 785632, India.

^b Assam Agricultural University, Jorhat, Assam, 785013, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i630693

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/84865>

Received 15 January 2022

Accepted 22 March 2022

Published 26 March 2022

Original Research Article

ABSTRACT

Experiment conducted to study the impact of weather parameters on brood area, pollen area and nectar area of artificially inseminated and naturally inseminated colonies of *Apis mellifera* L. and *Apis cerana indica* F. during April'16 to March'18. Investigation revealed significant negative correlation of brood area ($r = -0.57$ in both AI and NI colonies) of *A. mellifera* L., during 2016-17 and nectar area ($r = -0.71$ in AI and $r = -0.70$ in NI colonies) with maximum temperature and maximum relative humidity, respectively. Whereas, pollen area having significant positive correlation with bright sunshine hours (0.83 in AI and 0.84 in NI colonies) and significant negative correlation with maximum temperature ($r = -0.89$ in AI and $r = -0.88$ in NI colonies) and total rainfall ($r = -0.57$ in AI and $r = -0.60$ in NI colonies). During 2017-18, pollen area showed positive significant correlation with maximum relative humidity ($r = 0.57$ in AI and $r = 0.60$ in NI colonies) and negatively significant correlation with maximum temperature ($r = -0.87$ in AI and $r = -0.90$ in NI colonies) and total rainfall ($r = -0.74$ in AI and $r = -0.76$ in NI colonies). Nectar area having negatively significant correlation with maximum relative humidity ($r = -0.74$ in AI and $r = -0.77$ in NI colonies). However, *A. c. indica* F. revealed significant negative correlation of brood area ($r = -0.58$ in AI and $r = -0.60$ in NI colonies) and nectar area ($r = -0.66$ in AI and $r = -0.70$ in NI colonies) with maximum relative humidity during 2017-18.

Keywords: Weather; *A. mellifera* L.; *A. c. indica* F.; brood area; pollen area; nectar area.

1. INTRODUCTION

Weather influences the growth and development of honeybees. The weather parameters like temperature, relative humidity, rain fall, number of rainy days and sunshine hours might have positive or negative influence on developmental parameters of honeybees. The important developmental parameters of honeybees are brood rearing activity, pollen and nectar store. Weather conditions influences forage availability and thereby directly and indirectly influence honey bee health [1-3]. Weather factors can affect the onset and decline of specific foraging resources, lengthen or shorten the time in which resources are available for bees, change the quality of these resources, and alter the span during which bees can actively forage [4]. Reddy [5] reported that relative humidity and rainfall showed a positive relationship with the pollen gathering activity by the bees. Verma [6] noticed the increase in the pollen foraging activity with rise in the relative humidity, but not so affected by temperature. Mahrotra and Bisht [7] found significant correlation of foraging activity for pollen and nectar foragers with day temperature and humidity. Abrol and Bhat [8] reported that the foraging activity of *A. cerana* was significant and positive with temperature and non significant with relative humidity. The correlation between weather parameters and honeybees are critically important. Therefore the present investigation was aimed to study impact of weather on brood area, pollen area and nectar area of artificially and naturally inseminated colonies of *Apis mellifera* L. and *Apis cerana indica* F.

2. MATERIALS AND METHODS

An investigational data on weather parameters viz., maximum temperature, relative humidity, total rainfall and bright sunshine hours were recorded during the entire period of study from Department of Agro meteorology, AAU, Jorhat to determine the influence of weather parameters on brood area, pollen area and nectar area of artificially and naturally inseminated colonies of *Apis mellifera* L. and *Apis cerana indica* F.

2.1 Statistical Analysis

The data was analysed statistically using Randomized Block Design and correlation of the weather parameters was investigated for their influence on brood area, pollen area and nectar

area with independent variables to find out the influence of weather parameters. The statistical analysis was as follows:

2.2 Simple Correlation Studies

To calculate the correlation co-efficient(r), the following standard statistical formula by Karl Pearson was adopted [9].

$$r = \frac{\sum xy - \frac{\sum x \cdot \sum y}{N}}{\sqrt{\left(\sum x^2 - \frac{(\sum x)^2}{N}\right) \left(\sum y^2 - \frac{(\sum y)^2}{N}\right)}}$$

Where, r = Correlation co-efficient
 n = Number of observation
 x = Independent variables
 y = Dependent variables

Then the Correlation co-efficient(r) was tested for significant or non significant by Fisher 't' test, which is defined as follows :

$$t = \frac{r}{\sqrt{(1-r^2)}} \times \sqrt{n-2} \text{ with } n-2$$

degrees of freedom

For correlation studies, monthly average values of the environmental factors were taken into consideration to know the influence of weather parameters on brood area, pollen area and nectar area during the present investigation.

2.3 Simple Regression Analysis

Simple regression line was fitted to know the impact of independent variables on the dependent variables.

The regression line was given by the equation:

$$Y = a + bx$$

Where, for each unit increase in 'x' variable, the 'Y' variable increases by 'b' unit.

Y= Dependent variables
 x = Independent variables
 a =Intercept

3. RESULT AND DISCUSSION

The influence of various weather factors, viz; maximum temperature, maximum relative humidity, total rainfall and bright sunshine hours on brood area, pollen area and nectar area of *A. mellifera* L. and *A. c. indica* were worked out through simple correlation method. However, Karl Pearson's correlation co-efficient (r) which were found to be significant after being tested by Fisher's t were further studied through regression equation and illustrated graphically as follows:

3.1 Correlation of Weather Parameters with Brood Area, Pollen Area and Nectar Area of *A. mellifera* L.

Correlation studies of weather parameters with brood area, pollen area and nectar area were carried out in *A. mellifera* L. during April'16 to March'17 and the results have been discussed below (Table 1).

Relationship of brood area with maximum temperature: From the investigation, it was revealed that the relationship between average maximum temperature and brood area of *A. mellifera* L. was found to be significant, negative and linear with artificially inseminated colonies ($r = -0.57$, $P < 0.05$). The data revealed 32.9 per cent variation in brood area, which was observed to be controlled by maximum temperature ($R^2 = 0.329$). The regression of brood area on maximum temperature could be expressed by the equation $Y = -144.18X + 7390.1$ (Fig. 1A). Similarly, a significant, negative and linear relationship ($r = -0.57$, $P < 0.05$) was reported between average maximum temperature and brood area of naturally inseminated colonies of *A. mellifera* L. The data indicated 32.4 per cent variation in brood area which was controlled by maximum temperature ($R^2 = 0.324$). The regression of brood area on maximum temperature could be expressed by the equation $Y = -147.94X + 7316.1$ (Fig. 1B). The result was in conformity with the observations of Rahman [10] where he found significant correlation of maximum temperature with brood area of *A. mellifera* L.

Relationship of pollen area with maximum temperature: An investigation revealed that the relationship between average maximum temperature and pollen area was negatively significant and linear in artificially inseminated colonies of *A. mellifera* L. ($r = -0.89$, $P < 0.05$). The data indicated that 80.07 per cent variation in

pollen area was controlled by maximum temperature ($R^2 = 0.8007$). The regression of pollen area on maximum temperature could be expressed by the equation $Y = -108.66X + 3926.8$ (Fig. 1C). The relationship between average maximum temperature and pollen area of naturally inseminated colonies of *A. mellifera* L. was found to be significant, negative and linear ($r = -0.88$, $P < 0.05$). The data indicated 78.4 per cent variation in brood area was observed to be controlled by maximum temperature ($R^2 = 0.784$). The equation $Y = -103.99X + 3680.1$ indicated the magnitude of this association (Fig. 1D). Zapart et al. [11] found negatively significant correlation with maximum, minimum, and average temperatures as well as precipitation in the season.

Relationship of pollen area with total rainfall: A significant, negative and linear relationship was revealed between total rainfall and pollen area of artificially inseminated colonies of *A. mellifera* L. ($r = -0.57$, $P < 0.05$). The data indicated that 32.0 per cent variation in pollen area was controlled by total rainfall ($R^2 = 0.320$). The regression of pollen area on total rainfall could be expressed by the equation $Y = -1.1638X + 940.36$ (Fig. 1E). However, rainfall showed significant, negative and linear effect on pollen area of naturally inseminated colonies of *A. mellifera* L. ($r = -0.60$, $P < 0.05$). However, 37 per cent variation in pollen area was regulated by total rainfall ($R^2 = 0.370$) and the regression of pollen area on total rainfall could be expressed by the equation $Y = -1.2093X + 839.17$ (Fig. 1F). Contrary to the present findings Nag [12] documented positive impact of total rainfall on pollen area.

Relationship of pollen area with bright sunshine hours: From the investigation, it was revealed that bright sunshine hours has positive and linear ($r = 0.83$, $P < 0.05$) effect on pollen area of artificially inseminated colonies of *A. mellifera* L. The data indicated 68.9 per cent variation in pollen area was controlled by bright sunshine hours ($R^2 = 0.689$) and the regression of pollen area on bright sunshine hours could be expressed by the equation $Y = 177.34X + 98.326$ (Fig. 1G). Similarly, the relationship between bright sunshine hours and pollen area of *A. mellifera* L. was found to be significant, positive, and linear with naturally inseminated colonies ($r = 0.84$, $P < 0.05$). From the data it was revealed that 71.2 per cent variation in pollen area was regulated by bright sunshine hours ($R^2 = 0.712$). The regression of pollen area on bright sunshine hours could be expressed by the equation $Y =$

174.25X +193.17 (Fig. 1H). This was in conformity with the observations of Nag [12].

Relationship of nectar area with maximum relative humidity: A significant, negative and linear relationship ($r = -0.71$, $P < 0.05$) was came into light between maximum relative humidity and nectar area of artificially inseminated colonies of *A. mellifera L.* The data indicated 51 per cent variation in nectar area was found to be controlled by maximum relative humidity ($R^2 = 0.51$) where the equation $Y = -420.99X + 41875$ indicated the magnitude of the association (Fig. 1I). Similarly, the relationship between maximum relative humidity and nectar area of naturally inseminated colonies of *A. mellifera L.* was found to be negatively significant and linear ($r = -0.70$, $P < 0.05$). The data indicated that 49.1 per cent variation in nectar area was regulated by maximum relative humidity ($R^2 = 0.491$). The regression of nectar area on maximum relative humidity could be expressed by the equation $Y = -398.47X + 39515$ (Fig. 1J). Nag [12] also made similar observations of negative and significant impact of maximum relative humidity on nectar area.

3.2 Correlation of Brood Area, Pollen Area and Nectar Area with Different Weather Parameters of *A. mellifera L.*

During April'17 to March'18, correlation studies of brood area, pollen area and nectar area were again carried out with weather parameters in *A. mellifera L.* and the results discussed below (Table 2).

Relationship of pollen area with maximum temperature: The relationship between maximum temperature and pollen area of *A. mellifera L.* was found to be significant, negative and linear with artificially inseminated colonies ($r = -0.87$, $P < 0.05$) where 76.9 per cent variation in pollen area was observed to be controlled by maximum temperature ($R^2 = 0.769$). The regression equation $Y = -91.56X + 3443.8$ indicated the magnitude of this association (Fig. 2A). Similarly, the relationship between maximum temperature and pollen area of *A. mellifera L.* was found to be significant, negative and linear in naturally inseminated colonies ($r = -0.90$, $P < 0.05$) where 82.1 per cent variation in pollen area was observed to be controlled by maximum temperature ($R^2 = 0.821$). The regression of pollen area on maximum temperature could be expressed by the equation

$Y = -91.254X + 3314.4$ (Fig. 2B). Similar results were also obtained by Chaand et al. [13] in *A. mellifera L.*

Relationship of pollen area with maximum relative humidity: The relationship between maximum relative humidity and pollen area of *A. mellifera L.* was found to be significant, positive and linear with artificially inseminated colonies ($r = 0.57$, $P < 0.05$). The data indicated that 31.8 per cent variation in pollen area was found to be regulated by maximum relative humidity ($R^2 = 0.318$). The equation $Y = 84.73X + 7288.3$ indicated the magnitude of this association (Fig. 2C). Similarly, a significant, positive and linear relationship was existed between maximum relative humidity and pollen area of naturally inseminated colonies of *A. mellifera L.* ($r = 0.60$, $P < 0.05$). The data indicated 36.5 per cent variation in pollen area was observed to be controlled by maximum relative humidity ($R^2 = 0.365$). The regression of pollen area on maximum relative humidity could be expressed by the equation $Y = 87.607X + 7682$ (Fig. 2D). Gebremedhn et al. [14] also reported such observations in *A. mellifera L.* colonies.

Relationship of pollen area with total rainfall: The relationship between total rainfall and pollen area of *A. mellifera L.* was found to be significant, negative and linear with artificially inseminated colonies ($r = -0.74$, $P < 0.05$). The data indicated 55.1 per cent variation in pollen area was observed to be controlled by total rainfall ($R^2 = 0.551$). The regression of pollen area on total rainfall could be expressed by the equation

$Y = -1.841X + 1056$ (Fig. 2E). Similarly, the relationship between total rainfall and pollen area of *A. mellifera L.* was found to be significant, negative and linear with naturally inseminated colonies ($r = -0.76$, $P < 0.05$). The data revealed that 58.5 per cent variation in pollen area was regulated by total rainfall ($R^2 = 0.585$). The equation $Y = -1.829X + 933.58$ indicated the magnitude of this association (Fig. 2F). Such observations were in conformity with the results obtained by Chaand et al. [13].

Relationship of nectar area with maximum relative humidity: A significant, negative and linear relationship was existed between maximum relative humidity and nectar area of artificially inseminated colonies of *A. mellifera L.* ($r = -0.74$, $P < 0.05$). The data indicated 55.7 per cent variation in nectar area was observed to be controlled by maximum relative humidity

($R^2=0.557$). The regression of nectar area on rainfall could be expressed by the equation $Y= -343.63X +34290$ (Fig. 2G). Similarly, the relationship between maximum relative humidity and nectar area of *A. mellifera* L. was found to be significant, negative and linear relationship with naturally inseminated colonies ($r= -0.77$, $P<0.05$). The data indicated 59.8 per cent variation in nectar area was observed to be controlled by maximum relative humidity ($R^2=0.598$). The regression of nectar area on maximum relative humidity could be expressed by the equation $Y= -338.9X +33634$ (Fig. 2H). The results were in agreement with the findings of Gebremedhn et al. [14].


3.3 Correlation of Weather Parameters with Brood Area, Pollen Area and Nectar Area of *Apis cerana indica* F.

Correlation studies were also carried out in *A. c. indica* between weather factors and brood area, pollen area and nectar area during April'17 to March'18 and the results are discussed below (Table 3).


Relationship of brood area with maximum relative humidity: In *A. c. indica*, it was found that relative humidity has significant effect on brood area and has adverse and linear effect with artificially inseminated colonies ($r= -0.58$, $P<0.05$) where, 34.2 per cent variation in brood area was observed to be controlled by maximum relative humidity ($R^2=0.342$). As shown in figure 12A, the regression of brood area on maximum relative humidity could be expressed by the equation $Y= -248.82X +25463$ (Fig. 3A).

Similarly, the relationship between maximum relative humidity and brood area of *A. c. indica* was found to be significant, negative and linear with naturally inseminated colonies ($r= -0.60$, $P<0.05$) where, 36.3 per cent variation in brood area was observed to be regulated by maximum relative humidity ($R^2=0.363$). The regression of brood area on maximum relative humidity could be expressed by the equation $Y= -258.18X + 26175$ (Fig. 3B). Similar results were also obtained by Pastagia and Patel [15] and Painkra et al [16].


Relationship of nectar area with maximum relative humidity: Relative humidity revealed significant, negative and linear relationship with nectar area of artificially inseminated colonies of *A. c. indica* ($r= -0.66$, $P<0.05$). The data indicated 44 per cent variation in nectar area was observed to be controlled by maximum relative humidity ($R^2=0.440$). The regression of nectar area on maximum relative humidity could be expressed by the equation $Y= -122.31X +12277$ (Fig. 3C). Similarly, the relationship between maximum relative humidity and nectar area of *A. c. indica* was found to be significant, negative and linear with naturally inseminated colonies ($r= -0.70$, $P<0.05$). The data indicated 50.2 per cent variation in nectar area was observed to be controlled by maximum relative humidity ($R^2=0.502$). The regression of nectar area on maximum relative humidity could be expressed by the equation $Y= -130.22X +12901$ (Fig. 3D). Contrary to the present finding Painkra [16] documented positive significant correlation of maximum relative humidity with nectar area.




a) Relationship of brood area with maximum temperature in artificially inseminated colonies




b) Relationship of brood area with maximum temperature in naturally inseminated colonies




c) Relationship of pollen area with maximum temperature in artificially inseminated colonies




d) Relationship of pollen area with maximum temperature in naturally inseminated colonies




E. Relationship of pollen area with total rainfall in artificially inseminated colonies




F. Relationship of pollen area with total rainfall in naturally inseminated colonies




G. Relationship of pollen area with bright sunshine hours in artificially inseminated colonies



H. Relationship of pollen area with bright sunshine hours in naturally inseminated colonies




I. Relationship of nectar area with maximum relative humidity in artificially inseminated colonies




J. Relationship of nectar area with maximum relative humidity in naturally inseminated colonies


Fig. 1. Correlation of weather parameters with brood area, pollen area and nectar area of *Apis mellifera* L. (April'16 to March'17)




A) Relationship of pollen area with maximum temperature in artificially inseminated colonies




B) Relationship of pollen area with maximum temperature in naturally inseminated colonies




C) Relationship of pollen area with maximum relative humidity in artificially inseminated colonies




D) Relationship of pollen area with maximum relative humidity in naturally inseminated colonies




E) Relationship of pollen area with total rainfall in artificially inseminated colonies



F) Relationship of pollen area with total rainfall in naturally inseminated colonies



G) Relationship of nectar area with maximum relative humidity in artificially inseminated colonies



H) Relationship of nectar area with maximum relative humidity in naturally inseminated colonies

Fig. 2. Correlation of brood area, pollen area and nectar area with different weather parameters of *Apis mellifera* L. (April'17 to March'18)

Table 1. Correlation of brood area, pollen area and nectar area of *A. mellifera* L. L. with different weather factors

Month of observation	Brood area		Pollen area		Nectar area		Maximum temperature (°C)		Maximum relative humidity (%)		Total rainfall (mm)	Bright sunshine hours (hr)		
	AI	NI	AI	NI	AI	NI	AI	NI	AI	NI		AI	NI	
April'16	3829.8	3678.0	730.2	573.2	3619.8	3443.2	28.6		94		208.1	4.2		
May'16	4091.2	3882.4	629.8	519.4	3201.8	2903.4	30.4		94		249.7	4.1		
June'16	3517.8	3381.6	532.2	479.6	2333.8	2030.2	32.3		95		267.5	3.3		
July'16	2680.8	2524.6	497.4	362.2	1524.8	1323.8	33.0		96		576.0	4.9		
August'16	2427.2	2229.0	420.2	321.8	1233.6	1038.2	31.1		94		269.1	3.4		
September'16	2102.6	1946.2	436.8	342.6	992.2	847.8	32.2		96		249.2	2.0		
October'16	2039.2	1813.8	440.8	349.0	872.0	695.2	31.8		96		78.4	4.2		
November'16	2926.8	2507.0	527.8	486.4	845.0	654.4	28.8		97		16.5	5.5		
December'16	3449.8	3342.8	1287.4	1108.0	783.4	612.2	26.3		99		43.5	7.1		
January'17	3494.2	3368.4	1349.2	1295.2	671.8	556.0	25.2		98		0.1	7.5		
February'17	3581.8	3423.6	1052.4	922.8	832.8	625.2	26.8		95		39.4	5.6		
March'17	3672.6	3503.6	882.4	714.4	2712.6	2411.0	26.3		93		148.8	4.4		
Correlation with brood area (r=)							AI	NI	AI	NI	AI	NI	AI	NI
Correlation with pollen area (r=)							-0.57*	-0.57*	-0.20	-0.19	-0.19	-0.15	0.38	0.37
Correlation with nectar area(r=)							-0.89*	-0.88*	0.45	0.49	-0.57*	-0.60*	0.83*	0.84*
							0.10	0.09	-0.71*	-0.70*	0.36	0.35	-0.37	-0.36

* = Significant at 5% level of probability

Table 2. Correlation of brood area, pollen area and nectar area of *A. mellifera* L. L. with different weather factors


Month of observation	Brood area		Pollen area		Nectar area		Maximum temperature (°C)	Maximum relative humidity (%)		Total rainfall (mm)	Bright sunshine hours (hr)			
	AI	NI	AI	NI	AI	NI		AI	NI		AI	NI		
April'17	3888.6	3744.2	823.4	616.6	3620.8	3321.2	29.7	91		111.8		4.5		
May'17	4203.8	3836.8	758.0	595.2	3145.0	2989.8	30.7	92		226.9		4.2		
June'17	3674.8	3421.4	581.8	386.8	2391.2	2071.6	32.8	95		399.2		3.7		
July'17	2842.6	2564.0	478.2	384.6	1438.6	1286.4	33.6	94		275.1		4.0		
August'17	2376	2123.4	427.2	342.6	1310.2	1172.2	31.1	94		269.1		3.4		
September'17	2113.6	1974.6	419.0	340.4	1006.0	865.8	32.7	94		324.4		3.3		
October'17	2018.8	1944.2	395.4	337.2	867.8	742.4	31.1	96		192.8		5.2		
November'17	2953.0	2729.8	569.8	428.8	818.0	640.8	28.4	95		14.7		7.4		
December'17	3426.4	3239.4	1305.6	1115.0	796.0	605.0	25.8	99		0.0		5.4		
January'18	3483.0	3317.0	1382.2	1226.8	664.4	510.0	23.9	98		2.7		4.7		
February'18	3556.4	3423.2	1021.0	961.6	848.6	615.0	24.5	97		29.1		3.4		
March'18	3583.6	3463.0	971.6	952.4	2835.8	2436.0	27.3	95		76.0		4.0		
							AI	NI	AI	NI	AI	NI		
Correlation with brood area (r=)							-0.39	-0.45	-0.12	-0.08	-0.29	-0.35	-0.01	-0.01
Correlation with pollen area (r=)							-0.87*	-0.90*	0.57*	0.60*	-0.74*	-0.76*	0.10	0.05
Correlation with nectar area(r=)							0.28	0.31	-0.74*	-0.77*	0.25	0.27	-0.25	-0.25

*= Significant at 5% level of probability


Table 3. Correlation of brood area, pollen area and nectar area of *A. c. indica* with different weather factors

Month of observation	Brood area		Pollen area		Nectar area		Maximum temperature (°C)		Maximum relative humidity (%)		Total rainfall (mm)		Bright sunshine hours (hr)	
	AI	NI	AI	NI	AI	NI	AI	NI	AI	NI	AI	NI	AI	NI
April'17	2971.8	2837.2	1055.2	834.0	1328.8	1194.8	29.7		91		111.8		4.5	
May'17	3337.0	3250.4	839.8	645.0	1370.2	1257.2	30.7		92		226.9		4.2	
June'17	2850.4	2717.4	774.4	571.4	829.4	773.2	32.8		95		399.2		3.7	
July'17	1671.6	1559.4	569.8	515.8	548.8	451.2	33.6		94		275.1		4.0	
August'17	1443.6	1324.2	529.4	472.6	455.8	418.0	31.1		94		269.1		3.4	
September'17	1309.0	983.2	522.6	452.8	388.2	295.2	32.7		94		324.4		3.3	
October'17	735.0	607.8	464.4	306.2	265.4	105.2	31.1		96		192.8		5.2	
November'17	622.8	466.8	285.6	254.8	228.4	79.8	28.4		95		14.7		7.4	
December'17	539.0	427.2	352.2	302.4	179.0	48.0	25.8		99		0.0		5.4	
January'18	1421.2	1122.8	827.4	593.2	461.2	299.0	23.9		98		2.7		4.7	
February'18	2368.2	2153.6	966.6	881.2	679.8	484.2	24.5		97		29.1		3.4	
March'18	2632.6	2319.2	1233.8	1099.2	1158.6	958.4	27.3		95		76.0		4.0	
							AI	NI	AI	NI	AI	NI	AI	NI
Correlation with brood area (r=)							-0.13	-0.16	-0.58*	-0.60*	-0.27	-0.29	-0.53	-0.50
Correlation with pollen area (r=)							-0.25	-0.27	0.27	0.25	-0.09	-0.13	-0.49	-0.51
Correlation with nectar area(r=)							-0.09	-0.17	-0.66*	-0.70*	-0.13	-0.23	-0.37	-0.41


* = Significant at 5% level of probability




A) Relationship of brood area with maximum relative humidity in artificially inseminated colonies



B) Relationship of brood area with maximum relative humidity in naturally inseminated colonies



C) Relationship of nectar area with maximum relative humidity in artificially inseminated colonies



D) Relationship of nectar area with maximum relative humidity in naturally inseminated colonies

Fig. 3. Correlation of weather parameters with brood area, pollen area and nectar area of *Apis cerana indica* F. (April'17 to March'18)

4. CONCLUSION

The correlation study of *A. mellifera* during the year 2016-17 revealed significant negative correlation of brood area ($r = -0.57$ in both AI and NI colonies) and nectar area ($r = -0.71$ in AI and $r = -0.70$ in NI colonies) with maximum temperature and maximum relative humidity, respectively. Whereas, pollen area was found to have significant positive correlation with bright sunshine hours (0.83 in AI and 0.84 in NI colonies) and significant negative correlation with maximum temperature ($r = -0.89$ in AI and $r = -0.88$ in NI colonies) and total rainfall ($r = -0.57$ in AI and $r = -0.60$ in NI colonies). Again in 2017-18, pollen area showed positive significant correlation with maximum relative humidity ($r = 0.57$ in AI and $r = 0.60$ in NI colonies) and negatively significant correlation with maximum temperature ($r = -0.87$ in AI and $r = -0.90$ in NI colonies) and total rainfall ($r = -0.74$ in AI and $r = -0.76$ in NI colonies). Nectar area was found to have negatively significant correlation with maximum relative humidity ($r = -0.74$ in AI and $r = -0.77$ in NI colonies). *A. cerana indica* showed

significant negative correlation of brood area ($r = -0.58$ in AI and $r = -0.60$ in NI colonies) and nectar area ($r = -0.66$ in AI and $r = -0.70$ in NI colonies) with maximum relative humidity during 2017-18.

ACKNOWLEDGEMENT

The authors are thankful to Retd. Dr. Ataur Rehman, Department of Entomology and Retd. Dr. D. Das Borah, Department of Agricultural Statistics, Assam Agricultural University, Jorhat for their sincere help during the period of investigation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bisht DS. Pollen gathering activity of honey bees and *Halictus* sp. in roses, Indian Bee J. Keywords: *cerana / florea / honey / pollen / India*. 1975;37:1-3.

2. Bisht DS, Naim M. Pollen gathering capacity of the Indian honeybee, *Apis cerana* F., *Indian J. Entomol.* Keywords: *cerana* / foraging / pollen / India. 1979;41:295-296.
3. Das PK, Rahman A. Brood rearing activity of *Apis cerana indica* F. In Assam. *Crop. Res.* 2000;19(3): 469-473.
4. Calovi M, Grozinger CM, Miller DA, Goslee SC. Summer weather conditions influence winter survival of honey bees (*Apis mellifera*) in the northeastern United States. *Scientific Reports.* 2021;11:1553. Available: <https://doi.org/10.1038/s41598-021-81051-8>.
5. Reddy CC. The influence of various weather factors on the foraging behaviour of *Apis cerana indica*. *Proc. 4th Int. Symp. Pooln. College Park, Myrland, U.S.A.* 1979:341-345.
6. Verma S. Studies on foraging behaviour of *Apis cerana indica* F. in Jeolikote, Nainital, India. *Indian Bee J.* 1983;45:5-9.
7. Mehrotra KN, Bisht DS. Current status of the biology, physiology and behaviour of the Indian hive bee, *Apis cerana indica* Fab. *Indian Bee J.*, 1984;46:36-42.
8. Abrol DP, Bhat AA. Influence of atmospheric conditions and floral rewards on diurnal activity pattern, flower visitation rates and pollinating effectiveness of *Apis cerana indica* foragers. *Res. Develop. Reporter.* 1987;4(1):13-16.
9. Panse VG, Sukhatme PV. *Statistical methods for Agricultural Workers*, ICAR, New Delhi. 1985:87-93.
10. Rahman S. Comparative brood rearing and pollen gathering activities of *Apis cerana indica* F. and *Apis mellifera* L. under Jorhat conditions of Assam. Thesis, Assam Agricultural University, Jorhat; 1992.
11. Zapart KD, Chlopek K, Niedzwiedz T. The impact of meteorological conditions on the concentration of alder pollen in Sosnowiec (Poland) in the years 1997-2017. *Aerobiologia.* 2018;34:469-485.
12. Nag YK. Performance of *Apis mellifera* L. under different migratory mode in Northern parts of India. Thesis, Indira Gandhi Krishi Vishwavidyalaya, Raipur; 2016.
13. Chaand D, Sharma D, Ganai SA, Norboo T, Sharma S. Effect of colony strength on colony build up and foraging activity of *Apis mellifera* L. *J. Entomol. Zool. Stud.* 2017;5(6):1369-1373.
14. Gebremedhn HN, Tadesse A, Belay T. Flight intensity of honeybees (*Apis mellifera*) and its relationship with temperature, sunshine hours, cloudiness and relative humidity. *Livestock Research for Rural Development.* 2014; 26 (11).
15. Pastagia JJ, Patel MB. Influence of weather parameters on brood rearing and foraging activities of Indian bee, *Apis cerana*. *AGRES – An International Journal.* 2014;3:403-409.
16. Painkra MK, Shaw SS, Thakur RK, Nag Y. Influence of Weather Parameters on Colonies Performance Parameters of Indian Bee, *Apis cerana indica*, *International Journal of Current Microbiology and Applied Sciences* ISSN: 2319-7706. 2017;6:2375-2380.

© 2022 Borah and Deka; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/84865>