



# Structural Analysis of the Bagoué Furrow (South Boundiali) in the Northwest of Ivory Coast: Fracturing, Kinematics and Chronology

Lipoublida Djagre <sup>a\*</sup>, Barthélémy Gnammytchet Koffi <sup>a</sup>  
and Gbele Ouattara <sup>a</sup>

<sup>a</sup> *Department of Earth Sciences and Mining Resources (STeRMi) / Civil Engineering Laboratory, Geosciences and Geographical Sciences / Institut National Polytechnique Félix Houphouët-Boigny, Yamoussoukro, Côte d'Ivoire.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95982>

**Original Research Article**

**Received: 25/10/2022**  
**Accepted: 29/12/2022**  
**Published: 31/12/2022**

## **ABSTRACT**

Structural mapping from spatial imagery is an important technique in mastering lineament fracture networks. The purpose of this work is to make a structural analysis of the Bagoué furrow, using spatial imagery and field data. The methodological approach adopted in the work consists in the pre-processing and digital processing of images, and the extraction of lineaments by means of automatic and manual extraction techniques. Five major lineament directions are identified: the N-S, NNE-SSW, NE-SW, E-W and NW-SE directions. The field data is used to define their kinematics and chronology. This data reveal that the lineaments corresponding to the N-S trending fractures are intersected by the NE-SW trending fractures, which in turn are cut by the NNE-SSW trending fractures. The latter are intersected by the lineaments corresponding to the SE-NW trending

\*Corresponding author: Email: [lipoublida.dja@gmail.com](mailto:lipoublida.dja@gmail.com);

fractures. The SE-NW trending fractures are intersected by the E-W trending fractures. Ultimately, the N-S trending fractures are considered the oldest, followed by the NE-SW, NNE-SSW, SE-NW and E-W trending fractures.

**Keywords:** Lineament structures; Landsat 8; radar; Bagoué; Ivory Coast; West Africa.

## 1. INTRODUCTION

Structural analysis in geological sciences is of paramount importance in the exploration of mineral resources [1]. However, in tropical regions such as Côte d'Ivoire, it is not an easy task to identify lineament structures because of the thick lateritic cover over the geological substratum. The structural study from lineament mapping using spatial imagery data processing techniques has therefore become a considerable approach in geological studies. Lineaments are thus regarded as important structural and tectonic indicators to determine regional tectonic trends and fracture zones in rocks [2]. These structural features can then constitute preferential flow paths for hydrothermal fluids and groundwater [3-5]. By offering a synoptic view, satellite imagery allows for the study of vast geographical areas and it is also a powerful tool in analyzing fracturing [6].

The main objective of this work is to carry out a structural analysis of the Bagoué furrow using spatial images. This will specifically include the mapping of all the fracture networks of the Bagoué furrow (South Boundiali) by extracting the lineament networks from satellite images while validating those using structural field data, and then the establishment of their kinematics and chronology relative to the regional scale.

### 1.1 Study Area and Geological Context

The Bagoué furrow is located in the northwest of Côte d'Ivoire (West Africa) and more precisely in the Bagoué region. It is therefore located between longitudes 6°00 and 7°00 W and latitudes 8°00 and 10°00 N. In this area, the water regime is known as the transitional tropical regime, with the Bagoué River as its main watercourse.

The Bagoué Trench is an entity of the Boundiali-Bagoé Birimian unit, whose geology is an integral part of the Baoulé-Mossi domain of the West African Craton. The geological formations are of Paleoproterozoic era and are characterized by a succession of bands of schistose rocks, migmatitic rocks and plutonic rocks. Thus, one

can observe from west to east of the complex of undifferentiated granites with some areas of biotite granites, granites with marked migmatitic facies and porphyroid granite. In a few places, these granites are associated to basic rocks [7,8]. Next to these granites, there is a zone of undifferentiated schists and sericite schists. The geological formations are also characterized by the presence of a band of biotite granite, and granitoids and calc-alkaline granite with amphibole in the southeast (Fig.1).

## 2. MATERIALS AND METHODS

### 2.1 Materials

In this study, the mapping databases used include the topographical and geological maps of the Bagoué region at 1:200,000 scale and then the field data. The data used for the extraction of the linear network are from orthorectified Landsat 8 OLI optical images composed of four scenes (197-053 of 29/12/2019, 197-054 of 29/12/2019, 198-053 of 04/12/2019 and 198-054 of 04/12/2019) covering the study area, and also from radar images (SAT and ERS radar).

The geographic information system software used in this study includes: Envi 4.7, used to carry out various tasks of image preprocessing and processing; PCI's Geomatica for the automatic extraction of the lineament network through its LINE module; Arcgis 10.4, used to extract the statistical parameters of the lineaments and to dress maps. To these software we also used software such as Adobe Illustrator to digitize the lineaments extracted manually then Georient and Rockwork 17 for the realization of the directional rosettes.

### 2.2 Methodology

Two methods for extracting lineaments from spatial images were used in this work: automatic extraction by the use of the LINE module of Geomatica and visual extraction by which lineaments are detected by visual interpretation. The preprocessing and processing techniques have allowed for the automatic and manual extraction of lineaments from OLI images and radar images.

The methodological approach adopted for the mapping of geological lineament structures is summarized in the following diagram (Fig. 2).




Fig. 1. Geological map of the Bagoué furrow as modified by [9]




Fig. 2. Landsat 8 image and radar images processing methodology

### 3. RESULTS AND DISCUSSION

The mapping in our study area initially consisted in the digital processing of Landsat 8 optical satellite images and radar images, of which some results are shown in Fig.3.

#### 3.1 Extraction of Lineaments

In this study, the extraction of lineaments is based on Landsat 8 OLI satellite images along with radar images (SAT and ERS radar). This is

carried out both by automatic and manual extraction techniques.

The results obtained from the automatic extraction technique have allowed for the identification of 12930 lineaments (Fig. 4). This Fig. 4 shows that the automatically detected lineament map has a large number of structures to show the different directions of the geological lineaments, but only the N-S and NE-SW directions of the lineaments are well discriminable.





Fig. 3. Landsat 8 OLI image and radar image processing. (A): Color-coded (567) OLI image; (B): Image filtered by Gaussian high pass filter applied on band 8 of OLI image; (C): Radar SAT image; (D): Radar ERS image (negative print)



**Fig. 4. Map of the automatically detected lineaments**

The lineament texture analysis of the maps generated from manual extraction (Figs. 5 and 6) shows the following geometric shapes:

- Arc-shaped lineaments, ovoid to radial representing periplutonic to plutonic areas [10,11],
- Large rectilinear lineaments that most often correspond to large regional fractures, and
- Small straight lineaments.

Of the two lineament maps deriving from manual extraction, that of SAT radar image reveals more lineament structures with arc-like shape than that of ERS radar images. The latter also has fewer lineament structures compared to the former.

The manual extraction technique is used to extract lineaments from SAT and ERS radar imagery. This technique was useful in detecting long geological lineaments in the NE and NW directions and medium lineaments in the E-W and NS directions of the study area in both radar images (Figs. 5 and 6).

### 3.2 Directional Statistical Analysis of the Lineament Maps

The directional statistical analysis of the synthetic maps of the structural lineaments extracted from our study area (Fig. 7), shows a dispersion of structural lineaments in all directions ( $N0^\circ$  to

$N180^\circ$ ). Among these directions, five major groups of directions stand out in the diagrams that discriminate lineament orientations of the maps in Fig. 5 and 6: 1) N-S direction lineaments ( $N00^\circ$  to  $N10^\circ$ ), 2) NNE-SSW direction lineaments ( $N30^\circ$  to  $N40^\circ$ ), 3) NE-SW direction lineaments ( $N40^\circ$  to  $N50^\circ$ ), 4) E-W direction lineaments ( $N90^\circ$  to  $N100^\circ$ ), and 5) SE-NW direction lineaments ( $N130^\circ$  to  $N140^\circ$ ); and six minor direction groups:  $N10^\circ$  to  $N30^\circ$ ,  $N50^\circ$  to  $N70^\circ$ ,  $N70^\circ$  to  $N90^\circ$ ,  $N110^\circ$  to  $N120^\circ$ ,  $N120^\circ$  to  $N130^\circ$  and  $N140^\circ$  to  $N170^\circ$ .

The directional statistical analysis of the map of the automatically extracted lineaments (Fig. 7A) allows us to highlight four classes of preferential directions: 1) the class of N-S direction ( $N00^\circ$  to  $N10^\circ$ ), 2) the class of NE-SW direction ( $N40^\circ$  to  $N50^\circ$ ), 3) the class of E-W direction ( $N90^\circ$  to  $N100^\circ$ ) and 4) the class of NW-SE direction ( $N30^\circ$  to  $N40^\circ$ ).

The overall directional analysis of the three maps shows the following:

- The lineament maps in Fig. 4 and 6 show a significant number of lineaments with N-S and E-W directions compared to the lineament map in Fig. 5,
- The maps in Figs. 5 and 6 show more discriminable and concordant lineament directions compared to those of the map in Fig. 4,

- The NNE-SSW and NE-SW lineaments in Fig. 5 are more numerous than those of the lineaments in Fig. 6. Furthermore, they are more numerous in the E-W direction in Fig.6 than in Fig. 5,
- The map in Fig. 4 is highly saturated.

As a result of the directional statistical analysis of the lineament maps, one of the maps of the lineament extracted manually is retained and used for the control and validation of the lineaments. This is the lineament map extracted from radar SAT image (Fig. 5) which shows almost all the lineament structures that can be found in the field.

### 3.3 Control and Validation of Lineaments

The control and validation phase of the geological lineaments of our study area consisted in comparing the preferential directions of the lineaments extracted manually from the SAT radar image and in confronting them with the results of field works. This control phase is essential in assessing the relevance of the chosen extraction method and the results obtained.

Using field data in control phase is an adequate means to validate structural lineaments. The study site which is located in a basement

environment with a subhumid tropical climate, is largely covered with lateritic crust and laterite, which explains the scarcity of outcrops in this area. Few data have been collected in the field regarding this rarity of outcrops. The geotraverse carried out in the field made it possible to observe metasediments in Tarato in which schistosity networks develop with fractures trending N160-60W, N60-50NW (Fig. 9C). The N160° fractures intersect those of N60° following a sinistral movement. In Foutio, not far from Tarato, they are schistosity networks with fracture N60° intersected by N170° schistosity and N160 quartz veins dipping northeast which intersect the N90° quartz veins dipping vertically (Fig. 10A). In the field, several measurements were taken which led to the realization of the rose diagrams that discriminate the directions of fractures. The rose diagram (Fig. 8) shows seven families of fractures observed in the study area: 1) fractures trending N0° to N20°; 2) fractures trending N30° to N40°; 3) fractures trending N50° to N60°; 4) fractures trending N70° to N100°; 5) fractures trending N120° to N130°; 6) fractures trending N140° to N150 and 7) fractures trending N160° to N170°. Among these families, there are four preferential directions: fractures trending NNE-SSW (N10° to N20°), fractures trending NE to SW (N50° to N60°), fractures trending EW (N80° to N90 °) and fractures trending SE-NW (N120° to N130°).




Fig. 5. Map of the lineaments extracted manually from radar SAT image




Fig. 6. Map of the lineaments extracted manually from radar ERS image





Fig. 7. Directional statistical analysis. (A): Rose diagram of the lineaments extracted automatically; (B): Rose diagram of the lineaments extracted manually from radar Sat image; (C): Rose diagram of the lineaments extracted manually from radar ERS image



**Fig. 8. Directional rose of the fractures in the study area**

The spatial analysis in the field reveals that the NE direction fractures are intersected by the NNE direction fractures following a sinistral shear. The latter are in turn intersected by fractures trending SE in a sinistral shear context. SE trending fractures are intersected by E-W trending fractures by dextral shear. N-S trending fractures are intersected by NE trending fractures following a sinistral shear.

From a chronological point of view, N-S fractures are older, successively followed by NE-SW, NNE-SSW, SE-NW fractures and E-W fractures which are considered as the youngest according to the intersection principle.

The validation of the lineaments extracted by analogical analysis consisted in comparing the statistical data of the lineament network with those of the field and in establishing genetic links between them. In order to assess the contribution of the spatial imagery data, the lineaments extracted from them were compared to the structural data collected in the field. Two objectives are considered for this comparison: the reliability of the interpretation of the lineament analysis and the spatial precision of the data collected in the field.

The field data, by analogy with the orientations of lineaments, allowed to draw up the following synthesis: the direction families deriving from the lineament map are similar to those shown by the field data on the directional rose diagram

(Fig.11). Nevertheless, some lineaments from satellite images are not remarkable with the structural elements of the terrain. These lineaments belong to the following directional classes:  $N20^{\circ}$  to  $N30^{\circ}$ ,  $N40^{\circ}$  to  $N50^{\circ}$ ,  $N60^{\circ}$  to  $N70^{\circ}$ ,  $N100^{\circ}$  to  $N110^{\circ}$ ,  $N120^{\circ}$  to  $N130^{\circ}$ ,  $N140^{\circ}$  to  $N150^{\circ}$  and from  $N170^{\circ}$  to  $N180^{\circ}$ . Other families of lineaments, on the other hand, are superimposable on those of the structural elements of the terrain; these are particularly the families of directions  $N00^{\circ}$  to  $N20^{\circ}$ ,  $N30^{\circ}$  to  $N40^{\circ}$ ,  $N50^{\circ}$  to  $N60^{\circ}$ ,  $N70^{\circ}$  to  $N100^{\circ}$ ,  $N110^{\circ}$  to  $N120^{\circ}$ ,  $N130^{\circ}$  to  $N140^{\circ}$  and from  $N150^{\circ}$  to  $N170^{\circ}$ .

Following the validation of the lineaments by structural field data, the lineament map resulting from the SAT radar image represents the final map of the fractures affecting the Bagoué furrow.


### 3.4 Discussion

Satellite image processing for the purpose of structural mapping made it possible to detect the lineament structural elements of the Bagoué furrow. To ensure the reliability of this mapping technique, the structural field data was checked by reference to the lineaments extracted from spatial imagery through the comparison of the directional rose diagrams. The rose diagrams analysis shows that some families of lineaments extracted from satellite images are perfectly superimposable on the structural elements of the field, others are slightly offset and they do not




appear on the rose diagram of the field data and other families are in the same extension as the latter but are not superimposable on them. The discrepancy between satellite lineaments and the structural elements of the field is explained by the geocoding and resampling effect which cause inaccuracy in the processed images [12-14]. The absence of certain families of lineaments (N20° to N30°, N40° to N50°, N60° to N70°, N100° to N110°, N120° to N130° and N140° to N150°) on the rose diagram of the field data can be attributable to the scarcity of outcrops which did not allow for the observation of these families of structures during the geotraverse in the field. The N20° to N30° lineament classes have been highlighted on the kolia granite outcrops by [15]. Lineament classes N120° to N130° were observed by this same author on the Boundiali arkoses. These lineaments families which were not noticed in the field during our study were observed on the Bagoué furrow by [9] and [15]. From the above analyses, the similarity between the directional rose diagrams of the lineaments with the directional rose diagrams of the field data and the previous studies carried out on the Bagoué


furrow do confirm that the lineaments extracted from satellite images correspond to geological fractures. The lineament map (Fig 5) resulting from manual extraction using SAT radar imagery is then the structural map of the Bagoué furrow. On this map, the arc-shaped lineaments, ovoid to radial which can be observed in the northeast correspond to the relics of the periplutonic and plutonic zones whose laminations constitute real gold traps [10]. The directional analysis of the lineaments highlights five preferential directions: NS (N00° to N10°), NNE-SSW (N30° to N40°), NE-SW (N40° to N50°), EW (N90° to N100 °) and NE-SW (N130° to N140°). The NNE-SSW trending Lineaments might correspond to the direction of orientation of the general schistosity indicated by the mineral schistosity of direction N20° which can be observed on the granitoid outcrops located in the South-west of the study area and the schistosity of N20° to N30° trending fractures observed in volcano-sedimentary and sedimentary rocks. NE-SW trending lineaments define the direction of the high to medium density lineaments on the ground. They correspond to the Yérétiélé faults described in the Bagoué furrow by [15].



**Fig. 9. (A): Outcrop of fractured metasediments with intersecting fractures; (B): Outcrop of diorite affected by the N20° fractures intersecting the N40° fractures in dextral shear; (C): Folded metasediments presenting fractured networks, the N160° networks intersect the N60° networks; (D): Outcrop of fractured gneiss with N150° fractures intersecting the N20° fractures**



**Fig. 10.** Lithological and structural data about some of the outcrops in the study area. (A) : Outcrop of metasediments in which the N160 quartz veins intersect the N90° veins (B) : Fractured metasediment with N170° and N60° fractures intersecting; (C) : Fractured biotite granite outcrop, the N20° fractures are intersected by the N140° fractures and both fracture networks are intersected by N80° fractures; (D): Fractured biotite granite outcrop with directions: N20° intersected by N140°, N40° intersected by N20°



**Fig. 11.** Comparison of the directional rose diagrams. (A): the lineaments extracted from SAT radar image; (B): field data

#### 4. CONCLUSION

For the structural analysis of the Bagoué furrow using space imagery, we had to use Landsat 8 OLI satellite images together with radar images. The chosen methodological approach to achieve the objective of this study is based on image pre-processing and processing operations in order to enhance the images and extract lineament structures. The automatic and manual lineament extraction techniques led us to retain the lineament map manually extracted from the SAT radar image because of its conformity with the geological structures of the area of interest.

The statistical analysis of the lineament network using the directional rose diagrams showed a particular directional distribution of these lineament structures. They are organized into five main directional systems: N-S, NNE-SSW, NE-SW, E-W and NW-SE. These lineament directions correspond to zones with medium to high fracture density. From a chronological point of view, the lineaments corresponding to the N-S trending fractures are intersected by the NE-SW trending fracture zones moving in a sinister direction, which in turn are intersected by the NNE-SSW direction fracture zones moving in a sinister direction. The latter are intersected by the lineaments of the SE-NW trending fracture zones moving in a sinister direction. The SE-NW fracture zones are intersected by the E-W fractures in a dextral movement. In the end, the lineaments corresponding to the N-S direction fracture zones are considered to be the oldest, followed successively by the NE-SW, NNE-SSW, SE-NW and E-W direction fracture zones, which are considered to be the youngest.

The resulting structural map obtained using spatial imagery will constitute the structural map of the Bagoué trench and will help to update the previous structural maps of the area. It will serve as a guide for mineral and water resource exploration activities.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Marghany M, Hashim M. Lineament Mapping Using Multispectral Remote Sensing Satellite Data, Research J. of Applied Sciences. 2010;5(2):126- 130.
2. El-Sawy ESK, Ibrahim M, El-Bastawesy MA, El-Saud WA. Automated, manual lineaments extraction and geospatial analysis for Cairo-Suez district (Northeastern Cairo-Egypt), using remote sensing and GIS. 2016;3(5):10.
3. Christian AG, Sekouba O, Alexis KK., Bachir M, Fernand KK. Extraction automatique des linéaments à l'aide d'images satellitaires optique et radar en milieu de socle précambrien (Haute Marahoué, Côte d'Ivoire. 2019;8(01): 24-32.
4. Nouayti M, Khattach ND, Hilali M. Cartographie des zones potentielles pour le stockage des eaux souterraines dans le haut bassin du Ziz (Maroc) : Apport de la télédétection et du système d'information géographique. 2017;13.
5. Zazoun RS, Marok A, Samar L, Benadla M, Mezlah H. La fracturation et les bandes de déformation dans la région d'El Kohol (Atlas saharien central, Algérie): Analyse fractale, lois d'échelles et modèle de réseaux de fractures discrètes. 2015; 71(2):39. DOI:10.3989/egeol.42011.359.
6. Millogo C, Nakolendoussé S, Saga S. Cartographie des accidents géologiques par imagerie satellitaires et géophysiques aéroportée, et analyse statistique des réseaux de fractures du socle birimien du bassin versant du lac bam (centre nord du Burkina); 2018.
7. Beaudou AG, Sayol R. Etude pédologique de la région de Boundiali - Korhogo (Côte d'Ivoire): cartographie et typologie sommaire des sols, feuille Boundiali, feuille Korhogo à 1/200.000. 1980;58.
8. Eschenbrenner V, Badarello L. Étude pédologique de la région d'Odienné (Côte d'Ivoire) : carte des paysages morpho-pédologiques, O.R.S.T.O.M., Paris ; 1978.
9. Turner P. Evolution of the early Proterozoic Boundiali-Bagoe Supracrustal Belt and associated granitic rocks, northern Côte d'Ivoire, West Africa. Ph.D., University of Portsmouth; 1995.
10. Niamke K H. Télédétection et SIG dans la stratégie de prospection 1 minière en Côte d'Ivoire : cas de l'indice d'or baoulé-est 1 de la région du n 'Zi Comoé (centre de la Côte d'Ivoire) », université de Cocody ; 2008.
11. Gbele O. Structure du batholite de Ferkessedougou (secteur de zuenoula,

- Côte d'Ivoire) implications sur l'interprétation de la géodynamique du Paléoprotérozoïque d'Afrique de l'ouest à 2.1 Ga, Thèse de doctorat de l'Université d'Orléans.1998;344.
12. Guillaume R. Développement et évaluation de nouvelles méthodes de classification spatiale-spectrale d'images hyperspectrales. Thèse de doctorat de l'université de Toulouse. 2012 ;148.
  13. Sghaier MO. Extraction des structures linéaires à partir des images satellitaires à très haute résolution pour l'aide à la gestion des catastrophes majeures. Thèse de doctorat, École de technologie supérieure université du Québec. 2017;270.
  14. Yao KA. Le volcanisme du sillon de Boudali, phénomène principal du protérozoïque inférieur de cette région NNW de la Côte d'Ivoire. Thèse de doctorat, Université de Clermont-Ferrand 2. 1993;201.
  15. Gnammytchet Barthélémy Koffi. Possibilités et limites des images radar en cartographie géologique. Application en Côte d'Ivoire ». Thèse de doctorat, Université de Paris 6 ; 1998.

---

© 2022 Djagre et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/95982>