



Prevalence of Bacterial and Fungal Isolates Associated with Road Traffic Accident In-patients in General Hospitals in Niger State

P. T. Chukwuka¹, S. A. Garba², M. E. Abalaka² and V. C. Chukwu^{2*}

¹General Hospital, Minna, Niger State, Nigeria.

²Department of Microbiology, School of Life Sciences, Federal University of Technology, Minna, Niger State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author PTC designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors SAG and MEA managed the analyses of the study. Author VCC managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/MRJI/2019/v27i530111

Editor(s):

(1) Dr. Eliton da Silva Vasconcelos, Department of Physiological Sciences, Federal University of São Carlos – UFSCar, Rod. Washington Luiz, São Carlos, SP, Brazil.

Reviewers:

- (1) K. Anandakumar, Swamy Vivekanandha College Of Pharmacy, Tiruchengode, India.
(2) Leticia Ximena Barrientos Díaz, Universidad de La Frontera, Temuco, Chile.
(3) Vivek Kumar Singh, Public Health and Infectious Disease Research Center (PHIDReC), Kathmandu, Nepal.
Complete Peer review History: <http://www.sdiarticle3.com/review-history/48687>

Received 15 February 2019

Accepted 03 May 2019

Published 16 May 2019

Original Research Article

ABSTRACT

Aim: This study was carried out to determine the prevalence and antimicrobial susceptibility of bacteria and fungi isolated from wounds of in-patients with road traffic accidents at four General hospitals in Niger State.

Place and Duration of Study: Department of Microbiology, Federal University of Technology, Minna, Nigeria, between October 2017 and May 2018.

Methodology: Wound swabs were taken twice per patient, first swab at contact and second swab taken seven days post wound dressing. The outcome of 1000 wound swabs taken from 409 (267 male and 142 female) in-patients with road traffic accident wound treated in the General hospitals Bida, Minna, Kontagora and Suleja areas of Niger State, Nigeria.

Results: The results showed that 405 (40.5%) samples were positive for bacteria and fungi pathogens while 595 (59.5%) were negative. 262 (80.6%) swabs yielded single isolates while 46

*Corresponding author: E-mail: victorchukwuc@gmail.com;

(14.2%) yielded double organisms and 17 (5.2%) yielded three or more organisms. Of the 405 positive cultures, 347 (85.7%) were bacterial isolates while 58 (14.3%) were fungal isolates. *Staphylococcus aureus* predominates (37.7%), followed by *Pseudomonas aeruginosa* (15.6%), *E. coli* (11.4%), *Streptococcus pyogenes* (9.4%), *Bacillus subtilis* (7.2%), *Klebsiella pneumonia* (4.4%), *Aspergillus niger* (4.2%), *Candida albicans* (3.7%), *Aspergillus flavus* (2.7%), *Candida pseudotropicalis* (2.2%) and *Mucor pusillus* (1.5%).

Conclusion: Most bacterial isolates were sensitive to Levofloxacin, Ciprofloxacin, Streptomycin and Gentamycin, while others showed some degree of resistance to the remaining antibiotics. *Klebsiella pneumoniae* showed the highest resistance to all the antibiotics used. All the fungi isolated were highly sensitive to the antifungal drugs used except Griseofulvin.

Keywords: Bacteria; fungi; RTA wounds; antibiotics; general hospitals.

1. INTRODUCTION

Road traffic accident results in 1.2 million deaths and injuries about four times this figure worldwide every year [1]. The causes of traffic crashes and fatalities are varied. Three major categories have traditionally been identified as human, vehicle and highway infrastructure. Amongst these, the road user attitude and human disability have been found to account for more than 85 per cent [2]. Road accidents result in trauma. About half of deaths due to trauma are within the age range 15–45 years, and males are twice commonly affected than females [3]. Trauma is the fifth leading cause of significant disability [4], the sixth leading cause of death worldwide. The commonest causes of death due to trauma are central nervous system injury, followed by substantial blood loss [4]. Road traffic accident wound infection are wound infections sustained after automobile accident. Wound infection is caused by microorganisms ranging from bacteria to fungi [5]. Main predisposing factor to infection include poor personal hygiene, equipment, the environment and patients risk factors.

Despite the use of Modern technologies in the management of wound in our hospitals, wound infection still remains the most common cause of hospital infection; and are associated with increased morbidity and death [6-8]. In our local settings, the clinicians are faced with challenges on issues regarding wound infection due to road traffic accident injury [9]. There is the problem of poor or improper handling of road traffic accident victims with injury, the presence of nosocomial infection during admission in the hospital, the possibility of drug-resistant organisms with its resultant sequels such as prolongation of hospital stay, increasing cost of treatment and a possible loss of function as a complication [9].

Because of the high prevalence of wound colonization and / or infection in road accident

victims with injuries, it is important to study the different antimicrobial resistance patterns and the selective usage of antimicrobials only when absolutely necessary so as to avoid changing the normal flora of the skin, in order not to result in multidrug-resistant flora [10]. However, this necessitates periodical assessment of the causative agents of road accident wound infection and their sensitivity profiles which is going to be of great use in the comprehensive treatment of such wound in Niger state.

2. MATERIALS AND METHODS

2.1 Study Area

The study was carried out in three General Hospitals in Minna, Bida, Kontagora and Suleja Local Government Areas of Niger state. Minna is the capital of Niger state with two Local Government Areas, Bosso and Chanchaga. Niger state is part of North Central Geopolitical Zone of Nigeria and is located between latitudes 8°20' N and 11°30' N and longitude 3°30' E and 7°20' E. This study area is made up of people with different ethnic groups living together.

2.2 Study Population

Random sampling technique was used to collect samples from accident victims. A total of 500 wound swab samples were collected from in-patients of 15 to 70 years of age with fresh road accidents injury brought to accident and emergency unit at three general hospital considered in the study.

2.3 Demographic Information

Socio-demographic data such as age, sex, location of accident and vehicle (s) involved in the accident using standard questionnaires and kept confidential during the research.

2.4 Sample Collection

Five hundred fresh wound swabs were collected from 409 patients within the space of 6 months. Specimens were collected by rubbing the lesion with a sterile swab stick. The swab sticks were introduced into an ice pack and then transported to the microbiology laboratory of Federal University of Technology Minna Niger state for analysis.

2.5 Sample Processing

The swab sticks were inoculated onto Nutrient Broth and incubated at 37°C for 24 hours. It was then sub cultured onto Nutrient Agar and Mac Conkey Agar by picking the organism from the Nutrient Broth culture with wire loop and were incubated for 24 hours at 37°C. The colonies were then sub cultured onto Nutrient Agar and incubated at 37°C for 48 hours to obtain pure isolates. The pure isolates were then characterized by Gram's staining and biochemical tests using Bergey's Manual of Determinative Bacteriology [11]. Suspected bacterial species were characterized and identified according to standard bacteriological methods as highlighted by Chessbrough, [12] and Barrow and Feltham [13].

2.6 Antibiotic Susceptibility Test

Nutrient broth was prepared and inoculated with bacterial isolate (5 ml of nutrient broth) using wire loop, and was incubated at 37°C for 3 hours. After incubation, the turbid culture was compared with McFarland standard, and was smeared onto prepared nutrient agar using sterile cotton swab [14]. The commercial antibiotic susceptibility discs were picked with sterile forceps and placed on the surface of the inoculated nutrient agar under aseptic condition. The antibiotic discs used for Gram negative sensitivity were: Tarivid, pefloxacin, ciprofloxacin, pefloxacin, chloramphenicol, amoxicillin, Augmentin, gentamycin, streptomycin, septrin and Gram-positive sensitivity as follows: Norfloxacin, streptomycin, levofloxacin, rifampicin, erythromycin, ampiclox, chloramphenicol, gentamycin, ciprofloxacin, amoxicillin. The plate containing the discs was incubated at 37°C for 24 hours. The zones of inhibition were recorded accordingly and those that were resistant were also recorded. This was done for all the isolates.

3. RESULTS

A total of 1000 wound swabs from 409 patients with road accident wounds over the period of 6 months were sampled. All were in-patients in different wards in the hospital (i.e. Accident and Emergency ward, Emergency paediatric unit, intensive care unit (ICU), Surgical and Medical wards, Obstetrics and Gynaecology ward and Paediatrics ward).

The percentage of positive swab cultures as against negative swab cultures was shown in Fig. 1. Of the 1000 swab samples taken from 409 patients, 267 were male and 142 were female, 405 (40.5%) samples showed growth for microorganisms while 595 (59.5%) showed no growth. Of the 405 microbial positive cultures, 347 (85.7%) were positive for bacteria, while 58 (14.3%) were positive for fungi as seen in Fig. 2.

The number of organisms isolated per wound swab cultured, 262 (80.6%) of the microbial positive cultures showed single organism per wound swab, 46 (14.2%), showed double organisms per wound swab while 17 (5.2%) showed a mixture of three or more organisms per wound swab as shown in Fig. 3.

Table 1 shows the frequency of occurrence of the organisms isolated from road accident wounds over the period of 6 months. *Staphylococcus aureus* has the highest number of organisms with (37.7%), and is followed by *Pseudomonas aeruginosa* (15.6%), *E. coli* (11.4%), *Streptococcus pyogenes* (9.4%), *Bacillus subtilis* (7.2%), *Klebsiella pneumoniae* (4.4%), *Aspergillus niger* (4.2%), *Candida albicans* (3.7%), *Aspergillus flavus* (2.7%), *Candida pseudotropicalis* (2.2%) and *Mucorpusillus* (1.5%).

Table 2 shows the Antibiotics sensitivity pattern exhibited by these microorganisms isolated from Road Accident Wound infections. Antibiotics sensitivity disks were used to carry out sensitivity test on each bacterial isolate. Gentamycin was sensitive to all bacteria isolated but resistant to *Klebsiella spp.* The zone of inhibition was measured in diameter (mm) as: R-Resistance: 13 mm or less, I-Intermediate: 14-16 mm and S-Sensitive: 17 mm or more according to the methods described by CLSI [14] and Reller et al. [15]. The antibiotics used were: ST – Septrin, CH – Chloranphenicol, CX – Ciprofloxacin, SP – Sparfloxacin AX – Amoxicillin, AU – Augmentin,

CN – Gentamycin, OF – Travid S – Streptomycin, PF–Pefloxacin, NB-Norfloxacn, LF-Levofloxacin, RD-Rifampicin, E-Erythromycin, AX-Ampiclox (Table 2).

Furthermore, Table 3 shows the antifungal susceptibility pattern of the fungal isolates. Most of the antifungal drugs tested showed larger zones of inhibition, for instance itraconazole (40.5 mm), ketoconazole (40.0 mm), miconazole (39.0 mm) and fluconazole (39.0 mm). All the fungi

isolated, showed no zones of inhibition that is resistant to the antifungal drug Griseofulvin.

4. DISCUSSION

This study was meant to explore the most common causes of wound infections associated with road traffics accident injury by examination of wound swabs. Of the 1000 wound swab samples taken from 409 (267 males and 142 females) patients with road traffic accident




Fig. 1. Percentage of positive swab culture as against negative




Fig. 2. Percentage of positive bacterial isolates in relation to positive fungal isolates

wounds, 595 (59.5%) showed no microbial growth and 405 (40.5%) showed microbial growth. This could be attributed to the proper handling of accident wounds by health professionals in these Hospitals or the fact that the level of bacterial contamination present was naturally cleared by the host [16,17]. The 21-to-40 age group was mostly involved in road traffic

accident according to this study. This could be due to the fact that this age group constitute the most economically productive age bracket. This was similar to the findings of Chang [18], who attributed the reason why age group 21 to 40 are overrepresented in road accident fatalities to their economic productivity as they carry out their daily activities.




Fig. 3. Distribution of microorganism isolated per wound swab examined

Table 1. Frequency of occurrence of bacterial and fungal isolates

Isolates	Number of isolates	Frequency of occurrence (%)
<i>Bacillus subtilis</i>	29	7.2
<i>Escherichia coli</i>	46	11.4
<i>Klebsiella pneumoniae</i>	18	4.4
<i>Pseudomonas aeruginosa</i>	63	15.6
<i>Staphylococcus aureus</i>	153	37.7
<i>Streptococcus pyogenes</i>	38	9.4
<i>Aspergillus niger</i>	17	4.2
<i>Aspergillus flavus</i>	11	2.7
<i>Candida albicans</i>	15	3.7
<i>Candida pseudotropicalis</i>	9	2.2
<i>Mucor pusillus</i>	6	1.5
Total	405	100

Table 2. Antibiotics susceptibility test on bacterial Isolates from road accident wounds

Bacteria	OF	PF	CX	SP	CH	AX	AU	CN	S	ST	NB	LF	RD	E	AX
<i>Bacillus subtilis</i>	R	I	S	R	R	R	R	R	S	R	R	S	R	R	R
<i>Escherichia coli</i>	R	R	S	R	S	S	R	S	S	R	S	S	R	S	R
<i>Klebsiella pneumoniae</i>	R	R	R	R	R	R	R	R	R	R	R	I	R	R	R
<i>Pseudomonas aeruginosa</i>	R	I	S	S	R	R	R	R	S	R	R	S	R	R	R
<i>Staphylococcus aureus</i>	I	S	R	R	I	I	R	I	S	R	I	S	R	R	R
<i>Streptococcus pyogenes</i>	S	S	S	R	R	R	R	S	S	S	S	S	R	S	R

R-Resistance: 13 mm or less, I-Intermediate: 14-16 mm and S- Sensitive: 17 mm or more

Table 3. Antifungal susceptibility pattern on fungal Isolates from road accident wounds

Fungi	Itraconazole (mm)	Ketoconazole (mm)	Miconazole (mm)	Fluconazole (mm)	Griseofulvin (mm)
<i>A. niger</i>	39.00	33.00	32.00	32.00	0.00
<i>A. flavus</i>	39.50	34.00	33.00	33.00	0.00
<i>C. albicans</i>	39.50	34.50	33.45	33.00	0.00
<i>C. pseudotropicalis</i>	37.50	32.60	31.50	31.00	0.00
<i>M. pusillus</i>	40.50	40.00	39.00	39.00	0.00

The most frequently isolated microorganisms were *Staphylococcus aureus* (37.7%), *Pseudomonas aeruginosa* (15.6%) and *Escherichia coli* (11.4%). This observation was in agreement with previous reports in Akwa Ibom and Cross River States, Nigeria [19,20,8]. General Hospital Minna showed the highest number of patients with swab specimen. This may be due to the fact that General Hospital Minna is in the capital and is with the largest number of bed space and serves as the referral centre to the other General Hospitals in the State. General Hospital Bida showed the least number of patients with swab specimens and this may be due to the fact that General Hospital Bida is in the same location with Federal Medical Centre Bida which serves as a referral centre for most accident cases in the area.

Staphylococcus aureus was the commonest isolated organism as it confirms the ubiquitous nature of *Staphylococcus aureus*, which is found in the human skin as a normal flora [21,22]. *Bacillus subtilis* though not an organism commonly found in wound, but could be found in air, soil and water and hence can easily infect wound [23]. *Candida albicans* is found in the vagina, but its presence in wound may be traced to immunocompromised state such as patients with diabetes mellitus, on prolonged steroid or prolonged use of antibiotics [24,25,26].

Klebsiella pneumoniae showed the highest resistance to all the antibiotics used, except for Levofloxacin. This is similar to the findings of Fair and Tor [27], who reported the multidrug resistant ability of *Klebsiella pneumoniae* owing to its thick polysaccharide capsule that acts as an antiphagocytic factor [28] and was the first species that *qnr* quinolone resistance genes were isolated from [29] All the fungal isolates from this study showed marked resistance against the antifungal drug Griseofulvin. Griseofulvin is known to act by binding to keratin tissue which is absent in wound [30]. Other antibiotics and antifungals used in this study showed moderate to high sensitivity. Finally, the antibiotics:

Levofloxacin, Ciprofloxacin, Streptomycin, Gentamycin and the antifungal: Itraconazole, Miconazole and Fluconazole are the most sensitive drugs as shown in this study, hence are recommended for use as empirical drug treatment pending the outcome of laboratory sensitivity result.

5. CONCLUSION

Wound infections due to road traffic accidents is increasing and is becoming a problem in all the locations and areas studied. Many organisms were isolated during the period of study. Most of the bacteria were resistant to the antibiotics used, for instance the bacteria *Klebsiella pneumoniae* was resistant to all the antibiotics used except for Levofloxacin.

Fungi isolated in this study were resistant to the fungicide Griseofulvin, but highly sensitive to the antifungal drugs Itraconazole, Ketoconazole, Miconazole and Fluconazole. Levofloxacin, Ciprofloxacin, Streptomycin and Gentamycin were the most effective antibiotics used in this study and hence are the best drugs of choice for the empirical treatment of injuries and resistance due to road traffic accidents wound infection.

CONSENT AND ETHICAL APPROVAL

Wound swab samples were obtained with full informed written consent of the accident victims and for subjects less than 18 years had their consent sought from their relatives or guardian. Clearance to conduct this research was sought from the ethics and research committee of the hospital management board, Minna. Also, absolute confidentiality and privacy was respected.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organisation. World report on road traffic injury prevention: Summary. World Health Organization, Geneva, Switzerland; 2004.
2. Odero W, Khayes M, Heda PM. Road traffic injuries in Kenya: Magnitude, causes and status of intervention. *Inju Contr Saf Prom.* 2003;10:53-61.
3. Porter SJ, Greavis IK. Major trauma. UK: Oxford University Press; 2010.
4. Palmer C. Major trauma and the injury severity score—where should we set the bar? Annual Proceedings of the Association for the Advancement of Automotive Medicine. 2007;51:13–29. [PMC 3217501, PMID 18184482]
5. Bowler PG, Duerden BI, Amstrong DC. Wound microbiology and associated approaches to wound management. *Clin Microb Rev.* 2001;14:244–269.
6. Dionigi R, Rovera F, Dionigi G, Imperator A, Ferrari A, Dionigi R. Risk factors in surgery. *J Chemoth.* 2001;13:6-11.
7. Iroha IR, Amadi ES, Orji JO, Esimone CO. In vitro evaluation of the activity of colloidal silver concentrate against *Pseudomonas aeruginosa* isolated from postoperative wound infection. *Sci Res Essay.* 2008;3(5): 209-211
8. Akinjogunla OJ, Adegoke AA, Mbotto CI, Chukwudebelu IC, Udokang IP. Bacteriology of automobile accident wounds infection. *Int J Med Med Sci.* 2009;1(2):23-27.
9. Federal Road Safety Commission: Nigerian traffic accidents (January to October, 2009). FRSC report 2009. Abuja: Federal Road Safety Commission; 2009.
10. Shittu AO, Kolawole DO, Oyedepo ER. A study of wound infections in two health institutions in Ile-Ife, Nigeria. *Afr J Bio Res.* 2002;5:97-110.
11. Buchanan KE, Gibons NE. Bergey's manual of determinative bacteriology. (8th ed.) Williams and Williams Company, Baltimore; 1974.
12. Cheesbrough M. Microscopical techniques used in microbiology. In district laboratory practice in tropical countries (Part 2). Third edition United Kingdom: Cambridge University Press; 2010.
13. Barrow GI and Feltham RKA. Bacterial characters and characterization. In: Cowan and Steel's manual for identification of medical bacteria (3rd ed.) Cambridge University Press. The Edinburgh Building, Cambridge CB2 2RU, UK; 2003.
14. Clinical and laboratory standards institute (CLSI). Performance standards for antimicrobial susceptibility testing; Twenty-third informational supplement. CLSI document M100-S23. CLSI, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA; 2013.
15. Reller LB, Weinstein M, Jorgensen JH, Ferraro MJ. Antimicrobial susceptibility testing: A review of general principles and contemporary practices. *Clin. Inf. Dis.* 2009;49(1):1749–1755. Available: <https://doi.org/10.1086/647952>
16. Landis SJ. Chronic wound infection and antimicrobial use. *Advanced Skin Wound Care.* 2003;21:531-540.
17. Stotts NA. Wound infection: Diagnosis and management. In: Bryant R, Nix D (Eds). *Acute and chronic wounds: Current management concepts.* 4th ed. St. Louis, MO: Elsevier Mosby; 2012.
18. Chang D. Comparison of crash fatalities by sex and age group. Traffic safety facts, National center for statistics and analysis, 1200 New Jersey Avenue SE., Washington, DC. 2008;1-6.
19. Anbumani N, Kalyan J and Mallike M. Epidemiology and microbiology of wound infections. *Ind J Prac Doc.* 2006;3(5):55–73.
20. Isibor JO, Oseni A, Eyaufe A, Osagie R, Turay A. Incidence of aerobic bacteria and *Candida albicans* in post-operative wound infections. *Afr J Micr Res.* 2008;2:288-291.
21. Tong SY, Davis JS, Eichenberger E, Holland TL, Fowler VG. *Staphylococcus aureus* infections: Epidemiology, pathophysiology, clinical manifestations, and management. *Clin. Microbiol. Rev.* 2015; 28(3):603-61.
22. Rasigade JP, Vandenesch F. *Staphylococcus aureus*: A pathogen with still unresolved issues. *Infect. Genet. Evol.* 2014;21:510-514.
23. Edokpayi J, Oruko R, Odiyo J. The role of leather microbes in human health [Online first], Intech Open; 2019. DOI: 10.5772/intechopen.81125 Available: <https://www.intechopen.com/online-first/the-role-of-leather-microbes-in-human-health>
24. Allison DL, Willems HME, Jayatilake JAMS, Bruno VM, Peters BM, Shirtliff ME. *Candida*-bacteria interactions: Their impact

- on human disease. *Microbiol Spectr.* 2016; 4:103–136.
25. Schlecht LM, Peters BM, Krom BP, et al. Systemic *Staphylococcus aureus* infection mediated by *Candida albicans* hyphal invasion of mucosal tissue. *Microbiology (Reading, Engl)*. 2015;161(1):168–181.
 26. Malani PN, McNeil SA, Bradley SF, Kauffman CA. *Candida albicans* sternal wound infections: A chronic and recurrent complication of median sternotomy. *Clin Infect Dis.* 2002;35(11):1316-20.
 27. Fair RJ, Tor Y. Antibiotics and bacterial resistance in the 21st century. *Perspect Medicin Chem.* 2014;6:25–64.
 28. Podschun R, Ullmann U. *Klebsiella spp.* as Nosocomial pathogens: Epidemiology, taxonomy, typing methods, and pathogenicity factors. *Clin Microbiol Rev.* 1998; 11(4):589–603.
 29. Martinez-Martinez L, Pascual A, Jacoby GA. Quinolone resistance from a transferable plasmid. *Lancet.* 1998;351(9105): 797–9.
 30. Sobue S, Sekiguchi K, Nabeshima T. Intracutaneous distributions of fluconazole, Itraconazole, and griseofulvin in guinea pigs and binding to human stratum corneum. *Amer Soci Micro.* 2004;48(1): 216-223.

© 2019 Chukwuka et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48687>