



Effects of Organic and Inorganic Fertilizer on the Early Growth Response of *Azelia africana*

Adeboyin Funmi Aderounmu^{1*}

¹*Federal College of Forestry, P.M.B. 5087, Jericho Hill Ibadan, Nigeria.*

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/AJOB/2019/v8i230060

Editor(s):

(1) Dr. Xing Li, Division of Biomedical Statistics and Informatics, Department of Health Sciences Research, Mayo Clinic College of Medicine and Science, USA.

Reviewers:

(1) Cristiane Ramos Vieira, University of Cuiabá, Brazil.
(2) Maria Lúcia Pereira da Silva, University of São Paulo, Brazil.

(3) Natchimuthu Karmegam, Government Arts College, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50881>

Original Research Article

Received 13 June 2019
Accepted 24 August 2019
Published 30 August 2019

ABSTRACT

An experiment was conducted in the nursery of Department of Forestry Technology at the Federal College of Forestry Ibadan, Oyo state, Nigeria to determine the effect of organic and inorganic fertilizers on early growth response of *Azelia africana*. Seedlings were collected from Forestry Research Institute of Nigeria, Ibadan, Oyo State and were transplanted into 2kg perforated polythene pots. The experiment was laid out in a completely randomized design with five treatments replicated three times. The treatments were: 20 t ha⁻¹ water hyacinth compost (T1), 20 t ha⁻¹ water hyacinth + poultry manure compost (T2), 20 t ha⁻¹ poultry manure (T3) and 50 kg ha⁻¹ NPK 20:10:10 (T4) and T5 - control (no fertilizer application). The compost was applied two (2) weeks before planting, while NPK 20:10:10 was applied two 2 weeks after planting. The experiment was monitored for eight (8) weeks after transplanting (WAT), while growth parameters were measured. The results of the study showed that application of fertilizers gave significant (p=0.05) increase in plant height (cm), stem diameter (mm), leaf production, and leaf area (cm²) of *A. africana*. Plant height ranged from 41.43 cm in the control to 47.96 cm in the pots where 20t ha⁻¹ water hyacinth compost was applied. Stem diameter also increased across treatments, while leaf production ranged from 9 in the control treatment to 14 in the pots with 20t ha⁻¹ poultry manure. These results suggested that the incorporation of organic and inorganic fertilizers increased

*Corresponding author: E-mail: afaderounmu@gmail.com;

productivity of *A. africana*. However, application of 20 t ha⁻¹ water hyacinth compost is recommended because it is available, affordable and environmentally friendly for the production of *A. africana* especially in the study area.

Keywords: Water hyacinth compost; *Azelia africana*; NPK 20:10:10.

1. INTRODUCTION

Forests and trees perform various functions in the ecosystem namely; aesthetics, provision of food and medicine, provision of shelter to wildlife and hygienic purpose [1]. It is universally accepted that forests and trees carry out a fundamental role in soil and water resources conservation [2,3]. As population density increases and land for food production expands due to agricultural activities and urbanization, natural forests became degraded [4]. The degradation has led to the disappearance of most species including agro forest trees and causing difficulty in growing some seedlings including *Azelia africana* [5,6,7,8]. *A. africana* is a leguminous tree found in the humid and dry forest savannah borders or semi-deciduous forest [6]. It is used for soil conservation and improvement [9]. *A. africana* is a timber species with high forage, economic and pharmacological values. Its leaves are harvested for grazing during the dry season. The high demand for *A. africana* leaves, seeds, roots and barks for various uses has resulted in corresponding increase in the exploitation at such a rate that sustainability of this natural resource cannot be guaranteed [10,11]. Documented reports on the cultivation and seedling growth of this multipurpose tree known commonly as African mahogany are scarce [12,13,7,14]. If the benefits derivable from *A. africana* must continue especially for the future generations, there is the need to stimulate farmers' interest in the cultivation of *A. africana* thereby helping to reduce poverty, helping in conservation role as well as to boost the source of revenue for the government. In the same vein, information on the domestication of the plant seeds and seedlings are scanty, mainly due to degradation caused by human influences. Consequently, the need to acquaint farmers with the most successful soil or manures that could enhance the growth of *A. africana* seedlings cannot be overemphasized. This study was aimed at investigating the effect of organic and inorganic fertilizers on the seedling establishment of *A. africana* with a view to recommend the best fertilizer to *A. africana* growers especially at the nursery stage and to multiply this multi-purpose species.

2. MATERIALS AND METHODS

The experiment was carried out in 2018 at nursery site of the Department of Forestry Technology, Federal College of Forestry Jericho Ibadan (Latitude 07°27' N and longitude 03°53' E), Ibadan, Nigeria (FRIN Meteorological Station, 2018). The annual rainfall is 1250 mm with a bimodal pattern and has a minimum temperature of 21.9°C and maximum temperature of 35.5°C. The experiment was laid out in a Complete Randomized Design with five treatments replicated three times. The treatments were: 20 t/ha water hyacinth compost (T1); 20 t ha⁻¹ water hyacinth + poultry manure compost (T2); 20 t/ha poultry manure (T3); 50 kg/ha N:P:K 20:10:10 (T4); and control (no fertilizer application) (T5). The compost was applied two (2) weeks before planting. Eight week old seedlings of *A. Africana* were collected from Forestry Research Institute of Nigeria (FRIN), Ibadan, Oyo State, and potted into 2kg polythene pots filled with different media treatments, watered and allowed to stabilize for two weeks before the commencement of growth assessment. Plant height was measured with a meter rule at the distance from soil level to terminal bud. Leaf production was determined by counting. Stem diameter was measured with venier caliper. Data collected were analysed statistically using Genstat Software Package and were subjected to analysis of variance. Means were separated using Duncan's multiple range test (DMRT) at 5% level of significance.

3. RESULTS

The physical and chemical properties of the soil (0–15 cm depth) at the experimental site before planting are as presented in Table 1. The soil is sandy loam, slightly acidic, high bulk density (1.62 g cm⁻³) and has been classified as an Alfisol [15] with its distinctive characteristics. The data in Table 1 further confirms this assertion and also reveals that the soils are moderate in Zn, low in K (0.11 cmol_c kg⁻¹), organic carbon (9.0 g kg⁻¹), total nitrogen (1.0 g kg⁻¹) and P (6.0 mg kg⁻¹). Saturated hydraulic conductivity value of 12.4 cm hr⁻¹ indicated a well-drained soil.

Table 1. Pre-planting soil physical and chemical properties of the experimental site

Soil parameters	Content in soil
pH (H ₂ O)	5.8
Organic carbon (g kg ⁻¹)	9.0
Total nitrogen (g kg ⁻¹)	1.0
Available phosphorus(mg kg ⁻¹)	6.0
Exchangeable cations (cmol_c kg⁻¹)	
Ca	1.32
Mg	0.25
K	0.11
Na	0.31
Extractable micronutrient (mg kg⁻¹)	
Mn	302.0
Fe	265.0
Cu	3.39
Zn	1.2
Exchangeable Acidity (cmol _c kg ⁻¹)	0.40
Particle size distribution (g kg⁻¹)	
Sand	838
Silt	54
Clay	108
Textural class	Sandy loam
Bulk density (g cm ⁻¹)	1.62
Saturated hydraulic conductivity (cm hr ⁻¹)	12.4

Table 2. Chemical properties of the combinations of compost used

Parameter	Poultry manure	Water hyacinth compost	Water hyacinth + Poultry manure compost
pH (H ₂ O)	6.8	5.89	5.76
Organic carbon (%)	22.94	31.92	32.11
Total Nitrogen (%)	3.30	2.6	1.73
C:N	6.8	12.52	43.99
P (%)	0.83	1.24	1.55
K (%)	1.80	0.47	0.80
Ca (%)	2.56	1.6	1.80
Mg (%)	1.58	4.30	0.92

The chemical composition of water hyacinth compost, poultry manure and water hyacinth + poultry manure compost used is as presented in Table 2. The pHs of all the organic fertilizers were all acidic with the water hyacinth + poultry manure compost having the most acidic pH of 5.76. Organic carbon, C/N and P were highest in the mixture of water hyacinth + poultry manure compost. Poultry manure was highest in N, K and Ca content.

3.1 Effect of Organic and Inorganic Fertilizers on Number of Leaves of *Azelia africana* Seedling

Fig. 1 showed the growth pattern of leaf production of *A. africana* seedlings as influenced

by the application of organic and inorganic fertilizers. Results revealed that mean leaf production of *A. africana* increased across the study period for T3 (20 t ha⁻¹ poultry manure) which recorded the highest mean number of leaves across the weeks and except in the last week while T5 (top soil only) had the least mean number of leaves all through the duration of the experiment.

The mean effects of organic and inorganic fertilizers on the number of leaves of *A. africana* as presented in Fig. 2. Leaf production varied across treatments and differs significantly (p=0.05) throughout the experiment. Leaf production ranged from 9 in the control to 14 in the pots where 50 kg ha⁻¹ NPK 20:10:10 was

applied. The highest mean leaf production of *A. africana* was observed in T3 with 14 and it was closely followed by T1, T2, and T4 with 13 and the least was found in T5 (control) with 9.

3.2 Effect of Organic and Inorganic Fertilizers on Plant Height of *Azelia africana* Seedling

Results in Fig. 3 showed the growth pattern of plant height of *A. africana* seedlings. The control treatment followed similar trend with the growth pattern of number of leaves and performed poorly in comparison to others in terms of the

number of leaves produced. The highest mean height of *A. africana* was observed in T1 (20t/ha water hyacinth compost) with T2 (20 t ha⁻¹ water hyacinth + poultry manure compost) closely followed and T5 (control) performed least. The effect of organic and inorganic fertilizers on the plant height of *A. africana* is as presented in Fig. 4. Plant height increased across treatments but did not differ significantly (p=0.05) throughout the experiment. 20t ha⁻¹water hyacinth compost (47.96 cm) had highest plant height followed by 20 t ha⁻¹ water hyacinth + poultry manure compost (47.89 cm) and least by control pot (41.43 cm).




Fig. 1. Growth pattern of number of leaves of *Azelia africana* seedlings as influenced by fertilizers

T1 (20 t ha⁻¹ water hyacinth); T2 (20 t ha⁻¹ water hyacinth + poultry manure compost); T3 (20 t ha⁻¹ poultry manure); T4 (50 kg ha⁻¹ NPK 20:10:10); T5 (control-no fertilizer application)




Fig. 2. Effects of fertilizers on number of leaves of *Azelia africana* seedlings

T1 (20 t ha⁻¹ water hyacinth); T2 (20 t ha⁻¹ water hyacinth + poultry manure compost); T3 (20 t ha⁻¹ poultry manure); T4 (50 kg ha⁻¹ NPK 20:10:10); T5 (control-no fertilizer application)




Fig. 3. Growth pattern of height of *Azelia africana* seedlings as influenced by fertilizers
 Legend: T1 ($20t\ ha^{-1}$ water hyacinth); T2 ($20t\ ha^{-1}$ water hyacinth + poultry manure compost); T3 ($20t\ ha^{-1}$ poultry manure); T4 ($50\ kg\ ha^{-1}$ NPK 20:10:10); T5 (control-no fertilizer application)




Fig. 4. Effects of fertilizers on plant height of *Azelia africana* seedlings
 T1 ($20t\ ha^{-1}$ water hyacinth); T2 ($20t\ ha^{-1}$ water hyacinth + poultry manure compost); T3 ($20t\ ha^{-1}$ poultry manure); T4 ($50\ kg\ ha^{-1}$ NPK 20:10:10); T5 (control-no fertilizer application)




Fig. 5. Growth pattern of stem diameter of *Azelia africana* seedlings as influenced by fertilizers
 T1 ($20t\ ha^{-1}$ water hyacinth); T2 ($20t\ ha^{-1}$ water hyacinth + poultry manure compost); T3 ($20t\ ha^{-1}$ poultry manure); T4 ($50\ kg\ ha^{-1}$ NPK 20:10:10); T5 (control-no fertilizer application)




Fig. 6. Effects of fertilizers on stem diameter of *Afzelia africana* seedlings
 T1 (20 t ha^{-1} water hyacinth); T2 (20 t ha^{-1} water hyacinth + poultry manure compost); T3 (20 t ha^{-1} poultry manure); T4 (50 kg ha^{-1} NPK 20:10:10); T5 (control-no fertilizer application)

3.3 Effect of Organic and Inorganic Fertilizers on Stem Diameter of *Afzelia africana* Seedling

Results in Fig. 5 showed the growth pattern of stem diameter of *A. africana* seedlings, the control experiment performed well at the beginning but poorly at the end of the eighth week, with T1 (20 t ha^{-1} water hyacinth) performing best.

The effect of organic and inorganic fertilizers on the stem diameter of *A. africana* is presented in Fig. 6. Stem diameter increased across treatments but did not differ significantly ($p=0.05$) throughout the experiment. 20 t ha^{-1} water hyacinth compost (0.81 mm) had highest stem diameter followed by 20 t ha^{-1} water hyacinth + poultry manure compost (0.76 mm) and least by control plot (0.62 mm).

4. DISCUSSION

The major factor affecting plant growth in the tropics is the nutrient deficiency in tropical soils resulting from degraded farmland. The low levels of N, P, and OC observed in the experimental soil indicated that the soil had a low fertility status. The value obtained for N, P, K and OC is below the critical range [16,17], thus indicating poor soil fertility not suitable without the addition of external input for planting *A. africana*. The chemical composition of water hyacinth compost, poultry manure and water hyacinth + poultry manure compost used in the experiment was relatively high in major elements (N, P, K, Ca and

Mg). The OC content of all the organic fertilizers were less than the values obtained for the composts of *Azadirachta indica*, *Albizia lebbek* and *Khaya senegalensis* by Daldoum and Hammad [18].

The application of the various fertilizers increased the growth of *A. africana* this agreed with the findings of Uddin et al. [19] where organic fertilizers enhanced the seedling growth of some leguminous forestry species. This could result from the nutritional benefits of organic and inorganic fertilizers which include improvement of soil fertility. The result obtained from the plant height showed that water hyacinth compost significantly induced the shoot growth, leaf production and stem diameter of *A. africana* seedlings. This is in support with [20,21,22] that reported that plant height and number of leaves of plants treated with water hyacinth compost had been used and shows better performance than control. Lata [23] that experimented with water hyacinth manure on *Coriandrum sativum* revealed positive response with increase in manure rates. Supported by study done by Osoro [24] and Aboul-Enein et al. [25] who advocated that water hyacinth has good N, P, K absorbing capacity from water and thus can be used as a good source of compost material to serve as fertilizer in soil with poor amount of N, P, K and C values. Water hyacinth which used to be tagged as waste and nuisance to aquatic environment can be converted to compost for fertilizing plants at the nursery stage in order to improve early growth. This might replace use of expensive, scarce and environmentally

hazardous inorganic fertilizers in forest nursery work.

5. CONCLUSION

Organic and inorganic fertilizer had effect on the growth of *A. africana*. It could be observed from the result obtained that there were increases in plant height, stem diameter and number of leaves. However, there was poor performance throughout the assessment period in control treatment when compared to other treatments in terms of the number of leaves produced per plant. Seedlings in 50 kg ha⁻¹ NPK 20:10:10 (T4) presented best at week eight though it was poor at the early stage of the experiment. Seedlings with 20 t ha⁻¹ poultry manure (T3) performed well from the beginning of the experiment up to the penultimate week to the end of the experiment. Application of 20 t ha⁻¹ water hyacinth compost gave the highest plant height. Stem diameter also increased appreciably across treatments. Therefore, it can be concluded and recommended that 20t ha⁻¹ water hyacinth compost can be used by farmers to increase the growth of *A. africana*.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Agbogidi OM, Eshegbeyi OF. Forestry development for a safe environment. In: Onyekwelu JC, Adekunle VAJ, Oke DO. (ed.). Proceedings of the First National Conference of the forests and forest products society held at the Federal University of Technology, Akure, Ondo State. 2008; 95-98.
2. Broadhead J, Leslie R. Coastal protection in the aftermath of the Indian Ocean tsunami: What role for forests and trees? In: Proceedings of an FAO Regional Technical Workshop, Khao Lak, Thailand. FAO, Thailand; 2007.
3. Hamilton LS. Forests and water – A thematic study prepared in the framework of the Global Forest Resources Assessment 2005. Forestry Paper #155. Rome, Italy. 2008;78.
4. Salim E, Ullsten O. Our forest: Our future. A paper of the world commission on forest and sustainable development, Cambridge Press, U.K; 1999.
5. Nwoboshi LC. Teak (*Tectona grandis* Linn. F.) ten year after thinning. Nig. J. Forestry. 1985;15:76-79.
6. Keay RWJ. Tees of Nigeria: A revised version of Nigeria trees. 1989;1-2.
7. Etukudo I. Forests: Our divine treasure. Dorand Publishers, Uyo; 2000.
8. Ezenwaka J, Ekeke BA, Oyebade BA, Anyadiegwe AO. Identification and evolution of some threatened indigenous medical plants of the tropical rainforest of Eastern Nigeria. J. Conserv. F.P.L.G. Ibadan. 2004;2(122):131-141.
9. Agbogidi OM, Onomeregbor VA. Enhancing germination in seeds of *Azalia africana* (Sm. Expers.). In: Olufajo, OO, Omokore DF, Akpa GN, Sanni SA (ed.). Proceedings of the 41st Annual Conference of the Agricultural Society of Nigeria held in Zaria, Kaduna State. 2007;260-263.
10. Mtambalika K, Munthali C, Gondwe D, Missanjo E. Effect of seed size of *Azalia quanzensis* on germination and seedling growth. International Journal for Forestry Research. 2014;5.
11. Palgreave KC. Trees of Southern Africa. (3rd Ed). Revised by Coates MC. Struik Publishers, Cape Town; 2002.
12. Okeke RE. Comparative anatomy of trees in Nigeria: *Azalia africana* Species; 1996.
13. Burkill HM. The useful plants of West Africa. *Calesal piniaceae* Royal botanic Gardens. Longman publishing U.K;1999.
14. Agbogidi OM, Onomeregbor VA, Awwevughwar OI. Eco Physiological effects of crude oil pollution *Azalia africana* (Sm. Ex pers.) In: Ezike KNN, Osakwe II, Ekwu LG, Utobo E, Mba CN. (eds.). Proceedings of the 42nd Annual Conference of the Agricultural Society of Nigeria held at the Ebonyi State University, Abakakili, Ebonyi State. 2008;365-369.
15. Smyth AJ, Montgomery RF. Soil and landuse in central western Nigeria. Government of Western Nigeria, Ibadan, Nigeria; 1962.
16. Adeoye GO, Agboola AA. Critical level for soil pH, available P, k, Zn and maize ear leaf content for P, Cu and Mn in sedimentary soils of Southwestern Nigeria. 1985;6:65-71.
17. Akinrinde EA, Obigbesan JA, Onanuga OA, Extractable micronutrient in relation to physiochemical properties of soil in four ecological zones in Nigeria. Global Journal of Pure and Applied Science. 2005;2(11): 197-200.

18. Daldoum DMA, Hammad GH. Performance of *Acacia senegal* (L.) wild seedlings growth under some tree manures and NPK fertilizers in nursery Site. Journal of Forest and Environmental Science. 2015;31(4):303-311.
19. Uddin MB, Mukul SA, Arfin Khan MAS, Hossain MK. Effects of organic manure on seedlings growth and nodulation capabilities of five popular leguminous agroforestry tree components of Bangladesh. Journal of Forestry Science. 2014;28(4):212–219.
20. Razaq M, Zhang P, Sheng H. Influence of nitrogen and phosphorus on the growth and root morphology of Acer mono. PLoS One. 2017;12(2):8.
21. Talkah A. Effect of organic fertilizer water hyacinth on the growth and production plant taro (*Colocasia esculenta* L.). Journal of Environment and Earth Sciences. [Internet]. 2015;5(22):5. Available:<http://www.iiste.org/Journals/index.php/JESS/article/view/2015;27595/28309>
22. Cuesta B, Villar-Salvador P, Puertolas JF, Jacobs DM, Rey Benayas J. Why do large, nitrogen rich seedlings better resist stressful transplanting conditions? A physiological analysis in two functionally contrasting Mediterranean forest species. Forest Ecology and Management [Internet]. 2010;260:71-78. Available:<http://80.24.165.149/webproduccion/PDFs/10ART10.PDF>
23. Lata N. The impact of water hyacinth manure on growth attributes and yields in Coriandrum. IOSR Journal of Environmental Science, Toxicology and Food Technology. 2013;5(3):04-07.
24. Osoro N. Effects of water hyacinth (*Eichhornia crassipes* [Mart.] Solms) compost on growth and Yield Parameters of Maize (*Zea mays*). British Journal of Applied Science & Technology. 2014; 4(4):617-633.
25. Aboul-Enein A, Al-Abd A, Shalaby E, Abul-Ela F, Nasr-Allah A, Mahmoud A. *Eichhornia crassipes* (Mart) solms, Plant Signaling & Behavior. 2011;6(6):834-836.

© 2019 Aderounmu; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50881>