



Assessment of Soil Properties under Long Term Integrated Nutrient Management Using STCR-based Targeted Yield Equations under Rice-wheat Cropping System

Pragya Kurmi ^{a*}, Hitendra K. Rai ^a, Mahendra Anjna ^b, Nazmeen Khanam ^c,
Bharti Parmar ^d and Rupesh Yadav ^c

^a Department of Soil Science and Agricultural Chemistry, Jawaharlal Nehru Krishi Vishwavidhyalay, Jabalpur, Madhya Pradesh-482004, India.

^b Department of Agronomy, Jawaharlal Nehru Krishi Vishwavidhyalay, Jabalpur, Madhya Pradesh, India.

^c Department of Soil Science and Agricultural Chemistry, Rajmata Vijayaraje Scindia Krishi Vishwavidhyalay, Gwalior, Madhya Pradesh-474002, India.

^d Department of Agronomy, Rajmata Vijayaraje Scindia Krishi Vishwavidhyalay, Gwalior, Madhya Pradesh-474002, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i232462

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92380>

Original Research Article

Received 27 July 2022
Accepted 30 September 2022
Published 11 October 2022

ABSTRACT

The experiment was carried out to assess the effect of organic and chemical fertilizers on soil chemical properties under a nine year long experiment of AICRP on STCR initiated in *Kharif* 2008. The field experiment was laid out at research farm of JNKVV, Jabalpur after *Rabi* season of 2018-19. The study was framed in split plot design with four main treatments of nutrient management viz., T₁-control; T₂-GRD; T₃- Soil test based NPK application for yield of 6.0 t ha⁻¹ for each crop, and T₄- Soil test based NPK application for yield of 6.0 t ha⁻¹ for each crop + 10 t ha⁻¹ yr⁻¹ FYM and four soil depths as sub treatment from D₁- 0-15 cm; D₂- 15-30 cm; D₃- 30-45 cm and D₄ – 45-60 cm replicated three times. The results shows significant influence of INM on soil organic carbon (SOC),

*Corresponding author: E-mail: pragyakurmi14@gmail.com;

cation exchange capacity (CEC) and CaCO₃ content of soil. After nine years of chemical fertilizer applications on general recommended dose bases and soil test bases in combination with FYM reported no profound impact on soil pH, electrical conductivity.

Keywords: Chemical properties; organic carbon; electrical conductivity; CaCO₃; cation exchange capacity.

1. INTRODUCTION

“The rice (*Oryza sativa* L.) -wheat (*Triticum aestivum* L.) cropping system, as a source of the world's basic food, is essential for maintaining global food security” [1,2,3]. “The RWCS is technologically most advanced system in existence that has experienced extensive evolution. For more than 85 percent of the overall RWCS in South Asia, this system is among the most substantial cereal crops utilized in the Indo-Gangetic Plains” (IGP; Banjara *et al.*, [4]). This system depletes the soil nutrients heavily [5]. With emerging environmental footprints, there are stoking fears about yield stagnation or contraction in the R-W system [6,7]. In order to safeguard agricultural productivity and nutrition security in India, the quality of the soil must be restored and strengthened [8].

Through agricultural intensification, India became able to produce enough food, feed, bioenergy, and fibre to satisfy the requirements of its burgeoning population. The augment of synthetic agrochemicals and nutrient responsive high yielding crop varieties made us self-sufficient in the sector of food production. But the boost in agricultural production system has slowed down or even declines over last few years due to heavy removal of native nutrients which lead to diminish the soil health [9,10].

The real hurdle for the nation in the upcoming years will be how to produce enough food and fibre in along with high-quality produce, which can only be accomplished by boosting the productivity of various crops through a collaborative approach of adopting organic and chemical fertilizers. INM improves soil quality and sustainability while increasing crop yields by 8–150% when compared to conventional farming methods [11]. It also boosts farmers' economic returns [12] and the efficiency with which they use water and nutrients [13]. The primary factor contributing to the ongoing loss of soil fertility is changes in fundamental soil properties such as pH, cation exchange capacity (CEC), and soil organic carbon (SOC) concentration under

various fertiliser management schemes. To address these challenges, the recommendation of fertiliser based on soil test results and crop response offers a hint for prudent and balanced nutrient application to maintain production [14,15]. Due to their complementary effects, combining organic and inorganic nutrient aids in reaching the intended result and improving soil health (Antil *et al.*, 2011).

2. MATERIALS AND METHODS

2.1 Site and Soil

The present field experiment on rice-wheat cropping system was initiated in monsoon 2008 at research farm of JNKVV, Jabalpur under the network of All India Coordinated Research Project (AICRP) on Soil Test Crop Response (STCR). The experimental field was located at 23°10' N latitude and 79°57' E longitude and at an elevation of about 393.0 meter above mean sea level. The climate of the experimental site is sub-tropical with hot dry summers and cool winters with an average annual rainfall, average humidity and average evaporation are 1274 mm, 73 percent and 3.93 mm/day, respectively. The experimental area represents to *Kymore* Plateau and Satpura hills agro-climatic zone of Madhya Pradesh belongs to *Kheri* series. The soil is vertisol characterized as medium deep black soil. Based on the analytical report the soils are neutral in reaction, medium salt concentration, organic carbon, available phosphorus and potassium and low in available nitrogen. Clay mineralogy is dominated by fine montmorillonitic minerals. Taxonomically it belongs to of hyperthermic family of *Typic Haplustert*. The initial soil samples from surface (0-15 cm) and sub-surface (15-30 cm) were analysed for mechanical composition, pH, electrical conductivity (EC), cation exchange capacity (CEC), bulk density (BD), organic carbon (OC) and available nitrogen, phosphorous and potassium following standard procedure. Table 1 shows the physio-chemical characteristics of the initial soil under investigation.

Table 1. Physicochemical properties of the experimental soil before commencing the study

Soil characteristics	Soil depth		Method followed
	0-15 cm	15-30 cm	
Soil pH (1:2.5 at 25°C)	7.48	7.45	Glass electrode pH meter [16]
Electrical Conductivity (EC) (dSm ⁻¹ at 25°C)	0.285	0.267	Electrical conductivity meter [16]
Organic Carbon (g kg ⁻¹)	5.37	5.28	Potassium dichromate rapid titration method [20]
Available Nitrogen (Kg ha ⁻¹)	211.66	164.69	Alkaline potassium permanganate method (Subbiah and Asija, 1954)
Available Phosphorous (Kg ha ⁻¹)	22.95	14.97	Soil extracted with 0.5 M NaHCO ₃ and colour development by ascorbic acid [18]
Available Potassium (Kg ha ⁻¹)	307.53	282.76	Neutral normal ammonium acetate method by using flame photometer [16]

2.2 Experimental Design and Treatments

The field experiment on rice-wheat cropping system was designed under AICRP on STCR with three nutrient management practices along with a control in a split plot design with three replications. The treatments selected for this study consisted of (i) unfertilized control (control); (ii) General Recommended Dose (GRD) of NPK (120 N₂O: 60 P₂O₅: 40 K₂O ha⁻¹); (iii) Soil Test Based (STB) NPK + Targeted Yield (TY) of 6.0 t ha⁻¹; (iv) STB NPK + TY of 6.0 t ha⁻¹ + FYM @10.0 t ha⁻¹ yr⁻¹ (NPK+ FYM).

2.3 Soil Sampling and Analysis

After completion of nine cropping cycles of rice-wheat, soil samples from 0-15, 15-30, 30-45 and 45-60 cm soil depth were collected. In each plot, soil was collected from three points randomly and mixed into one sample. The samples were air-dried in shade, ground to pass through a 2 mm sieve and used for the estimation of soil chemical properties. The pH was measured in 1:2.5 soil: water suspension with a glass electrode. "The electrical conductivity was measured in supernatant liquid of soil: water (1:2.5) suspension with the help of conductivity meter as described" by Jackson [16]. The CaCO₃ content was determined following the rapid titration method [17] with 0.5N H₂ SO₄ in the presence of bromothymol blue and bromocresol green. The organic carbon in soil samples were determined using K₂Cr₂O₇ as oxidizing agent (1N) and back titrating with 0.5 N FAS method as suggested by Walkley and Black, 1934. The available major nutrients viz., nitrogen, phosphorous and potassium were estimated using the methods suggested by Subbiah and

Asija (1954); Watanabe and Olsen's [18] and Jackson [16] respectively.

2.4 Statistical Analysis

For statistical analysis of data, ANOVA for split plot design on Microsoft Excel were prepared. Analysis of variance (ANOVA) was drawn as described by Gomez and Gomez [19]. The differences of treatment means were tested by 'F' test of significance on the bases of null hypothesis at 5% level of significance.

3. RESULTS AND DISCUSSION

3.1 Soil pH

Integrated nutrient management practices and soil depth had no significant effect on soil pH (Table 2). The pH of soil under different management practices ranged from 7.47 to 7.57. Lower value of soil pH was recorded under treatments of soil test based NPK for targeted yield of 6 t ha⁻¹ and soil test based NPK+FYM@10.0 t ha⁻¹ yr⁻¹ for targeted yield of 6 t ha⁻¹ as compared to treatments with general recommended dose of NPK and control. Lower pH value of 7.53 at 0-15 cm was recorded. Soil pH followed the increasing trend with increasing depth. The release of carbon dioxide (CO₂) and organic acids into the soil during the breakdown of the manure may have contributed to the reduction in soil pH in the FYM treatments. As a result of FYM's breakdown, aliphatic and aromatic hydroxyl acids may also be produced, which may complicate free and exchangeable aluminium ions and lower pH levels [21,22,23,24].

3.2 Electrical Conductivity

“In different nutrient management systems, the electric conductivity as a measure of soluble salts or salinity changed from 0.25 to 0.28 dS m⁻¹ with highest value in general recommended dose of NPK and lowest under soil test based NPK+FYM@10.0 t ha⁻¹ yr⁻¹ for targeted yield of 6 t ha⁻¹ (Table 2). However, no significant differences in EC were observed among the treatments and soil depths. At depth level EC ranged between 0.26 to 0.28 dS m⁻¹. The interaction effect of both nutrient management and soil depth were also found not significant. The results under long-term fertilizer experiments also showed no appreciable change in salinity. The conductivity also followed the same trend as soil pH. The integration of FYM with inorganic fertilizers was found more effective to decrease the EC of soil” [25,26,27,28].

3.3 Soil Organic Carbon

It is evident that continuous application of FYM in combination with NPK resulted in considerable changes of SOC than that of unfertilized control as well as NPK treated plots on the bases of general recommended dose and soil test base (Table 2). In the present study, the plots that received FYM+NPK (6.62 g kg⁻¹) and soil test based NPK (6.10g kg⁻¹) had significantly higher build-up in SOC over general recommended dose of NPK (5.92g kg⁻¹) and unfertilized control (4.83g kg⁻¹).

At depth level 0-15 cm soil depth reported maximum retention of SOC (6.74 g kg⁻¹) followed by 15-30 cm (6.07 g kg⁻¹), 30-45 (5.54 g kg⁻¹) cm and 45-60 cm (5.11 g kg⁻¹). Integrated nutrient management practice either maintains or enhances the organic carbon content in soil [29,30,31] (Alaamer et al., 2020). Application of 100% NPK + FYM @ 10t ha⁻¹ increased the organic carbon content [21,22,32,33,34,35,36].

3.4 Cation Exchange Capacity

The cation exchange capacity of the soil in different treatments ranged from 34.3 C mol (P⁺) kg⁻¹ to 40.0 C mol (P⁺) kg⁻¹. Treatment received FYM+NPK reported maximum 40.0 C mol (P⁺) kg⁻¹ cation exchange capacity followed by 38.9 C mol (P⁺) kg⁻¹ in soil test based NPK, 37.4 C mol (P⁺) kg⁻¹ in general recommended dose of NPK and 34.3 C mol (P⁺) kg⁻¹ unfertilized control. Data also indicated that CEC of soil significantly decreased with increasing soil depth and ranged from 40.7 to 34.5 C mol (P⁺) kg⁻¹ with highest value at 0-15 cm depth which was statistically at par with that found at 15-30 cm but significantly higher than those obtained at 30-45 cm and 45-60 cm soil depths. Cation exchange capacity of the soil increased significantly with the application of organic manures along with recommended dose of fertilizers [37,38,29,39]. However, Hati et al. [40], Sarkar et al. [41] and Brar et al., [42] reported higher CEC of soil in 150% NPK treatment over 100% NPK+FYM treatment.

Table 2. Effect of INM and soil depth on pH, EC and organic carbon

Treatment	pH	EC (dS m ⁻¹)	SOC (g kg ⁻¹)
Nutrient management			
T ₁	7.57	0.27	4.83
T ₂	7.54	0.28	5.92
T ₃ T ₃	7.53	0.26	6.10
T ₄	7.47	0.25	6.62
SEm ±	0.049	0.014	0.053
CD (p=0.05)	NS	NS	0.155
Soil depth			
D ₁	7.53	0.26	6.74
D ₂	7.55	0.27	6.07
D ₃	7.54	0.27	5.54
D ₄	7.56	0.28	5.11
SEm ±	0.028	0.010	0.119
CD (p=0.05)	NS	NS	0.347
M x S			
SEm ±	0.063	0.013	0.308
CD (p=0.05)	NS	NS	NS

Table 3. Effect of INM and soil depth on CEC and CaCO₃ of soil

Treatment	CEC	CaCO₃
Nutrient management	[C mol (P⁺) kg⁻¹]	(g kg⁻¹)
T ₁	34.3	56.3
T ₂	37.4	62.5
T ₃ T ₃	38.9	60.6
T ₄	40.0	62.4
SEm ±	0.61	0.71
CD (p=0.05)	1.79	2.09
Soil depth, cm		
D ₁	40.7	53.5
D ₂	38.7	59.1
D ₃	36.6	63.8
D ₄	34.5	65.3
SEm ±	0.76	0.43
CD (p=0.05)	2.13	1.25
M x S		
SEm ±	1.17	0.79
CD (p=0.05)	NS	NS

3.5 Calcium Carbonate Content

Nutrient management practices and soil depths significantly affected the CaCO₃ content of the soil (Table 3). CaCO₃ content in soil under control (56.3 g kg⁻¹) was significantly lower than those obtained in other treatments. The CaCO₃ content under general recommended dose of NPK (62.5 g kg⁻¹), soil test based NPK (60.6 g kg⁻¹) and treatment received NPK+FYM (62.4 g kg⁻¹) reported statistically on par. Data further revealed that CaCO₃ content in soil significantly increased from 53.5 to 65.3 g kg⁻¹ with increasing soil depth from 0-60 cm with highest (65.3 g kg⁻¹) value at 45-60 cm and lowest (53.5 g kg⁻¹) at 0-15 cm depth. The downward movement of Ca and its subsequent precipitation as carbonate or decomposition of CaCO₃ increases its content in the pedons with soil depth [43,44,45]. Dubey et al. [46] reported that application 100% NPK with FYM significantly increased the calcium carbonate content in sub-surface soil.

4. CONCLUSION

The current study showed that a targeted yield approach based on STCR boosted the utilization of integrated FYM and NPK fertiliser. After 9-year of application of various fertilizer treatments, it was observed that SOC, cation exchange capacity and CaCO₃ content of soil showed significant differences among the treatments, while soil pH and electrical conductivity were not

found significantly different. The addition of FYM along with NPK fertiliser elevated soil organic carbon levels compared to NPK alone treated plots, which may have contributed to the better soil properties in FYM treated plots against NPK treated plots.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Timsina J, Connor DJ. Productivity and management of rice-wheat cropping systems: Issues and challenges. *Field Crops Research*. 2001;69:93–132.
2. Lalik R, Sharma S, Idris M, Singh AK, Singh SS, Bhatt BP, Saharawat Y, Humphreys E, Ladha JK. Integration of conservation agriculture with best management practices for improving system performance of the rice-wheat rotation in the eastern Indo-Gangetic Plains of India. *Agriculture Ecosystem and Environment*. 2014;195: 68–82.
3. Banjara TR, Bohra, JS, Kumar S, Ram A, Pal V. Diversification of rice-wheat cropping system improves growth, productivity and energetics of rice in the

- Indo-Gangetic Plains of India. Agricultural Research. 2021a;10:1–10.
4. Banjara TR, Bohra JS, Kumar S, Singh T, Shori A, Prajapat K. Sustainable alternative crop rotations to the irrigated rice-wheat cropping system of Indo-Gangetic Plains of India. Archives of Agronomy and Soil Science. 2021b;67: 1–18.
 5. Bhatt R, Kukal SS, Busari MA, Arora S, YadavM. Sustainability issues on rice–wheat cropping system, International Soil and Water Conservation Research. 2016; 4(1):64-74.
 6. Gupta R, Seth A. A review of resource conserving technologies for sustainable management of the rice–wheat cropping systems of the Indo-Gangetic plains (IGP). Crop Protection. 2007;26(3):436-447.
 7. Sharma S, Padbhushan R, KumarU. Integrated nutrient management in rice–wheat cropping system: An evidence on sustainability in the Indian subcontinent through meta-analysis. Agronomy. 2019;9: 71.
 8. Basak N, Mandal B, Rai AK, Basak P. Soil quality and productivity improvement: Indian story. Proceedings of the Indian National Science Academy. 2021;87: 2–10.
 9. Alaamer SA, Alsharifi SK, Shtewy N. Wheat parameters under influence of the depths and cultivation machinery, IOP Conf. Series: Earth and Environmental Science 1060; 2022.
 10. Subbiah BV, Asija GL. A rapid method for the estimation of available nitrogen in soils. Current Science. 1956;25:259–260.
 11. Jat L, Singh Y, Meena S, Meena S Parihar M, Jatav H, Meena R, Meena V. Does integrated nutrient management (INM), enhance agricultural productivity?. Journal of Pure and Applied Microbiology. 2015; 9(2):1211-1221
 12. Ramalingam K, Velayudham K, Somasundaram E, Muthukrishnan P. Effect of Integrated Nutrient Management on Yield and Economics of Banana. Global Journal of Biology and Health Science. 2020;2(4):191-195.
 13. Rathore VS, Singh P, Gautam RC. Productivity and water use efficiency of rainfed pearl millet (*pennisetum glaucum*) as influenced by planting pattern and integrated nutrient management. Indian Journal of Agronomy. 2006;51(1):46-48.
 14. Antil RS, Bar-Tal A, Fine P, Hadas A. Predicting nitrogen and carbon mineralization of composted manure and sewage sludge in soil. Compost Science & Utilization. 2013;19: 33-43.
 15. Singh G, Jalota SK, Singh Y. Manuring and residue management effects on physical properties of a soil under the rice-wheat system in Punjab, India. Soil and Tillage Research. 2007;94:229-238.
 16. Jackson ML. Methods of chemical analysis. Prentice Hall of India (Pvt.) Ltd., New Delhi; 1973.
 17. Puri AN. A new method of estimating total carbonates in soils. Pusa Bulletin, No. 73, Imperial Agriculture Research, New Delhi; 1930.
 18. Watanabe FS, Olsen SR. Test of an ascorbic acid method for determining phosphorus in water and NaHCO₃ extracts from the soil. Soil Science Society of America Journal. 1965;29: 677-678.
 19. Gomez KA, Gomez AA. Statistical procedures for agricultural research. New York: Wiley Interscience; 1984.
 20. Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. Soil Science. 1934;37:29–38.
 21. Swarup A, Wanjari RH. Three decades of all India coordinated research long term fertilizer experiments for studying changes in fertility crop productivity and sustainability. Bhopal, India: Indian Institute of Soil Science; 2000.
 22. Swarup A, Yaduvanshi NPS. Effect of integrated nutrient management on soil properties and yield of rice in alkali soil. Journal of the Indian Society of Soil Science. 2000;48:279-282.
 23. Hati KM, Swarup A, Mishra B, Manna MC, Wanjari RH, Mandal KG, Misra AK. Impact of long-term application of fertilizer, manure and lime under intensive cropping on physical properties and organic carbon content of an Alfisol. Geoderma. 2008; 148:173–179.
 24. Yadav A, Kumar A, Prakash V, Kumar N, Tiwari A, Yadav RK. Effect of integrated nutrient management on soil properties, yield attributes and yield of wheat (*Triticum aestivum* L.). International Journal of Current Microbiology and Applied Science. 2017;6(10):225-228.
 25. Lal Bahadur, Tiwari DD, Mishra J, Gupta BR. Effect of integrated nutrient

- management on yield, microbial population and changes in soil properties under rice-wheat cropping system in sodic soil. *Journal of the Indian Society of Soil Science*. 2012;60(4):326-329.
26. Parvathi E, Venkaiah K, Munaswamy V, Naidu MVS, Tand GK, Prasad TVNKV. Long-term effect of manure and fertilizers on the physical and chemical properties of an alfisol under semi-arid rainfed conditions. *International Journal of Agricultural Sciences*. 2013;3(4): 500-505.
 27. Gudadhe N, Dhonde MB, Hirwe NA. Effect of integrated nutrient management on soil properties under cotton-chickpea cropping sequence in vertisols of Deccan plateau of India. *Indian Journal of Agricultural Research*. 2015;49 (3):207-214.
 28. Tiwari A, Tiwari A, Singh NB, Kumar A. Effect of Integrated nutrient management (INM) on soil properties, yield and economics of rice (*Oryza sativa* L.). *Research in Environment and Life Sciences*. 2017;10(7): 640-644.
 29. Singh AK. Evaluation of soil quality under integrated nutrient management. *Journal of the Indian Society of Soil Science*. 2007; 55:58-61.
 30. Manna MC, Swarup A, Wanjari RH, Ravankar HN, Mishra B, Saha MN, Singh YV, Sahi DK, Sarap PA. Long-term effect of fertilizer and manure application on soil organic carbon storage, soil quality and yield sustainability under sub-humid and semi-arid tropical India. *Field Crops Research*. 2004;(93):264-280.
 31. Shtewy N, Ibrahim JH, Alsharifi SK. Effect of mechanical properties on some growth characteristics for wheat crop .*Plant Archives*. 2020;20(1);3141-3148.
 32. Santhy P, Muthuvel P, Selvi D. Status and impact of organic matter fractions on yield, uptake and available nutrients in a Long-Term Fertilizer Experiment. *Journal of the Indian Society of Soil Science*. 2001; 49:281-285.
 33. Mandal B. Soil organic carbon research in India - A way forward. The 29th Prof. J.N Mukhaljee ISSS Foundation Lecture. Delivered on 16th Nov. 2011 at the 76'h Annual Convention of Indian Society of Soil Science held at University of Agricultural Sciences, Dharwad, Karnataka; 2011.
 34. Verma G, Sharma RP, Sharma SP, Subehia SK, Shambhavi S. Changes in soil fertility status of maize-wheat system due to long-term use of chemical fertilizers and amendments in an alfisol. *Plant Soil Environment*. 2012;58(12):529–533.
 35. Kumari G, Thakur SK, Kumar N, Mishra B. Long term effect of fertilizers, manure and lime on yield sustainability and soil organic carbon status under maize (*Zea mays*) – wheat (*Triticum aestivum*) cropping system in Alfisols. *Indian Journal of Agronomy*. 2013;58(2):152-158.
 36. Khan AM, Kirmani NA, Wani FS. Effect of INM on soil carbon pools, soil quality and sustainability in rice-brown sarson cropping system of Kashmir Valley. *International Journal of Current Microbiology and Applied Sciences*. 2017; 6(7):785-809.
 37. Prakash YS, Bhadoria PBS and Rakshit A. Comparative efficacy of organic manures on the changes in soil properties and nutrient availability in Alfisol. *Journal of the Indian Society of Soil Science*. 2002; 50:219-221.
 38. Laxminarayana K. Effect of nitrogen, phosphorus and potassium on yield performance of maize under mid hills of Mizoram. *Indian Journal of Hill Farming*. 2001;14 (1):132-135.
 39. Rathore DS, Purohit HS, Yadav BL, Sharma SR. Effect of Integrated Nutrient Management on Soil Properties and Crop Yield under Black Gram-Wheat Cropping System in a Typic Haplustept. *Annals of Arid Zone*. 2011;50(1):21-26.
 40. Hati KM, Swarup A, Singh D, Misra AK and Ghosh PK. Long-term continuous cropping, fertilization and manuring effects on physical properties and organic carbon content of a sandy loam soil. *Austria Journal of the Soil Research*. 2006;44: 487–495.
 41. Sarkar AK, Mahapatra P, Kumar A. Management of macronutrients in acid soils. In Rattan, R.K. (Nutrient management acid soils. *Bulletin of the Indian Society of Soil Science*. 2007;25: 10-26.
 42. Brar BS, Singh J, Singh G, Kaur G. Effects of long term application of inorganic and organic fertilizers on soil organic carbon and physical properties in maize–wheat rotation. *Agronomy*. 2015;5:220-238.

43. Sharma SS, Totawat KL, Shyamura RL. Characterization of salt affected soils of Southern Rajasthan. *Journal of Indian Society Soil Science*. 2004;52(3): 209-214.
44. Sarkar R, Basavarajl B, Kar S. Influence of calcium on distribution of different forms of iron in Vertisols. *Agropedology*. 2006; 16(1):32-36.
45. Ingole AJ, Vaidya PH, Aundhakar AV, Ghode MK. Effect of moisture and calcium carbonate content on chemical properties of soil and yield of cotton in Nanded District of Maharashtra. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(5): 1971-1976.
46. Dubey L, Dwivedi AK, Dubey M. Long term application of fertilizer and manures on physico-chemical properties and n, p and k uptake in soybean-wheat cropping system. *Plant Archives*. 2015;(15)1: 143-147.

© 2022 Kurmi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92380>