



Comparative Analysis on the Proximate Composition of Processed Cassava Products Obtained from January to March, 2023 in Lafia Town, Nigeria

**Idongesit V. Edet ^{a*}, Timothy M. Akpomie ^a,
Abel U. Augustine ^a, Teghtegh F. Samoh ^b,
Ekponoudim J. Ekam ^c and Dumebi D. Nzegbuna ^c**

^a *Department of Chemistry, Faculty of Science, Federal University of Lafia, PMB 146, Lafia, Nassarawa State, Nigeria.*

^b *Department of Chemistry, Faculty of Physical Sciences, University of Ilorin, PMB 1515, Kwara State, Nigeria.*

^c *Department of Chemistry, University of Lagos, PMB 12003, Lagos State, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Author IVE designed the study, performed the experimental analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors TFS, EJE and DDN performed the statistical analysis. Authors TMA and AUA managed, supervised and proofread the work. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CSJI/2023/v32i4854

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103334>

Original Research Article

Received: 19/05/2023

Accepted: 21/07/2023

Published: 28/07/2023

ABSTRACT

The extensive applicability of cassava and its derivatives in various industries in Nigeria is constantly increasing, thus the necessity to evaluate the chemical composition in order to ascertain the nutritional value of these products. This study investigated and compared the proximate composition of eight samples of processed cassava products (Niji[®] Foods Cassava Flour, IFGREEN[®] Odourless Fufu Flour, Ayoola[®] Fufu Flour, Aiteefills[®] Fufu flour, Niji[®] Foods Garri, Golden Penny Garri, GGEE[®] foods Ijebu Garri and local brand cassava starch) obtained from

*Corresponding author: E-mail: idongesitval2@gmail.com;

supermarkets and local markets in Lafia town from January to March, 2023 using standard official methods for proximate analysis. The results were analyzed by Minitab version 20.0 by one way ANOVA and pair-wise comparison was made post hoc using Tukey t-tests. The moisture, ash, crude protein, crude fibre, crude fats and carbohydrate ranged from 4.34 – 12.70 %, 0.34 – 1.63 %, 1.30 – 10.06 %, 0.02 – 0.81 %, 4.01 – 12.53 % and 70.83 – 85.73 %. The study revealed that the results of the proximate composition in all the samples varied significantly ($P \leq .05$). Crude protein and Crude fibre contents in all the samples agreed with the recommended limits by FAO/WHO and SON. The high moisture content in Aiteefills Fufu Flour may impart a shorter shelf life on the product. Findings from the study also suggested the possibility of formation of metal-ion pigment complexes in GGEE® foods Ijebu Garri, Ayoola® Fufu Flour, Niji® Foods Cassava Flour and Local Brand Cassava Starch due to high ash contents. However, all samples meet the basic nutritional requirements for crude protein, crude fibre, fats and carbohydrate contents in cassava products. The study recommended regular routine proximate composition checks on new and existing products available to consumers to further maintain high nutritional standards in processed cassava products.

Keywords: Cassava; proximate composition; processed; nutritional value.

1. INTRODUCTION

“Cassava, *Manihot esculanta* Crantz, now widely grown in the Pacific is a domesticated plant derived from one or more species of the Genus *Manihot* in the Euphorbiaceae family” [1]. “The plant parts used are the storage root (tuber) and leaves. The *Manihot* genus is reported to have about 100 species, among which the only commercially cultivated species is *Manihot esculanta* Crantz. There are two distinct types of cassava plants: erect, with or without branching at the top type and the spreading type” [2]. “The plant is characterized by palmate lobed leaves, inconspicuous flowers and a large, starchy, tuberous root with a tough papery brown bark and white to yellow flesh” [3]. “It is one of the most perishable tuber crops with a high postharvest loss” [4,5].

“Being a drought-tolerant crop, cassava can be grown in areas with uncertain rainfall patterns which usually results in unsuccessful cultivation of many other crops. Recently, the world cassava production stands at 302.66 million metric tonnes with leading countries like Nigeria, Congo DR, Thailand and Indonesia ranked 1st, 2nd, 3rd and 4th respectively” [6]. “In Africa, the production of cassava stood at 193.62 million metric tonnes in 2020, thus positioning Africa as the world’s largest cassava growing region. Unarguably, Nigeria remains the highest producer of cassava in the world with about 60 million metric tonnes in 2020” [6]. “Cassava is a starchy staple whose roots are very rich in carbohydrates, a major source of energy. Apart from sugarcane, cassava has the highest carbohydrate contents amongst the crop plants” [7]. In the aspect of food sufficiency and food crisis alleviation in Africa,

the role of cassava cannot be over emphasized [8]. Considering the huge instrumentality of Cassava in this regard, it has been recognized as Africa’s food security crop.

“Food processing involves the transformation of food from its raw form into a new form by the application of varying techniques” [9]. Food processing is vital since it ensures the improvement of shelf life of the products, reduce spoilage and increase the availability and accessibility of the food products to people. Fresh cassava roots contain about 70 % moisture content thereby increasing its bulkiness and this poses serious transportation problems which may affect the availability of the food crop to users. Additionally, studies have confirmed the high health risks associated with the consumption of cassava in its raw form owing to the presence of high level of toxic Hydrogen cyanide in the fresh roots. In recent years, the processing of cassava into various products has gained significant attention due to its versatility, cyanide reduction capacity, long shelf life, and potential for value addition thus ensuring food security. The processed forms of cassava finds wide applications both as food and useful industrial products. Tapioca, farina, garri, fufu, starch etc. are some of the product often gotten from processing of cassava tuber locally. The different processed forms are the cheapest source of staple food in Africa and Nigeria in particular. Nigeria consumes nearly all that it produces, regardless of the high level of cassava production [10,11]. The processing of cassava into various forms involves the application of different techniques, each of which affects significantly the chemical composition of the final products. Traditionally, soaking, fermentation,

cooking, steaming and chipping, frying, drying and roasting are the different processing steps employed in the processing of the cassava roots. The factors affecting the chemical composition of processed cassava products include the cultivar, geographical location, maturity stage of the plant, environmental conditions and processing methods [12,13].

Proximate composition refers to the basic components of a food product, including moisture, protein, fat, fibre, ash, and carbohydrates. Over the years, research efforts have been directed towards characterizing the proximate composition of various species of cassava roots and derived products to unravel the effects of processing methods, environmental factors, and genetic diversity on their nutritional profiles [14,15,16]. Recent advances in analytical techniques and innovative research have provided valuable insights into the proximate composition of processed cassava products. In 2020, significant contributions were made to this field, shedding light on the nutritional content and potential health benefits of cassava-based foods. Findings from a study conducted by Okonkwo et al. [15] where advanced spectroscopic techniques were used to analyze the starch composition of cassava chips, revealed correlations between processing parameters and starch characteristics. Meanwhile, the work of Nzuta et al. [16] explored the nutritional variations of cassava flours derived from different cultivars and regions, revealing distinctive nutrient profiles among products.

Processed cassava products have been reported by several researchers to possess high carbohydrate content and low protein and fat content. In a study by Nwaliowe et al. [17], the proximate composition of cassava flour and its derived products was evaluated. The researchers reported that cassava flour contained approximately 80% carbohydrates, 2% protein, and negligible amounts of fat, making it a valuable source of energy. Similarly, Sajeev et al. [18] investigated the proximate composition of cassava chips and found that they contained about 80-85% carbohydrates, 1-2% protein, and low levels of fat. Additionally, research has focused on the nutritional attributes of other cassava-based products, such as cassava starch and tapioca pearls. A study by Nzuta et al. [16] analyzed the proximate composition of tapioca pearls commonly used in bubble tea. The researchers found that tapioca pearls contained approximately 85% carbohydrates, 1% protein,

and negligible amounts of fat, further emphasizing the predominance of carbohydrates in processed cassava products. The high carbohydrate profile associated with cassava products makes it a common household food in Nigeria generally, and Lafia town in particular.

Recently, the increased production of processed cassava products, poses a necessity for detailed evaluation of the chemical composition of the final products as a basis for ascertaining the nutritional value of the products. Studies on the proximate composition of fresh cassava roots have been done, but none has been done on the different processed cassava products available in local markets and supermarkets in the city of Lafia, Nassarawa State, Nigeria. Being a low-income town, many households in Lafia depend on these processed cassava products available including Gari, Fufu flour, Cassava flour and starch for their nutritional needs since they are the cheapest sources of carbohydrate. The present work reports on the proximate composition of eight different processed cassava products including Garri (Niji[®] Foods Garri, Golden Penny Garri, GGEE[®] foods Ijebu Garri), Fufu flour (IFGREEN[®] Odourless Fufu Flour, Ayoola[®] Fufu Flour, Aiteefills[®] Fufu flour), high quality cassava flour (Niji[®] Foods Cassava Flour) and starch (Local Brand Cassava starch) obtained from local markets in Lafia town. The study also compares the proximate composition of processed cassava products to ascertain their nutritional significance based on the standard health rating index.

2. MATERIALS AND METHODS

2.1 Study Site and Sample Collection

The present study was conducted in Lafia town during the period between January to March, 2023. Lafia town is the headquarters of the Lafia Local Government Area, and also the capital city of Nassarawa State in the Guinea Savanna vegetation of north central Nigeria. It is located between latitudes 8° 28' N and 8° 30' N to longitudes 8° 29' E and 8° 32' E. The local government area has a population of 361 000 [19], and is bordered by three local government areas; Wamba in the north, Nassarawa Eggon in the north-west, Obi in the south, Doma in the south-west, and Plateau state in the east (Fig. 1). Lafia is geologically a part of the lower Benue trough and the major occupations of the inhabitants are civil service, farming, mining, artisanry, and fishing. Samples consisting of

products containing cassava as primary ingredients were randomly obtained directly from supermarkets and local markets in Lafia town from January to March, 2023. The eight samples of processed cassava products were divided into three groups based on the method of processing: garri samples, fufu flour, high quality composite flour and locally processed starch. The samples include Niji[®] Foods Cassava Flour, IFGREEN[®] Odourless Fufu Flour, Ayoola[®] Fufu Flour, Aiteefills[®] Fufu flour, Niji[®] Foods Garri, Golden Penny Garri, GGEE[®] foods Ijebu Garri and local brand cassava starch. They were designated as NFCF, IGFF, AYFF, ATFF, NFG, GPG, GIG and LBCS respectively. The samples were transported directly to the Chemistry laboratory, Faculty of Agriculture, Nassarawa State University, Shabu Campus, for analysis.

2.2 Preparation of Samples

100 g of each processed cassava product sample was put in tightly sealed envelopes and kept in field cellophane bags before analysis to prevent environmental contamination.

2.3 Proximate Analysis

Proximate composition, namely, moisture, crude protein, crude fibre, crude fats, ash and nitrogen free extract (carbohydrate contents) in portions of the various samples were measured according to the standard methods as described in AOAC [21] and other standard techniques obtained from the literature.

2.3.1 Moisture determination (% MD)

The moisture content was determined according to AOAC [21] method 925.10. 2 g of the samples was accurately weighed into a pre-labelled, preweighed beaker and transferred to a vacuum dry oven to dry at a temperature of 130°C. The samples were heated within a time range of 1 h, 1 h -30 min, 2 h, 2 h -30 min, and 3 h respectively, and weighed till constant weight was achieved. All sampling and analysis were done in triplicate [13]. The formula used for the calculation of moisture content and dry matter can be seen below:

Sample weight – moisture content = dry matter

$$\% \text{ moisture content} = \frac{\text{dry matter}}{\text{weight of sample}} \times 100 \dots 2.1$$

2.3.2 Ash (% Ash)

The ash content was determined according to AOAC [21] method 923.03. Prepared samples were weighed into preweighed, porcelain crucibles. The samples were transferred to a muffle furnace (J M Ney furnace, model 2-525) and ashed at 550°C for 8 h. The crucibles were allowed to cool in desiccators and then weighed [22]. The formula that was used for the calculation of ash content can be seen below:

$$\% \text{ Ash} = \frac{\text{Wt of Ash}}{\text{weight of sample}} \times 100 \dots\dots\dots 2.2$$

2.3.3 Fat (% Fat)

The fat content in the samples were determined according to 920.39 of AOAC [21] by dissolving 8 g of the cassava samples in a 200 cm³ beaker containing 8.4 cm³ of hydrochloric acid and heated in a water bath for 1 h. After heating, the sample solution were allowed to cool and then extracted with petroleum ether in a separating funnel. After extraction, the sample solution were heated to dryness and the weight collected after cooling [22]. The formula that was used for the calculation of fat content can be seen below:

$$\% \text{ Fat} = \frac{\text{Weight loss of sample (extract)}}{\text{weight of sample}} \times 100 \dots\dots 2.3$$

2.3.4 Crude protein (% C.P)

The crude protein content was determined as described in Nuwamanya et al. [23] using Dumas combustion method of nitrogen content analysis (Leco Truspec Model FP-528, St Joseph Mi, USA) by taking about 0.3 g of sample and using the conversion factor:

$$\% \text{ protein} = \% \text{ N} \times 6.25 \dots\dots\dots 2.4$$

2.3.5 Crude fibre content (% C.F)

The crude fibre content was determined using the method 962.09 of AOAC [21]. About 0.5 g of the sample was boiled in 50 mL of 0.3 M H₂SO₄ under reflux for 30 min, followed by filtering through a 75 mm sieve under suction pressure. The residue was washed with distilled water to remove the acid. The residue was then boiled in 100 mL, 0.25 M sodium hydroxide under reflux for 30 min and filtered under suction. The insoluble portion was washed with hot distilled water to free the alkaline. The insoluble portion




Fig. 1. Map of Nassarawa state showing Lafia Local Government Area [20]

was dried to the constant weight in the oven at 100 °C, for 2 h, then cooled in the desiccator. The dried sample was ashed in a muffle furnace to subtract the mass of ash from the fibre after then the % of fibre was determined [24].

2.3.6 Nitrogen Free Extract (% NFE) as carbohydrate content

NFE was determined by mathematical calculation. It was obtained by subtracting the sum of percentages of all the nutrients already determined from 100.

$$\% \text{ NFE} = 100 - (\% \text{ moisture} + \% \text{ CF} + \% \text{ CP} + \% \text{ EE} + \% \text{ Ash}) \dots\dots\dots 2.5$$

% NFE represents soluble carbohydrates and other digestible and easily utilizable non-nitrogenous substances in the samples.

2.4 Data Analysis

Data was analyzed using a one-way analysis of variance (ANOVA). The mean differences were determined using the Tukey's Least Significance Difference test at 5% significant level. Values of $p \leq 0.05$ were considered statistically significant.

All data were expressed as the mean \pm standard deviation (SD) of three observations. All calculations were done using the Minitab version 20 software.

3. RESULTS AND DISCUSSION

The proximate composition of the cassava products is shown in Fig. 2. Eight samples of different processed cassava products were randomly selected from the supermarkets and local markets in Lafia town, Nassarawa State, Nigeria and analyzed for their proximate composition using the standard methods as described by AOAC [21].

3.1 Moisture Content (% MD)

The moisture content of the processed cassava samples ranged from 4.34 - 12.70% and varied ($P = .05$) among the products (see Fig. 3). LBSCS and AYFF were not significantly different ($P > .05$). ATFF, AYFF, NFCF and LBSCS gave relatively higher moisture content compared to NFG, GPG, GIG and IGFF. As expected, the garri samples (NFG, GPG and GIG) had low values of moisture content due to the roasting stage involved during their processing stage.



Fig. 2. Proximate composition of the eight processed cassava products




Fig. 3. Moisture contents of the eight processed cassava products

Conversely, the moisture content in IGFF was lower than ATFF and AYFF. The moisture content of Aitefills Fufu flour (ATFF) was significantly the highest ($P = .05$). This is followed by LBCS, AYFF, NFCF, NFG, GPG, GIG and IGFF being the least. Manano et al. [25] reported moisture contents of cassava varieties in Uganda in the average range 5.43–10.87% which is similar to results obtained from the current study. The differences in moisture

contents could be attributed to differences in chemical constituents and processing methods. Moisture is an important parameter in the storage of cassava flour. Very high levels greater than 12% allow for microbial growth and thus low levels are favourable and give relatively longer shelf life [26]. The low content of moisture in IGFF may be attributed to the effect of commercial production considerations and shelf life where the intention of the producers must

have been to provide products that can store for a longer duration before purchase by consumers or before spoilage. The effect of insufficient drying during the processing stage may have contributed to the high moisture content in ATFF (12.70%). All the fufu flour samples (AYFF and IGFF) except ATFF had moisture contents within the permissible limit as stipulated by the Standard Organization of Nigeria (SON) and FAO/WHO limits. Similarly, all the garri samples, (GIG, GPG and NFG), cassava flour (NFCF) and cassava starch (LBCS) showed moisture contents that were within the standard limits of SON and FAO/WHO.

3.2 Ash Contents (% Ash)

The ash contents were in the average range of 0.34% – 1.63% and varied ($P = .05$) among the samples (see Fig. 4). GIG (1.63%) had the highest ash content while IGFF (0.34%) had the lowest. Ash contents of the cassava fufu flour samples (IGFF, AYFF, ATFF) except the high quality cassava flour sample (NFCF) were relatively lower than the garri samples (GIG, GPG, NFG). However, the ash contents of IGFF, AYFF, GPG and NFG were within the standard limits of SON while NFCF, AYFF, LBCS and GIG had values beyond the standard limits of SON and FAO/WHO. Crude ash content is usually indicative of inorganic constituents (minerals such as K, Zn and Ca) and generally including cassava, it ranges from 1% to 2%. Ash contents represent the total mineral content in food after it has been burnt at a very high temperature. The ash contents reported in previous studies: 1.46 – 2.71% [27], 1.90 – 2.84% [28] and 1.44 – 2.35% [29] are relatively higher as compared to the result obtained in the current study. Omowonuola et al. [30] also reported ash content in samples of cassava flour as 3.49% which is significantly higher than that obtained in this study. The differences in reported ash contents could be attributed to differences in dry matter contents, genotypic form of the raw cassava roots and their proximate composition. It has been reported that higher dry matter contents were associated with lower ash contents [30]. This could also be attributed to higher fibre contents in GIG (0.81%) as shown in Table 1. Fibre was a major contributor to ash contents in the samples. In a related study, wheat flour varieties with higher fibre contents had higher ash contents [31]. In the food industry, ash contents may influence the quality of flours. High ash contents can impact the whiteness of flours. Since ash content is also

an indicator of mineral contents, increased mineral content may promote metal chelating activities to form metal-ion pigments complexes [32] which can confer greenness/redness or yellowness colour on the final product. Justifiably, the possibility of formation of metal ion-pigment complexes in NFCF, AYFF, LBCS and GIG is high since the product samples had ash contents above the recommended limits.

3.3 Crude Protein (% CP)

The protein contents were in the range of 1.30% – 10.06%. IGFF and LBCS as well as ATFF and GIG were comparable ($P > .05$) (see Fig. 5). The protein in other samples were significantly different ($P = .05$). The composite flour sample, NFCF (10.06%) had the highest crude protein content significantly, shortly followed by GPG (5.25%), a garri sample. GIG had the least crude protein content in this study. The introduction of palm oil during the roasting stage is likely to be responsible for the value of the protein content in GPG. Manano et al. [25] reported protein in the range 0.74% - 1.52%. Emmanuel et al. [28] reported in the range 1.76% - 3.46% protein. Other authors have reported lower protein values in the range 0.3% - 0.6% protein [33,34,35] and 0.72% protein [36]. The differences in protein contents can be accounted for in terms of environmental conditions such as soil fertility [13,12] from which the cassava tubers used in making the products were obtained from. The application of nitrogen rich fertilizer to the soil contributes to the increased protein contents in cassava as shown by Shittu et al. [37] where the protein range increased from 4.3% - 19.30% in unfertilized cassava varieties to 9.6 – 20.9% in fertilized varieties. However, these results are alarmingly too high levels for protein in cassava flours, garri and fufu [38]. The high levels however, could be attributed to additional nitrogen from cyanides during alkaline distillation of acid-digested samples. While it is not completely and clearly understood, the nitrogen in cyanide compounds can contribute to the crude content of nitrogen levels attributed to proteins. During drying, saccharides replaces water molecules bonded to proteins. The elimination of water may alter the binding sites of proteins which affect their activities, and presumably decreasing the protein contents. In the current study, all samples had protein contents above the minimum values (0.5% for starch and 1.0% for other cassava products) as recommended by SON.




Fig. 4. Ash contents of the eight processed cassava products




Fig. 5. Crude protein contents of the eight processed cassava products

3.4 Crude Fats (% Fats)

The crude fats contents were in the range of 4.01% - 12.53% and varied significantly ($P = .05$) among the samples (see Fig. 6). All fufu flour samples (IGFF and AYFF) recorded high fat contents except ATFF. NFCF and LBCS also had high fat contents. All the garri samples (GIG, GPG and NFG) had relatively lower fat contents. Previous studies have reported fats in the range 0.1% - 0.3% [39,35], 0.74% - 1.49% [28], 0.41%

[36] which are significantly lower than results obtained in the present study. Fat can act as an alternative energy source. Studies have shown that lower fats had decreased moisture contents which proximately increases dry matter contents, and subsequently total carbohydrates. Fats such as monoglycerides and phospholipids can form liquid-crystalline phase with water through hydrophilic (polar-heads) or hydrophobic (methyl) groups. The high fats contents in the cassava fufu flour samples may be due to microbial

metabolism spin off associated with the fermentation period. Most of the garri samples were not yellow garri indicating that the samples might not have been fortified with palm oil which are rich in β -carotene as well as fat matter during the processing stage.

3.5 Crude Fibre (% CF)

The fibre contents ranged from 0.02% - 0.81%. Fibre contents of NFCF, ATFF, AYFF and LBCS are significantly ($P > .05$) comparable (see Fig. 7). Expectedly, the fibre contents of the garri samples (GIG, GPG and NFG) was relatively higher compared to LBCS, NFCF, IGFF, AYFF

and ATFF. There was a relative comparability amongst NFCF-ATFF-AYFF and LBCS-AYFF-NFCF. High fibrous cassava would be characteristically coarse while less fibrous is likely to be finer. Higher dry matter contents are likely to be associated with high amounts of fibre and larger flour particle size. Fibre contents is related to ash contents. High fibrous cassava roots during dewatering process could indicate loss of mineral contents [38]. There could also be increased rate of nutrients release (loss) in highly permeable fibres during processing [40]. Edible fibres are mainly composed of polysaccharides such as cellulose, hemicellulose and pectins.




Fig. 6. Fat contents of the eight processed cassava products




Fig. 7. Crude fibre contents of the eight processed cassava products

Fibre can take up water easily like a wick, however this water is loosely bound in the fibre structure and can easily be lost during drying resulting in decreased moisture contents. When fibre is present along with starch, it competes for the limited amount of water available in food system. Pectin functions as a plasticizer and control porosity [41], and depending on porosity, there could be differential moisture responses among the product samples. The proximate increase in fibre resulted in decreased protein and lipid contents of flour samples. Fibre contents in cassava flours were observed to increase while protein and lipid contents decreased. This was similar to the results reported by Oluwaniyi and Oladipo [42]. The variation in fibre can be alluded to differential genetic varieties of cassava. Excess fiber in the diet will increase fecal nitrogen, cause intestinal irritation and reduce protein digestibility used for production [43].

3.6 Total Carbohydrates (% NFE)

The total carbohydrate content was expressed in form of % NFE and was obtained by difference. Carbohydrate contents ranged from 70.38% - 85.73% (see Fig. 8). The total carbohydrate contents were high in all the garri samples (GPG, GIG and NFG) compared to the flour samples. However, one of the fufu flour samples, IGFF, had a high carbohydrate content of 80.59% which was close to the range of the carbohydrate content in the garri samples. All samples were significantly different ($P = .05$). Similar results

were reported by Charles et al. [44] in the range 80.1 - 86.3% carbohydrate and 84.32 - 86.57% [45]. Protein, lipids and moisture contents are the major components impacting carbohydrates and decrease in these molecules would lead to significant increase in the total carbohydrates [14]. Carbohydrates binds proteins through hydrogen bonding via hydroxyl group on saccharides and amine group on proteins [44] which may result in highly carbonyl substituted carbohydrate and subsequently loss of protein activity and availability. Carbohydrate interact with lipids to form glycolipids through glycosidic bond [45] which reduces free lipids. The carbohydrates binds water molecules through hydrogen bonding [46] hence limiting water mobility which justifies the inverse relationship between moisture and carbohydrates as shown in Table 1.

Tharise et al. [47] showed that composite cassava flour had 8.51% moisture, 10.6% ash, 4.98% protein, 0.65% fat as well as 2.62% fibre in its composition which were quite different as compared to values obtained in this current study. The difference in the values may be attributed to the fact that the composite flour contains a mix of other blends which will impact additional nutritive value to the flour as opposed to the cassava flour in the current study (NFCF) which is purely high quality cassava flour and contains no mix. It may be argued that the results obtained by the researchers was due to the non-processing effects, since current research only involved processed samples.




Fig. 8. Nitrogen Free Extract (Carbohydrate content) of the eight processed cassava products

4. CONCLUSION

The eight processed cassava products showed more comparable results with each other based on the analyzed nutritional composition implying that the various methods of processing and genotypic nature of the raw forms of the cassava samples affected the nutritional contents of the products respectively. All samples showed crude fibre contents within the recommended levels and protein contents above the recommended limits of SON. The moisture contents of all the samples except that of ATFF were within the standard limits of SON and FAO/WHO implying that the products will all have a longer shelf life except ATFF which may have a shorter shelf life since the moisture content is greater than 12 % as recommended by CODEX Alimentarius and SON. The present study also revealed that the possibility of formation of metal ion-pigment complexes in GIG, AYFF, NCF and LBCS is high due to high ash contents and may affect the primary desirable quality of flours and starch (AYFF, NCF and LBCS) suitable for use in the food and non-food industry. Furthermore, IGFF, AYFF, NCF and LBCS had high fat contents while GIG, GPG and NFG had relatively lower fat contents. The carbohydrate contents of all samples were within acceptable ranges except for LBCS which had a carbohydrate content lower than the minimum carbohydrate level recommended by the benchmark used in this study.

5. RECOMMENDATIONS

1. Enhanced and strategic effort should be made by the relevant agencies of the government in Nigeria such as National Agency for Food and Drug Administration Control (NAFDAC) to educate, monitor and sensitize the local manufacturers and farmers on the need to use more improved varieties, better cultivation methods and modern techniques of cassava processing to further minimize the production of raw cassava roots with poor nutritional quality and the health risk posed to the consumers for adopting short-cut processing techniques.
2. Proximate composition studies should be carried out on new and existing processed cassava products available in other local markets in Nigeria.

ACKNOWLEDGEMENTS

We are sincerely grateful to Mr Tsaku, the Chief Laboratory Technologist and the entire

management of the Chemistry Laboratory, Faculty of Agriculture, Nassarawa State University, Shabu Campus, Lafia for granting access to carry out the experimental analysis of the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Nyamekye CA. Health issues related to the production and consumption of cassava as a staple food (Master's thesis, Norwegian University of Life Sciences, As); 2021.
2. Dhasan DB, Karthik K, Reddy ML, Yadav MGS. Identification and classification of cassava plant leaf disease using deep learning technique. In AIP Conference Proceedings. AIP Publishing LLC. 2022; 2519(1):050026.
3. Kidasi PC, Chao DK, Obudho EO, Mwang'ombe AW. Farmers' sources and varieties of cassava planting materials in Coastal Kenya. *Frontiers in Sustainable Food Systems*. 2021;5:611089.
4. Musah S, Kwakye E, Kudaddze S, Adzawla W. Analyzing cassava processors'preferred output and profitability of cassava processing in Hohoe municipality. *UDS International Journal of Development*. 2020;7(1):262-272.
5. Adjei LP, Arthur EK, Gikunoo E, Foli G, Quaye-Ballard JA, Agyemang FO, Baah DS. Cyanide contamination assessment via target survey and physicochemical and bacteriological characterization: A case study of Akrofrom-Techiman cassava processing area in Ghana. *Environmental Monitoring and Assessment*. 2023;195(4): 482.
6. FAOSTAT [Internet]; 2020. [Cited 24 February 2020]. Available:<http://www.fao.org/faostat/en/#data/QC/visualize>
7. Chandrasekara A, Kumar TJ. Roots and tuber crops as functional foods: a review on phytochemical constituents and their potential health benefits. *International Journal of Food Science*; 2016. Available:<https://doi.org/10.1155/2016/3631647>
8. Alamu EO, Ntawuruhunga P, Chibwe T, Mukuka I, Chiona M. Evaluation of

- cassava processing and utilization at household level in Zambia. Food Security. 2019;11(1):141-150.
9. Singla M, Sit N. Application of ultrasound in combination with other technologies in food processing: A review. Ultrasonics Sonochemistry. 2021;73:105506.
 10. Isabirye Z. The distribution and abundance of bondar's nesting whitefly (*paraleyrodes bondari*) on cassava in busukuma sub county, wakiso district, uganda (Doctoral dissertation, Makerere University); 2021.
 11. Ogunyinka O, Oguntuase A. Analysis of cassava production and processing by various groups in support of cassava value chain in the south west of Nigeria. ISABB Journal of Food and Agricultural Sciences. 2020;9(1):11-19.
 12. Agiriga A, Iwe M. Optimization of Chemical Properties of Cassava Varieties Harvested at Different Times using Response Surface Methodology. Am J Adv Food Sci Technol. 2016;4:10–21.
 13. Burns AE, Gleadow RM, Zacarias AM. Variations in the chemical composition of cassava (*Manihot esculenta* Crantz) leaves and roots as affected by genotypic and environmental variation. J Agric Food Chem. 2012a;60:4946–4956.
 14. Chisenga SM, Workneh TS, Bultosa G, Laing M. Proximate composition, cyanide contents, and particle size distribution of cassava flour from cassava varieties in Zambia. AIMS Agriculture & Food. 2019; 4(4).
 15. Okonkwo CE, Olaniran AF, Adeyi O, Adeyi AJ, Ojediran JO, Adewumi AD, Erinle OC. Drying characteristics of fermented-cooked cassava chips used in the production of complementary food: Mathematical and Gaussian process regression modeling approaches. Journal of Food Process Engineering. 2021;44(7):e13715.
 16. Nzuta RA, Elechi JO, Isa IS, Kittika'a HJ, Thomas CT. Flour and Cassava (*Manihot* spp) Cultivars in Nigeria: Understanding their Chemical Functionalities and Acceptability in Food Systems. Int. J. Sci. Res. in Chemical Sciences. 2022;9(2).
 17. Nwaliowe GN, Alamu EO, Sanusi RA, Maziya-Dixon B. Evaluation of improved cassava (*Manihot esculenta* Crantz) varieties and associated products for proximate, cyanogenic potential and glycemic indices. CyTA-Journal of Food. 2023;21(1):20-30.
 18. Sajeev MS, Padmaja G, Jyothi AN, Krishnakumar T, Pradeepika C. Tropical Tuber Crops: Nutrition and Entrepreneurial Opportunities. In Vegetables for Nutrition and Entrepreneurship. Singapore: Springer Nature Singapore. 2023;409-438.
 19. Demographic Statistics Bulletin. National Bureau of Statistics; 2020.
 20. Ezekiel A, Dahiru MK, Abdullahi SB. Households Perception on the Generation and Disposal of Solid Waste in Lafia Metropolis of Nasarawa State, Nigeria. African Journal of Environment and Natural Science Research. 2021;4(4):16-26. DOI: 10.52589/AJENSRIFFSP0F7D
 21. AOAC. Official Methods of Analysis of AOAC international. Gaithersburg, Maryland, USA: 19th edition, AOAC International; 2012.
 22. Kenneth VAR. Evaluation of three cassava varieties for tuber quality and yield. Gladstone Road Agricultural Centrecrop Research Report No.4; 2011.
 23. Nuwamanya E, Baguma Y, Rubaihayo P. Physicochemical and functional characteristics of cassava starch in Ugandan varieties and their progenies. J Plant Breed Crop Sci. 2010;2:001–011.
 24. Manano J, Ogwok P, Byarugaba-Bazirake GW. Chemical composition of major cassava varieties in Uganda, targeted for industrialisation. J Food Res. 2017;7:1.
 25. Turner ER, Luo Y, Buchanan RL. Microgreen nutrition, food safety, and shelf life: A review. Journal of Food Science. 2020;85(4):870-882.
 26. Rojas CC, Nair B, Herbas A, Bergenstahl B. Proximal composition and mineral contents of six varieties of cassava (*Mannihot esculenta*, Crantz), from Bolivia. Revista Boliviana de Química. 2007; 24(1):71-77.
 27. Emmanuel O, Clement A, Agnes S. Chemical composition and cyanogenic potential of traditional and high yielding CMD resistant cassava (*Manihot esculenta* Crantz) varieties. Int Food Res J. 2012; 19:175–181.
 28. Eleazu C, Eleazu K. Determination of the proximate composition, total carotenoid, reducing sugars and residual cyanide levels of flours of 6 new yellow and white cassava (*Manihot esculenta* Crantz) varieties. Am J Food Technol. 2012;7:642–649.
 29. Kanwate BW, Ballari RV, Kudre TG. Influence of spray-drying, freeze-drying

- and vacuum-drying on physicochemical and functional properties of gelatin from *Labeo rohita* swim bladder. *International Journal of Biological Macromolecules*. 2019;121:135-141.
30. Omowonuola AOA, Mary EM, Fidelis AF, Olalekan ASA, Sunday AA. Quality Characteristics of Fermented Cassava Flour (Lafun) Produced Using Backslopping Method. *EC Nutrition*. 2017;7(2):52-57.
 31. Pavlovich-Abril A, Rouzaud-Sández O, Romero-Baranzini AL. Relationships between chemical composition and quality-related characteristics in bread making with wheat flour—fine bran blends. *J Food Qual*. 2015;38:30–39.
 32. Yu L, Nanguet AL. Comparison of antioxidant properties of refined and whole wheat flour and bread. *Antioxidants*. 2013;2:370–383.
 33. Charles AL, Sriroth K, Huang T. Proximate composition, mineral contents, hydrogen cyanide and phytic acid of 5 cassava genotypes. *Food Chem*. 2005b;92 615–620.
 34. Montagnac JA, Davis CR, Tanumihardjo SA. Nutritional value of cassava for use as a staple food and recent advances for improvement. *Compr. Rev. Food Sci. Food Saf*. 2009a;8:181–194.
 35. Somendrika MAD, Wickramasinghe I, Wansapala MAJ, Peiris S. Analyzing Proximate Composition Of Macro Nutrients of Sri Lankan Cassava Variety Kirikawadi; 2016.
 36. Shittu TA, Raji AO, Sanni LO. Bread from composite cassava-wheat flour: I. Effect of baking time and temperature on some physical properties of bread loaf. *Food Research International*. 2007;40(2):280-290.
 37. Grundy MML, Edwards CH, Mackie AR, Gidley MJ, Butterworth PJ, Ellis PR. Re-evaluation of the mechanisms of dietary fibre and implications for macronutrient bioaccessibility, digestion and postprandial metabolism. *British Journal of Nutrition*. 2016;116(5):816-833.
 38. Gojiya D, Davara P, Gohil V, Dabhi M. Process standardization for formulating protein-augmented corn-based extrudates using defatted sesame flour (DSF): Sesame oil industry waste valorization. *Journal of Food Processing and Preservation*. 2022;e17203.
 39. Do DT, Singh J, Oey I, Singh H. Biomimetic plant foods: Structural design and functionality. *Trends in Food Science & Technology*. 2018;82:46-59.
 40. Oluwaniyi O, Oladipo J. Comparative studies on the phytochemicals, nutrients and antinutrients content of cassava varieties. *Journal of the Turkish Chemical Society Section A: Chemistry*. 2017;4(3): 661-674.
 41. Jha R, Fohse JM, Tiwari UP, Li L, Willing BP. Dietary fiber and intestinal health of monogastric animals. *Frontiers in Veterinary Science*. 2019;6:48.
 42. Charles A, Chang Y, Ko W, Sriroth K, Huang T. Influence of amylopectin structure and amylose content on the gelling properties of five cultivars of cassava starches. *Journal of Agriculture and Food Chemistry*. 2005a;53:2717-2725.
 43. Shadrach I, Banji A, Adebayo O. Nutraceutical potential of ripe and unripe plantain peels: A comparative study. *Chem Int*. 2020;6(2):83-90.
 44. Del Carmen Fernández-Alonso M, Díaz D, Alvaro Berbis M, et al. Protein-carbohydrate interactions studied by NMR: from molecular recognition to drug design. *Curr Protein Pept Sci*. 2012;13:816–830.
 45. Pomeranz Y, Chung O. Interaction of lipids with proteins and carbohydrates in breadmaking. *J Am Oil Chem Soc*. 1978;55:285–289.
 46. Li L, Wang N, Ma S, et al. Relationship of moisture status and quality characteristics of fresh wet noodles prepared from different grade wheat flours from flour milling streams. *J Chem*. 2018;2018:1–8.
 47. Tharise N, Julianti E, Nurminah M. Evaluation of physico-chemical and functional properties of composite flour from cassava, rice, potato, soybean and xanthan gum as alternative of wheat flour. *International Food Research Journal*. 2014;21(4).

APPENDIX

Table 1. Proximate composition of the processed cassava products

	Garri samples				Cassava flour samples			Composite flour sample		Starch sample		
	Standard	GIG	GPG	NFG	Standard	IGFF	ATFF	AYFF	Standard	NFCF	Standard	LBCS
% MD	7.0 (max)	5.41 ^f ± 0.01	5.49 ^e ± 0.01	6.84 ^d ± 0.02	10.0 (max)	4.34 ^e ± 0.02	9.32 ^b ± 0.01	12.70 ^a ± 0.01	12.0 (max)	9.07 ^c ± 0.06	12.0 (max)	9.35 ^b ± 0.02
% Ash	1.5 (max)	1.63 ^a ± 0.02	1.54 ^b ± 0.02	1.39 ^c ± 0.01	0.6 (max)	0.34 ^b ± 0.01	0.50 ^e ± 0.01	0.99 ^e ± 0.01	0.7 (max)	1.05 ^d ± 0.01	0.1 (max)	0.89 ^f ± 0.01
% CP	1.0 (min)	1.30 ^f ± 0.02	5.25 ^b ± 0.01	3.50 ^c ± 0.01	1.0 (min)	2.19 ^d ± 0.01	1.75 ^e ± 0.01	1.30 ^f ± 0.01	1.0 (min)	10.06 ^a ± 0.02	0.5 (min)	2.18 ^d ± 0.02
% Fat	-	5.13 ^e ± 0.01	4.01 ^h ± 0.02	5.82 ^f ± 0.01	-	12.53 ^a ± 0.01	10.46 ^b ± 0.01	6.61 ^e ± 0.01	-	9.39 ^d ± 0.01	-	9.43 ^c ± 0.01
% CF	2.0 (max)	0.81 ^a ± 0.01	0.74 ^b ± 0.02	0.53 ^c ± 0.01	2.0 (max)	0.02 ^f ± 0.01	0.07 ^{de} ± 0.01	0.05 ^e ± 0.01	1.5 (max)	0.06 ^{de} ± 0.01	0.2 (max)	0.08 ^d ± 0.01
% NF	65 – 70 (min)	85.73 [±]		81.93 ^c ± 0.03	65 – 70 (min)	80.59 ^d ± 0.03	77.90 ^e ± 0.02	78.35 ^e ± 0.02	65 – 70 (min)	70.38 ^e ± 0.09	85 (min)	78.09 ^f ± 0.02

Values are means of triplicate determinations ± standard deviation. Means on the same column with different superscript differ ($p \leq 0.05$) significantly. Max = Maximum; Min = Minimum; Standard = FAO/WHO and SON nutritional standards.

© 2023 Edet et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/103334>