



Studies on Mortality and Histopathological Alteration on the Gills of *Oreochromis niloticus* Juveniles Following Exposure to Ethanolic Extract of *Phragmites capitata* under Laboratory Conditions

U. U. George ^{a*}, A. J. Otoh ^a, O. O. Ajayi ^b and I. E. George ^c

^a Department of Fisheries and Aquaculture, Faculty of Agriculture, Akwa Ibom State University, Ikot Akpaden, Akwa Ibom State, Nigeria.

^b Department of Pathology, Faculty of Basic Clinical Sciences, University of Uyo, Akwa Ibom State, Nigeria.

^c Department of Zoology, Akwa Ibom State University, Ikot Akpaden, Mkpata Enin, Akwa Ibom State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Authors UUG and AJO did conception, design and development of the topic, data collection and analysis, initial drafting and reviewing the manuscript and final approval of the prepared manuscript. Author OOA did conception, design and development of the protocol, supervision of the experiments, data analysis and reviewing the manuscript. Author IEG is student who actually played the supporting role in the experiments and reviewing the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2023/v24i3634

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103963>

Original Research Article

Received: 30/05/2023
Accepted: 03/08/2023
Published: 10/08/2023

*Corresponding author: Email: ubonggeorge@aksu.edu.ng;

ABSTRACT

The study was conducted using two replicate (Batch A and B) for 96 hours under controlled laboratory conditions, with five different concentrations of the extract ranging from 0 to 40 mg/l. The results of the study revealed that the extract had a concentration-dependent toxic effect on the test organisms, with induced behavioural changes such as abnormal and uncoordinated swimming movement, restlessness, respiratory difficulties, and attempts at jumping out. The alterations of physico-chemical parameters of the test water were also observed, indicating the adverse effects of the extract on the aquatic environment. The 96 hours LC50 for *Oreochromis niloticus* was determined to be 22.124 mg/l representing a log-transformed concentration of 1.345 mg/l. This is the concentration where 50% of the test organisms is expected to die at the end of the 96 hours' bioassay. The histopathological examination of the gills revealed pathological changes in the test organisms at concentrations of 20, 30, and 40 mg/l whereas no histomorphological changes were observed at 0 and 10 mg/l concentrations of the extract. The result of the present findings suggests that ethanolic extract of *P. capitata* had severe impacts on the test organisms resulting in mortality and gill damage.

Keywords: *Phragmenthera capitata*; mortality; gills; histomorphological; *Oreochromis niloticus*.

1. INTRODUCTION

The study aims to assess the toxic impacts of ethanolic extract of mistletoe on the survival and histopathology of the gills of Nile tilapia (*Oreochromis niloticus*). This research is important because it helps determine the safety of using mistletoe extract as a medicinal substance and also as a plant-based additive in fish feed formulation. By evaluating the survival rate and examining the gill tissue of tilapia exposed to different concentrations of the extract, the study provides valuable information on the toxicity of mistletoe extract on aquatic species.

Mistletoe is a parasitic plant commonly used in traditional medicine for its medicinal properties. However, the safety and efficacy of using mistletoe extract as a therapeutic agent have not been thoroughly studied. In particular, its potential impact on the gill, a vital organ for respiration, is not well understood.

Phragmenthera capitata is a species of plant in the family Loranthaceae, commonly known as the African mistletoe. It is native to Africa and temperate Asia. The genus contains over 35 species (Flanagan, 1993; Cernusak, 2004). It is a hemi-parasitic plant that grows on various host trees, mostly in tropical and subtropical regions of Africa (Flanagan, 1993; Cernusak, 2004). *Phragmenthera capitata* has oval-shaped leaves that are about 1-3 cm long and 0.5-1.5 cm wide. Its flowers are small and inconspicuous, with greenish-yellow petals that are fused at the base. The plant produces small, spherical berries that

are about 0.5 cm in diameter and contain one or two seeds (Flanagan, 1993; Cernusak, 2004). In traditional African medicine, various parts of the *Phragmenthera capitata* plant are used to treat various ailments, including diarrhea, dysentery, stomach pains, and skin diseases. The plant also has cultural significance in some African communities and is used in various rituals and ceremonies.

Nile tilapia is a widely cultivated and consumed fish species, making it a valuable model organism for studying the toxic effects of substances in aquatic environments. The study of the toxic impacts of ethanolic extract of mistletoe on the survival and histopathology of the gills of Nile tilapia (*Oreochromis niloticus*) aims to fill this gap in knowledge and contribute to the safe use of mistletoe extract as a feed additive. In addition to monitoring survival and mortality, the study also examines the gill tissue of the fish to determine the effects of the extract on the gill morphology. This includes evaluating changes in the structure and function of the gill's cells and the presence of any pathological conditions.

However, the safety and efficacy of mistletoe extract as a therapeutic agent have not been thoroughly studied, and more research is needed to fully understand its potential health benefits and toxic effects. Despite its widespread use in traditional medicine, the use of mistletoe extract in medical practice should be done with caution, and its safety and efficacy should be established through rigorous scientific investigation. It is important to carry out independent toxicity studies

on the extract of *Phragmenthera capitata* to ascertain its level of toxicity and make useful recommendations for its effective use in the treatments of ailments and as a source of additive in fish feed formulation.

2. MATERIALS AND METHODS

2.1 Collection of Test Organism

Juveniles of *Oreochromis niloticus* were collected from Akwa Ibom State University fish farm, Obio Akpa Akwa Ibom State, Nigeria located within 4^o 57'52" N and 7^o 45'29" E. The climate of the area is tropical and is characterized by distinct wet and dry seasons. The vegetation of the study area is a rainforest close to the mangrove belt. Human activities in the area include farming, hunting, boat building and sand mining. A total of two hundred (200) Juveniles were collected and used for the toxicity study.

2.2 Acclimatization of Specimen's

The juveniles were acclimatized in a recirculatory glass aquarium measuring 96 x 50 x 29 cm containing fresh water for 24 hours in the fisheries and aquaculture laboratory of Akwa Ibom State fish farm. This enhanced the stability of the juveniles from stress of collection and transportation [1,2].

2.3 Collection of Plant Sample

Fresh leaves of mistletoe (*Phragmenthera Capitata*) growing on Avocado Leaf (*Persea Americana*) was collected for the study. The collection site of the plant was Ikot Udota in Afaha Eket Local Government Area, Akwa Ibom State. The date of Collection was 20th January, 2023. The plants material was taken for identification and authentication by a plant systematics at the Department of Botany Herbarium, Akwa Ibom State University, Ikot Akpaden, Mkpate Enin Local Government Area.

2.4 Preparation of Plant Material

After the identification, the leaves were washed and sun dried. The leaves were shredded and spread on cellophane and allowed to dry for 72 hours under room temperature. The dried leaves were pulverized (grinded) into fine powder using wooden pestle and mortar.

2.5 Preparation of Ethanolic Extract (Maceration and Extraction)

Cold extraction method (Maceration) was used in this research according to Hidayat and Wulandari [3]. In the extraction procedure, 1000ml of 99% Concentrated Ethanol was used to Macerate 240g of the plant materials in an airtight container and kept in the laboratory under room temperature for 72 hours (3 days). In the due date of filtration, the mixture was filtered with Muslim cloth to acquire the filtrate. The extract was stored in 250ml conical flasks. The conical flask was well labelled, the mouth of the conical flask was covered with foil paper and masking tape rapped around the mouth to ensure that it is tightly covered.

2.6 Preparation of Experimental Aquaria

Ten (10) rectangular plastic aquaria measuring 25 x 10 x 15 cm were thoroughly washed with tap water and properly rinsed with fresh water of similar salinity and allowed to drain dry for 24 hours on the laboratory bench based on Dede and Kaglo [4].

2.7 Stocking of Specimen

Each of the Ten (10) plastic aquaria was filled with two liters of fresh tap water and 10 juveniles of *Oreochromis niloticus* was stocked in each aquarium. The ethanolic extract of mistletoe (*P. capitata*) with varying concentrations was added to each stocked aquarium and allowed to stand for 96 hours for mortality examination. A preliminary test was conducted to give the actual variations in concentration to be used for the bioassay. Each of the aquarium had a replicate to ensure accuracy.

2.8 Monitoring of Water Quality

Water Quality Parameters was monitored prior to commencement of the experiment and also periodically according to Standard Method [5]. Parameters that were monitored include dissolve Oxygen (DO), pH, And Temperature (^oC). Temperature and pH were measured using portable pH /Ec/ TDs/ Temperature HANNA, H1 991301 Model instrument while oxygen was measured using digital portable analyser JPB - 607A from "Search Tech Instrument".

2.9 Monitoring of Specimen for Mortality

The effects of the various concentration of the ethanolic extract of mistletoe (*P. capitata*) on the juveniles of *O. niloticus* was monitored on a 24 hours' basis for 96 hours as recommended by Udo et. al. [2] and Ekanem and Ekpo [6].

2.9.1 Determination of mortality and survival rates of juveniles

The percentage mortality and survival rates of the juveniles of *O. niloticus* in the different concentrations of the ethanolic extract of *P. capitata* during the period of study was determine using the formula;

$$\% \text{ mortality} = n/N \times 100 \text{ (Chan, 1977).}$$

Where;

n = number of dead fish per aquarium per concentration

N = Total Individual Stocked

The difference between dead fish and survivors will give the percentage survival of the juveniles at the end of the experiment (96 hours) [2].

2.9.2 Determination of mortality lethal median concentration (96 hours LC_{50})

The effects of the various concentrations of the ethanolic extract of plant (*Phragmanthera capitata*) on the juveniles of *O. niloticus* was determined by graphical method (Probit Level Determination as recommended by Omoregie [7], Omoregie and Ufodike [8], Ekanem and Ekpo [6] and Udo et al. [2]. At Lethal Median Concentration LC_{50} , after 96 hours of test, the number of juveniles that are expected to die was determined from the graph. Similarly, the concentration that will kill 5% of the stocked juveniles at the end of the test (96 hours) was determined at the probit level (Omoregie, [7] Omoregie and Ufodike [8], Udo et al. [2]; Ekanem and Ekpo [6].

2.10 Collection of Sample for Histopathology Examination

The gill's tissues of *O. niloticus* juveniles were isolated from the test animal and fixed in formalin-saline for 48 hours. The fixed tissue was processed manually through graded ethanol,

cleared in xylene impregnated and embedded in paraffin wax, sections of the tissue sample were cut with a rotary microtome, stained by hematoxylin and eosin technique, prepared tissues were finally observed using a microscope for pathological changes at x100 and x400 magnification.

2.11 Data Analysis

The results of the respective concentration effects of the ethanolic extract of *Phragmanthera capitata* was presented in tables. Two-way analysis of variance (ANOVA) was used to test for significant ($P=0.05$) difference considering the extract concentration and mortality time. Also, the LD_{50} was determined using Probit analysis. All statistics were carried out using SPSS version 20.0.

3. RESULTS

3.1 Initial Water Quality Parameters

The initial water quality parameters prior to stocking are shown in Table 1. Dissolved oxygen had a value of 5.2 mg/l, with a value of 29.8°C for Temperature and 6.77 for pH.

3.2 Variation in water Quality (Physico-chemical parameters) in the test media with *Oreochromis niloticus* as test organism (Batches A and B) during the experimental period (96 Hours)

Table 2 shows the variation recorded in the different physico-chemical parameters for *O. niloticus* (Batches A and B) in the different concentration of the toxicant and time.

Dissolved oxygen concentrations ranged between 4.00 – 5.2 mg/l in the Omg/l concentration of the toxicant. The highest DO value was recorded at the 24 hours of test with the least value recorded during the 96 Hour of Test.

In the 10 mg/l concentration of toxicant, dissolved oxygen ranged between 2.8 – 4.8 mg/l. the least value was recorded during the 96 hours of test while the highest value was recorded during the 24 hours of test.

Table 1. Initial Physico-chemical parameters of the test water prior to stocking of test organism

Fish Species	Initial physico-chemical parameters prior to stocking		
	DO (mg/l)	Temp (°C)	pH
<i>Oreochromis niloticus</i>	5.2	29.8	6.77

Table 2. Summary of the variations in the physico-chemical parameters in the test media during the experimental period

Parameters	Conc. (mg/l)	Batch A				Batch B			
		24 hrs	48 hrs	72 hrs	96 hrs	24 hrs	48 hrs	72 hrs	96 hrs
Dissolved Oxygen (mg/l)	0	5.2	4.7	4.4	4.0	5.2	4.7	4.4	4.0
Initial: 5.2 mg/l	20	4.8	4.0	3.2	2.8	4.8	4.0	3.2	2.8
	30	4.1	3.6	3.6	3.4	4.1	3.6	3.6	3.4
Initial: 5.2 mg/l	40	3.8	3.6	3.4	2.8	3.8	3.6	3.4	2.8
	50	3.2	3.0	2.6	2.4	3.2	3.0	2.6	2.4
Temperature (°C)	0	27.7	27.6	27.5	27.6	27.7	27.6	27.5	27.6
	20	27.5	27.4	27.3	27.2	27.5	27.4	27.3	27.2
	30	27.6	27.5	27.3	27.3	27.6	27.5	27.3	27.3
	40	27.4	27.4	27.2	27.3	27.4	27.4	27.2	27.3
Initial: 29.8 °C	50	27.4	27.3	27.1	27.1	27.4	27.3	27.1	27.1
pH	0	6.31	6.25	6.22	6.12	6.31	6.25	6.22	6.12
	20	6.26	6.22	6.12	6.00	6.26	6.22	6.12	6.00
Initial: 6.77	30	6.21	6.20	6.00	5.80	6.21	6.20	6.00	5.80
	40	6.21	6.22	6.22	6.12	6.21	6.22	6.22	6.12
	50	6.16	6.05	6.02	6.00	6.16	6.05	6.02	6.00

In the 20 and 30 mg/l concentration of toxicant, dissolved oxygen ranged 3.4 – 4.1 mg/l and 2.8 – 3.8 mg/l respectively. The highest and least value of DO were observe at the 24 hours of test and 96 hours of test for both concentrations during the study duration.

In the 40 mg/l concentration of toxicant, dissolved oxygen value ranged from 2.4 – 3.2 mg/l. The highest value was recorded during the 24 hour of test and the least value was observed during the 96 hours of test.

Temperature value were observed to range between 27.1 – 27.7°C during the 96 hours' bioassay. The least value of 27.1°C was recorded during the 72nd and 96th hours of test in the 40 mg/l concentration of the extract while the highest value of 27.7°C was recorded during the 24 hours of test in the 0 mg/l concentration of the toxicant (control).

The value of pH was observed to range between 5.80 – 6.31 during the 96 hours' experimental bioassay. The least value of 5.80 was recorded during the 96th hours of test in the 20 mg/l concentration of the toxicant while the highest value of 6.31 was recorded during the 24 hours of

test in the 0 mg/l concentration of the toxicant (control).

3.3 Summary of the Percentage Mortality and Survivors of *O. niloticus* in the Different Concentrations of the Ethanolic Extract of *P. capitata* at the end of the Experiment (96 hours)

The percentage mortality and survivors of *O. niloticus* at the end of the test period in each of the concentrations are shown in Table 3 for the two batches of the experiment.

In the 0 mg/l concentration of the extract, no mortality was recorded throughout the test period in both batches A and B. Similar observations was recorded for 10 mg/l concentration of the extract in both bathes.

During the 96-hour bioassay for 20 mg/l concentration of the extract, 40 % mortality and 60 % survivors were recorded for both batch A and B.

In the 30 mg/l concentration of the extract, 80 % mortality and 20 % survivors were recorded while in the 40 mg/l concentration of the extract all the

test organisms were observed dead leaving 0 % survivors in both batches (Table 3). Statistical Analysis using Anova (SPSS 20.0) showed that there was no significant difference ($P = .05$) in mortality between the two batches.

3.4 96 Hours LC_{50} Determination

The 96 hours LC_{50} for *O. niloticus* exposed to the different concentrations of the ethanolic extract of *P. capitata* is shown in Table 4 for both batches. The 96 hours LC_{50} is given at 22.124 mg/l representing a log transformed concentration of 1.345 mg/l a point where 50 % of the test organisms would be killed at the end of the experiment.

3.5 Histopathology of the Gill of *O. niloticus* Exposed to the Different Concentrations of the Ethanolic Extract of *Phragmenthera capitata*.

The gills of the control group displayed histological features within the normal range. The primary filament epithelium, secondary lamella, supporting cartilage epithelium, muscle fibers, and the capillary-rich apical part of the filament were all observed to be in a healthy state. The primary filament epithelium exhibited its characteristic structure and integrity, while the secondary lamella maintained its normal morphology. The supporting cartilage epithelium

provided structural support, and the presence of well-developed muscle fibers contributed to the functionality of the gills. Additionally, the apical part of the filament exhibited a high cellularity and was abundantly supplied with capillaries, indicating an active and well-perfused region. Similar to the control group, Group 2 demonstrated normal histological features. The primary filament epithelium, secondary lamella, supporting cartilage epithelium, muscle fibers, and the capillary-rich apical part of the filament all appeared within the expected range. No significant deviations or abnormalities were observed in this group, suggesting that exposure to the ethanolic extracts of *P. capitata* did not induce histological changes in the gills. In both Group 3 and Group 4, histological analysis revealed diffuse epithelial degeneration in both the primary and secondary filaments. The degeneration appeared to affect the overall integrity and structure of the epithelial layer. Moreover, the apical part of the filament in these groups showed severe epithelial degeneration. The severe degenerative changes observed in these regions suggested a compromised condition of the gill tissue. In Group 5, complete degeneration of the secondary lamella epithelium and complete cellular loss in primary lamella was observed. The complete degeneration of the secondary lamella epithelium further emphasized the severity of the epithelial damage in the gill tissue of this group.

Table 3. Summary of the Percentage Mortality and survivors of *O. niloticus* in the different concentrations of the ethanolic extract of *P. capitata* at the end of the experiment (96 hours)

Conc. of extract (mg/l)	Batch A				Batch B			
	Mortality (M)	% M	Survivors (S)	% S	Mortality (M)	% M	Survivors (S)	% S
0	0	0	10	100	0	0	10	100
10	0	0	10	100	0	0	10	100
20	4	40	6	60	4	40	6	60
30	8	80	2	20	8	80	2	20
40	10	100	0	0	10	100	0	0

Table 4. LC_{50} determination for *O. niloticus* at the end of the 96-hours bioassay

Plant	Species	Probit	S. E	LC_{50} (mg/l)	Log Con. (mg/l)
<i>Phragmenthera capitata</i>	<i>Oreochromis niloticus</i>	$P = -10.735 + 7.982X$	0.111	22.124	1.345




Fig. 1. Photomicrograph of Gill arch tissue sections of Group 1-5. Haematoxylin and Eosin (H&E) stain. Each group was shown x100 and x400 magnification

Control Group (Ai&Aii) showed gills with normal primary filament epithelium (thick black arrow), secondary lamella (thin arrow), the supporting cartilage epithelium (thick blue arrow), muscle fibre (black arrowhead) and the capillary-rich apical part of the filament (blue arrowhead). (Bi&Bii) showed normal histological features as the control. (Ci&Cii) and (Di&Dii) showed both primary and secondary filament but diffused epithelial degeneration, the apical part of the filament showed severe epithelial degeneration (red arrowhead). (Ei&Eii) depicted complete disruption of the secondary lamella epithelium and acellular primary lamella (thick black arrow)

4. DISCUSSION

Three basic physico-chemical parameters were taken in line with standard practice in toxicological studies prior to stocking and introduction of the extract into the test media. A value of 5.2 mg/l was recorded for dissolved

oxygen with a value of 29.8°C recorded for temperature and a value of 6.77 was recorded for pH.

In aquaculture operations, there are standard threshold values for these parameters. For dissolved oxygen a range of between 4.0 – 6.0

mg/l is suitable, 6.7 – 8.6 for pH and 25.0 – 30.0°C for temperature are recommended values for standard operation of aquaculture [9-14].

The initial values of the physico-chemical parameters of the experimental water were found to fall within the recommended threshold limits prior to the commencement of the experiment as previously reported by the authors under reference.

The unperturbed values of physico-chemical parameters observed prior to the commencement of the experiment might be linked to the absence of impurities or the toxicant and the organisms themselves [15-17]. Impurities, pollutants and toxicants are known to play a role either to elevate or reduce the different physico-chemical parameters in aquatic environment [18-20].

Variations in physico-chemical parameters were observed in the experimental aquaria in both batches during the experimental period. The values of physico-chemical parameters recorded varied depending on time and concentration of the extract. As the concentration of the toxicant increased with time, the values of the physico-chemical parameters were observed to sway when compared with the control.

This occurrence has been previously reported in the physico-chemical parameters of the test water media by Ayotunde and Ofem [21] when reporting on the acute and chronic toxicity of *Carica papaya* seed powder to Nile tilapia (*Oreochromis niloticus*), Ayotunde et al. [22] when investigating toxicity of *Carica papaya* seed powder to *Clarias gariepinus* fingerlings and effects on haematological parameters, Ayotunde et al. [23] when investigating the toxicity of *Carica papaya* on adult catfish (*Clarias gariepinus*), and Cagauan et al. [24] when evaluating botanical piscicides on *Oreochromis niloticus* and mosquito fish *Gambusia affinis*.

It is a commonly acceptable scientific finding that concentration stimulates the elevation and / or reduction in physico-chemical parameters of test water during an experiment [25,23] coupled with the fact that the organisms will also spend their absorbing oxygen in particular for survival [26,27].

The percentage mortality of *O. niloticus* in the ethanolic extract of *phragmenthera capitata* ranged from 0 – 100 % in both batches A and B

at the end of the 96-hours bioassay. No mortality was recorded in the 0 mg/l and 10 mg/l concentration of the toxicant. However, 40 % mortality was recorded in the 20 mg/l concentration in each of the batches while 80 % mortality was recorded in the 30 mg/l concentration of the toxicant. No survivor was recorded in the 40 mg /l concentration of the toxicant. The results of the present findings agree with earlier assertion by George et al. [11] when reporting on the laboratory bioassay of the potential effect of rubber extract (*Hevea Brasiliensis*) on the Survival of fingerlings of *Oreochromis niloticus*; George et al. [12] during their studies on the effect of lethal concentrations of rubber extract (*Hevea Brasiliensis*) on the survival of fingerlings of *Clarias gariepinus* under laboratory condition; George et al. [14] when working on the toxic effect of crude oil on hatchery reared *Oreochromis niloticus* fingerlings and George et al. [28] when investigating on the acute toxic effect of crude oil on the gills of *Clarias gariepinus* juveniles.

Results from the present findings shows that percentage mortality was concentration dependent. The higher the concentration of the extract, the higher the percentage mortalities of the test organism. Similar results have been reported by different authors; Ogundiran et al. [26] when investigating toxicological impacts of detergents effluents in juveniles of African catfish (*Clarias gariepinus*), Calta, et al. [29] when studying the acute toxicity of the synthetic pyrethroid deltamethrin to young minnow carp (*Cyprinus carpio*), Ayuba et al. [30] when investigating on the acute toxicity of the root of Jimson's weed (*Datura innoxia*) to the African catfish (*Clarias gariepinus*) fingerlings and Adedeji et al. [31] when investigating acute toxicity of diazinon to African catfish (*Clarias gariepinus*) fingerlings.

The 96 hours LC₅₀ of any toxicant is the dose or concentration which kills 50 % of the stocked organisms at the end of the experimental period of 96 hours (4 days) [16,32,2,11,12,28 and 14].

The 96 hours LC₅₀ is known to vary from toxicant [5,16] and from concentration to concentration of the toxicant [24], (Ayotunde et al. 210).

In the present study the 96 hours LC₅₀ was 22.124 mg/l representing a log concentration of 1.345 for both batches (A and B). The 96 hours LC₅₀ of toxicants are known to vary as previously reported by the authors earlier cited above. In a

related study, Ogundiran, et al. [26] reported 96 hours LC₅₀ of 0.0166 mg/l and 0.0038 mg/l for batch A and B *Clarias gariepinus* fingerlings under the toxicity effects of detergent effluents, 96 hours LC₅₀ of 0.1 mg/l and 0.03 mg/l was reported by Adewoye et al. [27] when working on the effects of soap and detergent effluents on *Clarias gariepinus* fingerlings. Again Ayotunde et al. [23] reported the 96 hours LC₅₀ of 0,033 – 0.33 mg/l on *Clarias gariepinus* adults using *Carica papaya* extract. The varied 96 hours LC₅₀ values usually obtained from different toxicants and test organisms is again reported by Ekanem et al. [33], when they reported a 96 hours LC₅₀ of 5.0 ± 1.76 and 4.0 ± 1.76 mg/l for *Macrobrachium macrobrachion* and *Macrobrachium vollenhovenii*.

In this study the 96 hours LC₅₀ of 22.124 mg/l obtained for both batch A and B may be attributed to the concentration of the toxicant finally used for the bioassay.

The effects of the ethanolic extract of *Phragmenthera capitata* showed pathological effects on the gill lamellae of *Oreochromis niloticus* juveniles. However, the gill lamellae in the control (0 mg/l) and 10 mg/l concentration of the toxicant were not affected. In the 20 mg/l, 30 mg/l and 40 mg/l concentration both primary and secondary lamellae of the gills especially the apical part showed epithelial degeneration which is evident in the decreased or total loss of cellularity along the length and the apical part of the lamellae.

Pathological changes and degeneration of gill lamellae has been reported by several authors. Diana et al. [34] when investigating on the biochemical and histological effects of deltamethrin on *Carassius auratus gibelio* with different effects such as lamellae cells hypertrophy and nuclear pycnosis in the basal cells, Gabriel et al. [35] reported histopathological changes in the gills of *Clarias gariepinus* exposed to refined petroleum oil and kerosene under laboratory conditions.

The histological changes observed in the present study were concentration dependent with severe alteration been pronounced at higher concentration. The results of this findings are similar to earlier assertion reported by George et al. [36] when reporting on the acute toxic effects of *Hevea brasiliensis* on the gills of hatchery reared *Oreochromis niloticus* fingerlings and observed histological changes in the gills of the

exposed organisms which were concentration dependent, George et al. [28] when investigating on the acute toxic effect of qua iboe light crude oil on the gills of *Clarias gariepinus* juveniles; Idowu et al. [37] when studying the effect of *Euphorbia hirta* leaf extract on histopathology of juveniles *Clarias gariepinus* and George et al. [13] when reporting on the histopathological alterations in gills of fingerlings of *Clarias gariepinus* following sub-lethal acute exposure to *Hevea brasiliensis*, Audu et al. [38] reported on the histopathological effects of unrefined water fractions of the foliage of *Balanites aegyptiaca* on gills, kidney and liver of *Oreochromis niloticus* fingerlings. Similarly, Audu et al. [39] examined histological changes in gills and liver of *C. gariepinus* intoxicated with acute concentrated grades of *Vernonia amygdalina*, Adesina, et al. [40] evaluated the effect of acute toxicity of *Moringa oleifera* root extract on *O. niloticus*, Nasiruddin et al. [41] investigated the histological alterations in organs of *Heteropneustes fossilis* intoxicated with extracts of three dry seed, while Oluwatoyin, [42] studied the toxic effect of *Ipomoea aquatica* leaf extract on histopathology of *O. niloticus*. The result of the present findings on effects of toxicants on gills histomorphology agrees favourably with earlier assertion as reported by authors under reference [43].

5. CONCLUSION

Effects of ethanolic extract of *Phragmenthera capitata* (mistletoes) on the survival and histopathology of the gills of *Oreochromis niloticus* juveniles were investigated using static bioassay under laboratory condition. Prior to the toxicity test of the extract on the test organism physico-chemical parameters were taken before stocking of the experimental fish. Variations were observed in the physico-chemical parameters of the test media during the experimental period. The physico-chemical parameters (DO, temperature and pH) were observed to fluctuate with increased concentration of the extract with time when compared to the control. The variation observed in the test media during the 96 hours' bioassay is attributed to the introduction of the toxicant into the experimental aquaria which resulted in the mark shift or variation of these parameters as recorded within the period under study. Percentage mortality recorded in this study was observed to be concentration dependent with higher mortality recorded at higher concentrations in both batches. From the mortality's ratio the 96-hour LC₅₀ was 22.124 mg/l for both batches. The results of

histopathology which showed pathological changes in the gills of the test organisms in the 20, 30 and 40 mg/l concentration of the toxicant except at 0 mg/l and 10 mg/l concentration of the toxicant which no pathological changes was observed depicts the mortality results which shows that percentage mortalities was concentration dependent with high mortality values recorded at higher concentrations. Also, histopathological results confirm the toxic effect of the extract at higher concentration which was evidence in the degeneration of the primary and secondary gill lamellae observed during the period of the study. Based on the result of this findings, it is imperative to implement ecologically friendly methods to control invasive species within our environment. Additionally, due to the high cost of feed in aquaculture operations, further research on toxicity studies of common plants within our environment is recommended. This would help to identify which plants has the potentials to be used as a plant-based additive in the formulation of fish feed.

6. RECOMMENDATIONS

Based on the results of the findings which showed medium to high percentage mortalities when exposed to the ethanolic extract of *Phragmenthera capitata*, it is imperative that ecological friendly methods should be put in place to checkmates invasive species within our environment. However, further research is recommended on the use of *Phragmenthera capitata* in fish feed formulation at different inclusion level to confirm the results of the present findings. This recommendation stems from the fact that the toxicity level observed during the study was low except at higher concentrations. Also, based on the high cost of feed in aquaculture operations, this study recommends further research on the toxicity studies of common plant within our environment. This will help to know which of them can be used as a plant-based component in the formulation of fish feed.

ACKNOWLEDGEMENTS

The authors sincerely acknowledge Akwa Ibom State University for Providing an ambience environment for the research. We also convey our heartfelt thanks to the laboratory technicians of the department of Medical Laboratory Science, histopathology unit, who were involved in the histo-morphological examination of the gills of the experimental animal.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Smith LS. Introduction to Fish Physiology. T. F. H. Publications Incorporated, 211 West Sylvania Avenue, Neptune. New Jersey. 1982;352.
2. Udo PJ, Ekanem AP, Eze EE. Toxicity of crude oil to early life stages of *Heterobranchus longifilis* (Cruveier and Valenciennes) Pisces: Bagridae). Tropical Environmental Research. 2006;1:450-459.
3. Hidayat R, Wulandari P. Methods of Extraction: Maceration, percolation and Decoction. Eureka Herba Indonesia. 2021;2(1):68-74.
4. Dede EB, Kagbo HD. Aqua toxicological effects of water-soluble fraction (WSF) of diesel fuel on *Oreochromis niloticus* fingerlings. Journal of Applied Science and Environmental Management. 2001;5(1): 93-96.
5. APHA (American Public Health Association). Standard Methods for the examination of water and waste waters. 20th Edn. New York. 1998;314.
6. Ekanem AP, Ekpo IA. Effects of commercial detergents on the juvenile of *Herterobranchus longifilis* (Curvier and Valenciennes). African Journal of Environmental Pollution and Health. 2008;6(1): 18 – 23.
7. Omoriegie E. Acute Toxicity of Water-Soluble Fraction of Crude Oil to the Nile Tilapia (*Oreochromis niloticus*). Bulletin of Environmental Contamination and Toxicity. 2002;68:623 – 629.
8. Omoriegie E, Ufodike BC. Effects of water-soluble fraction of crude oil on growth of the Nile Tilapia (*Oreochromis niloticus*) (L). Bulletin of Environmental Contamination and Toxicity. 2000;64:601 – 605.
9. Udo PJ. Techniques in Fish farming (Practice and Management). Wusun Publishers, Calabar, Nigeria. 2007;100.
10. Ajah PO. Fish feeding and hatchery management. Calabar, Nigeria. Jerry Commercial Productions. 2007;178.
11. George UU, Asuquo FE, Idung JU, Andem AB. Effects of Lethal Concentration of Rubber Extract (*Hevea brasiliensis*) on the survival on fingerlings of *Clarias gariepinus* Under Laboratory Conditions. Journal of

- Natural Sciences Research. 2013a;3(9): 56-60.
12. George UU, Asuquo FE, Idung JU, Andem AB. A Laboratory Bioassay of the Potential Effects of Rubber Extract (*Hevea brasiliensis*) on the survival of fingerlings of *Oreochromis niloticus*. Journal of Biology, Agriculture and Health Care. 2013b;3 (11):70-74.
 13. George UU, Joseph A, Andy JA. Histopathological Alterations in Gills of Fingerlings of *Clarias gariepinus* (Burchell, 1822) Following Sublethal Acute Exposure to *Hevea brasiliensis*. International Journal of Scientific and Technology Research. 2014b;3(9):252- 255.
 14. George UU, Etim IN, Ekanim MP, Akpan MK. Toxic Effect of Crude Oil on Hatchery Reared *Oreochromis niloticus* Fingerlings. Journal of Academia and Industrial Research. 2015;3(11):573-576.
 15. Vogels S. Quantitative chemical analysis. In Mentham, J, Denney RC, Barnes JD, Thomas, MSK (Eds.). 2000;806.
 16. Samabaswa KS, Rao KR. Toxicity of Elsan to the Indian Snakehead (*Channa punctatus*). Indian Journal of Fisheries. 1985;3:153 – 159.
 17. World Health Organization (WHO). Guidelines for drinking water quality. Recommendations. WHO Geneva. 1998;1:130.
 18. World Health Organization (WHO). Guidelines for drinking water quality. Recommendations. WHO Geneva. 1984; 1:130.
 19. Idoho-Umeh G. Pollution Assessment of Olamoro water bodies using physical, chemical and biological indices. Ph.D Thesis, University of Benin, Benin City, Nigeria. 2002;485.
 20. Gbem TT, Balogun JK, Lawal FA, Annune PA. Trace metal accumulation in *Clarias gariepinus* (Teugels) exposed to sub-lethal levels of tannery effluent. Science Total Environmental. 2001;271: 1 - 9.
 21. Ayotunde EO, Ofem BO. Acute and Chronic toxicity of pawpaw (*Carica papaya*) seed powder to Nile tilapia (*Oreochromis niloticus* Lime 1757). Advance Environmental Biology. 2005;7:101-107.
 22. Ayotunde EO, Ofem BO, Okey IB, Ikpi GU, Ochang SN, Agbam NE, Omini DO. Toxicity of pawpaw (*Carica papaya*) seed pro to sharp-tooth catfish (*Clarias gariepinus*) fingerlings and effects on haematological parameters. International Journal of Fisheries and Aquaculture. 2010;2:71-78. Available:https://doi.org/10.5897/IJFA.9000 038.
 23. Ayotunde EO, Ofem BO, Bekah AF. Toxicity of *Carica papaya* Linn: Haematological and piscidal effects on adult's catfish (*Clarias gariepinus*). Journal of Fisheries and Aquatic Science. 2011;6(3):291-308. Available:https://doi.org/10.3923/jfas.2011. 291.308
 24. Cagauan AG, Galaites MC, Fajardo LJ. Evaluation of botanical piscicides on Nile tilapia (*Oreochromis niloticus*) and mosquito fish (*Gambusia affinis*, Baird and Girard). Proceedings of the 6th International Symposium on tilapia in aquaculture, September 12-16, Manila, Philippines, American Tilapia Association and Philippine Bureau of Fisheries and Aquatic Resources. 2004;179-187.
 25. Heijenick DG, Janssen CR, Decren WM. The combined effects of pH, hardness and dissolved organic carbon on the chronic toxicity of zinc to *Daphnia magna*: Development of the surface response model. Archives Environmental Contamination Toxicology. 2003;44:210 – 217.
 26. Ogundiran MA, Fawole OO, Aderoye SD, Ayandiran TA. Toxicological impacts of detergents effluent on juveniles of African catfish (*Clarias gariepinus*) (Burchell, 1822). Agriculture and Biology Journal of North America. 2010;1(3):330 – 342.
 27. Adewoye SO. A Comparative study on the behavioural responses of *Clarias gariepinus* on exposure to Soap and detergent effluents. Advances in Applied Research. 2010;1(1):89-95. Corpus ID: 85921404
 28. George UU, Urom SE, Etanketuk N. Acute Toxic Effect of Qua Iboe Light Crude Oil on the Gills of *Clarias gariepinus* Juveniles. International Journal of Environment and Pollution Research, 2014;2(2):16-30.
 29. Calta M, Ural MS. Acute toxicity of the synthetic pyrethroid deltamethrin to young mirror carp, *Cyprinus carpio*. Fresen Environmental Bulletin. 2004;13(11a): 1179–1183
 30. Ayuba JO, Ofojekwu PC. Acute toxicity of the root of Jimson weed (*Datura innoxia*) to the African catfish (*Clarias gariepinus*)

- fingerlings. Journal of Aquatic Science. 2002;17:131-133.
Available:<https://doi.org/10.4314/JAS.V17I2.19927>
31. Adedeji BA, Adedeji AO, Adeyemo OK, Agbede SA. Acute toxicity of diaziam to the African catfish *Clarias gariepinus*. African Journal of Biotechnology. 2008;7(5): 651-654.
Available:<http://www.academicjournals.org/AJB>
32. Akpan ER, Ekanem SB, Odaro JA. Toxicity of crude oil to freshwater algae. African Journal of Fisheries and Aquaculture. 1999;1:56-61.
Available:
<http://dx.doi.org/10.1007/s00128-011-0229-8>
33. Ekanem AP, Asuquo FE, Ndiok EJ. Toxicity of crude oil to freshwater shrimp *Macrobrachium macrobrachion* and *Macrobrachium Vollenhovenii* from Nigeria Coastal water. Bulletin Environmental Contaminant and Toxicology. 2011;86:394 – 397.
34. Diana C, Andreea SC, Diana D, Huculeci R, Marieta C, Anca D. Biochemical and histological effects of deltamethrin exposure on the gills of *Carassius auratus gibelio* (Pisces: Cyprinidae) Biotechnology. 2007;40(1):65-72.
35. Gabriel UU, Ezeri CN, Amakiri EU. Haematology and gill pathology of *Clarias gariepinus* exposed to refined oil and kerosene under laboratory conditions. J. Anim. Vet. Advances. 2007;6:461-465
36. George UU, Etim IN, Ekanim MP, Akpan MK. Acute Toxic Effects of *Hevea brasiliensis* on the Gills of Hatchery Reared *Oreochromis niloticus* Fingerlings. Journal of Academia and Industrial Research. 2015b;3 (11): 562-566.
37. Idowu AA, Soetan MO, Akinde A, Popoola OC. Effect of *Euphorbia hirta* leaf extracts on histopathology of juvenile *Clarias gariepinus*, Nigerian Journal of Animal Science. 2019;21(1):96-109
38. Audu BS, Wakawa IA, Omirinde JO, Garba U, Damshit M. Histopathological alterations in organs of Nile tilapia fingerlings exposed to sublethal concentrations of aqueous crude leaves extract of desert date. Pan Afri. J Life Sci. 2020;4:59-67.
Available:[http://dx.doi.org/10.36108/pajols/0202/40\(0240\)](http://dx.doi.org/10.36108/pajols/0202/40(0240)).
39. Audu BS, Omirinde JO, Gosomji IJ, Wazhi PE. Histopathological changes in the gill and liver of *Clarias gariepinus* exposed to acute concentrations of *Vernonia amygdalina*. Ani. Res. Int. 2017;14:2576-2587
40. Adesina SK, Illoh HC, Johnny II, Jacobs IE. African mistletoes (Loranthaceae); ethnopharmacology, chemistry and medicinal values: an update. African Journal of Traditional, Complementary and Alternative Medicine.0 2013;10(3):161-170.
Available:<http://www.DOI:10.4314/AJTCA M.V10I4.26>
41. Nasiruddin M, Azadi MA, Jahan A. Histopathological changes in gill, liver and intestine of *Heteropneustes fossilis* (bloch) treated with three dry seed extracts. J Asia Soc Banglades Sci. 2012;38:217-226.
42. Oluwatoyin AS. Histopathology of Nile tilapia (*Oreochromis niloticus*) juveniles exposed to aqueous and ethanolic extracts of Ipomoea aquatic leaf. International Journal of Fisheries and Aquaculture. 2011;3:244-257.
43. Deeni YY, Sadiq NM. Antimicrobial properties and phytochemical constituents of the leaves of African mistletoe (*Tapinanthus dodonifolius* (DC) Danser) (Loranthaceae): An ethnomedicinal plant of Hausa land, Northern Nigeria. Journal of Ethnopharmacology. 2002;83:235 – 240.

© 2023 George et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/103963>