



Standardization of Rose Flower Propagation by Improving the Climatic Conditions in India

M. K. Nehra ^{a*}, Gulfam Ali ^b and Joginder Singh ^c

^a Department of Horticulture, Kisan Degree College, Simbhaoli, Hapur, U.P., India.

^b Department of Botany, Kisan Degree College, Simbhaoli, Hapur, India.

^c Department of Horticulture, J. V. College, Baraut, U.P., India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i193680

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104729>

Original Research Article

Received: 21/06/2023

Accepted: 25/08/2023

Published: 04/09/2023

ABSTRACT

To get the better quality flowers, the agricultural / Horticultural practices have to be applied at proper time and in proper way. In the present investigation the planting time of rose root-stock was standardized alongwith the way of planting in the soil beds. Planting time was taken as January, February and March because plants are very sensitive to the temperature, photo period and the air moisture. Any deviation in then even a week or a fortnight may affect yield and quality of flowers in ease of flowering plants. Another important factor in plant growth is soil, where root established and moride nourishment to the plant. In this experiment, the rose cuttings were planted on ridges, flat beds. and in polythene bags. The observations recorded showed that cuttings when planted on ridges and in flat bed yielded better flower and good plant growth. February was found to be suitable time for the better and quicker growth time.

*Corresponding author: E-mail: chmanishnehra@gmail.com;

Keywords: Rose; root stock; ridges; sprout.

1. INTRODUCTION

Every body like flowers and use them at many occasions for enjoyment, thrill and decoration. Rose is a prominent flower as far as its variety and fragrance is concerned. It is the "Queen of the flowers". In India, it was grown starting from 1840. Mary it is said that, rose growing is the happiest hobby of poor and riches around the world. Rose is a source of sweetness and release tension. Its petals are used for oil extraction. Healthy food preparations. It contains vitamin C Its cultivation was spread near the big cities to a great extent. It is grown mainly by seedlings, own root plants, budded plants where bud of one plant is transferred to other plant. These plants have two parts, the base is called root stock and scion (top shoot) Root stock is fast growing for its roots, shoots, disease free and homogenous growth. The quality of plant and flower will depend on the soil, climate and time of planting. The efforts should be made to minimize the time of growth, reduce the cultivation cost and to get better yield was the aim of this investigation. Many other workers have worked on these aspects earlier also Gorld [1], Vatsik 1973, Namjan et al 1971, Korobov, [2], Malik [3].

2. MATERIALS AND METHODS

The experiment of standardization was conducted at the Horticultural Research Station of Meerut University during the winter season of 1987-88. The climate area which is stimulated at 29°01' N latitude and 71°43'E longitude of 222 m above sea level, is semi arid with extreme weather conditions. Annual rainfall is 700 mm which occur from may to October. During this research no rain fall was required.

Temperature ranges from 45.8 to 29.4 °C from May to June and 20.6 to 4.8°C during December to January.

Treatments:

Method of planting and cuttings:

1. Planting of cutting on ridges - P₁
2. Planting of cutting on flat beds - P₂
3. Planting of cutting in polythene bags - P₃
4. One year old roofed root stocks - P₄

Time of budding:

January - T₁, February T₂, March - T₃

Treatment combinations:

P₁T₁, P₁T₂, P₁T₃, P₂T₁, P₂T₂, P₂T₃, P₃T₁, P₃T₂, P₃T₃, P₄T₁, P₄T₂, P₄T₃

Seven treatments with two factors were laid in 3 replications. So total combinations were 12. The number of cutting planted per treatment were 60 with total cuttings of 720. These were planted in the month of November 1987.

Nursery beds were prepared well by mixing the compost. Root stock (cutting of one-year-old rooted) used in trial were similar in all characters. Healthy and disease free cutting were planted in al beds. Healthy bed cuttings from mother plants were selected with well developed buds.

2.1 Cultural Operations

Cuttings were irrigated as and when required taking operations were done by hand after every irrigation to remove weeds.

2.2 Characters Studied

- A. Root stock characters:
 - No. of sprouts per cuttings
 - Length of longest sprout
 - Diameter of thickest sprout
 - No. of leaves per sprout
- B. Scion characters:
 - Days of bud break
 - Survival percent of buds.
 - Length of longest sprout,
 - Diameter of thickest sprout
 - Average no. of leaves per sprout
- C. Flowering characters:
 - Appearance of first flower bud
 - Time between bidding to opening of flower
 - Folding time of the flower
 - Diameter of full open flower
 - No. of flower per sprout

3. RESULTS AND DISCUSSION

The result of rose root stock grow parameters was recorded the different treatments and planting time were recorded in the given Table 1.

It was found that the treatment P₁T₂, P₂T₂ and P₄T₂ gave maximum survival of root stocks which was 88.66, 88.33 and 83.88 respectively. Except polythene bag planting all three methods of February planting showed better results. Bud

Table 1. The effect of planting method and time of planting on rise root stock growth and related parameters

Treatments	Percent survival	Bud break time (days)	Mean sprout no. per cutting	Longest sprout length/ cutting (cm)	No. of leaves per sprout	Thickest sprout diameter (cm)
P ₁ T ₁	50.00	24.6	1.75	4.5	10	1.26
P ₁ T ₂	88.66	12.4	1.97	7.16	13	2.83
P ₁ T ₃	10.00	25.2	5.33	12.33	16	3.73
P ₂ T ₁	43.33	24.2	2.50	6.60	16	1.20
P ₂ T ₂	88.33	12.6	2.84	11.66	13	2.83
P ₂ T ₃	33.00	25.4	8.33	24.17	15	3.66
P ₃ T ₁	08.33	26.2	1.73	4.7	10	1.16
P ₃ T ₂	06.66	22.4	2.33	9.33	13	2.30
P ₃ T ₃	08.66	21.2	4.00	7.83	11	2.16
P ₄ T ₁	51.66	12.5	6.67	41.63	25	6.50
P ₄ T ₂	83.38	10.4	7.66	41.70	33	7.35
P ₄ T ₃	55.00	20.9	7.63	56.50	37	7.50

Table 2. The effect of method and time of planting on different parts of rose cutting and root stock growth

Treat-ments	Longest sprout length (cm)	Thickest sprout diameter (cm)	Average no. of leaves per sprout	Days needed to appear flower bud	Time taken from budding to open flower	Fielding time of flower (days)	Flower dia-meter (cm)	No. of flower per sprout (No.)
P ₁	16.38	5.25	10.35	48	54	5	6.83	5*
P ₂	11.31	5.19	9.55	51	56	5	6.26	3
P ₃	4.15	3.17	5.66	53	58	4	4.83	1
P ₄	18.63*	7.18*	11.99*	47*	53*	6*	6.99*	3
T ₁	15.86	4.17	8.4	49	56	5	6.30	4
T ₂	16.99*	6.03*	10.41*	46*	52*	7*	6.55*	4*
T ₃	7.05	4.29	8.99	49	55	5	5.39	2

break time break time was also less in treatments mentioned above than other treatments i.e. 12.4, 12.6 and 10.4 days respectively. The mean number of sprouts per cutting and length sprout showed the maximum value in P₁T₂, P₂T₂, P₃T₂ and P₄T₂ treatments. But the sprout diameter and sprout thickness did not give any regular pattern. The number of leaves per sprout were more in P₁T₃, P₂T₁, P₂T₃, P₄T₁, P₄T₂, P₄T₃ treatments in comparison to other treatment P₁T₃, P₂T₂, P₂T₃, P₃T₂, P₃T₃ and maximum sprout thickness was observed in treatments P₄T₁ (6.50 cm), P₄T₂ (7.3 cm) and P₄T₃ (7.50 cm) (Table 1)[4-6].

Sprouting character of the root stock observed was in the month of March, 1988. It was one-year-old on different growth parameters.

3.1 Buds Growth Character

The longest sprout was obtained in treatment P₄ (one-year-old root stock) (18.63 cm) and smallest in polythene bags (4.15cm) similarly the budding of February (T₂) showed better growth than other planting time (16.79 cm) sprout thickness (7.18 cm), average no. of leaves (11.99) were also more in P₄ treatment and in February planting (Table 2). In these treatments the flower bud appears first at 47 and 46 days in one-year root stock and February planting respectively.

The time taken to open the flower from bud was also less than other treatments in one-year root stock planting (53 days) and February planting (52 days). Similar observations were recorded for yielding time of flower was higher in treatment P₄ (6 days) and T₂ (7 days). Diameter of flower showed the better dimensions in these treatments than other which was (6.99 cm) and 6.55 cm in P₄ and T₂ treatments respectively. The number of flowers however has not shown the better yield (Table 2).

4. CONCLUSION

In the present work the "Standardization of rose propagation" conducted to check the proper planting time of root stocks as well as the cutting of rose the way of planting in the soil for a better and quick growth of plant and bigger size of rose flowers. The better survival of cutting was found to be February planting. It is because at that time the sun shine is not more [7]. The survival percent was poorest in the month of March either on ridges or in polythene bags due to the increasing temperature and loss of humidity in

the following months. This time developed the outbreaks of buds, reduced the sprout numbers, length also. However, the sprout length was better when cuttings were planting on ridges and root stocks were used in the month of February. Sprout diameter, no. of leaves reduction is budding time and flower diameter was also more in case of root stock planting in the month of February. Our results are in accordance with the yielding of other people [8], Raj as 1972, and Thebcult 1978, Singh [9].

CONFERENCE DISCLAIMER

Some part of this manuscript was previously presented in the conference: 6th International Conference on Strategies and Challenges in Agricultural and Life Science for Food Security and Sustainable Environment (SCALFE-2023) on April 28-30, 2023 in Himachal Pradesh University, Summer Hill, Shimla, HP, India. Web Link of the proceeding: <https://www.shobhituniversity.ac.in/pdf/Souvenir-Abstract%20Book-Shimla-HPU-SCALFE-2023.pdf>

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Gorid KH. Time for planting out rose root stocks. *Hort. Abstract.* 1978;49(11): 4420.
2. Korobov VI. Certain characteristics of root stock, scion union in garden roses budded at different times. *Agri. J. Japan.* 1976;26(1):138-141.
3. Malik RG. Effect of time and method of budding of H.T. and flori bunaroses. *South. Ind. Hort.* 1974;23(304): 125-126.
4. Lutsik ED. Quick growing of roses. *Hort. Abstract.* 1973;44(6):4144.
5. Nnjan K, Horizon MK, Ravinder R. A new technique in the propagation of roses. *Madros, Agri Y.* 1971;58: 526-528.
6. Rojas MLA. A new method of grafting for the rapid propagation of roses *Rovista. ICA.*1971;7(2):111-119.
7. The Bault G, A rose grower sprouts his mind. *Hort. Abst.* 1978;49(9):3680.

8. Mottos JR. Perrante OD. Rapid rose propagation. Hort Abst. 1971:43(7): 4708.
9. Singh SN. Studies on root stocks and budding time of roses. Hort. Advan. 1978;9:39-42.

© 2023 Nehra et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104729>