



# **Influence of Seed Storage and Priming on Naturally Aged Seeds of Aromatic Joha Rice**

**Partha Pratim Bora <sup>a</sup>, Meghali Barua <sup>a\*</sup>, Sharmila D. Deka <sup>a</sup>,  
Priyanka Das <sup>b</sup> and Purna K. Barua <sup>a</sup>**

<sup>a</sup> Department of Plant Breeding & Genetics, Assam Agricultural University, Jorhat-785 013, Assam, India.

<sup>b</sup> Department of Biochemistry & Agricultural Chemistry, Assam Agricultural University, Jorhat-785 013, Assam, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Author MB did the conceptualization and experimental designing. Author PPB did the field/ lab work and data collection. Authors PKB and SDD did the suggestions and facilities provided for field/ lab work. Author PD did the guidance regarding biochemical analysis. Authors PKB, MB and PPB did the statistical analysis. Authors MB and PPB did the Manuscript preparation. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/IJECC/2023/v13i113180

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107251>

**Original Research Article**

**Received: 30/07/2023**

**Accepted: 04/10/2023**

**Published: 09/10/2023**

## **ABSTRACT**

Joha is a fine grain aromatic rice of Assam, well known in the world market. This rice is protected and tagged as a geographical indication. This study evaluated priming effects on naturally aged seeds of two varieties, *Kon Joha* and *Keteki Joha*, after 0, 3, 6 and nine months of storage under ambient conditions. Seeds were hydro-primed, osmo-primed with 5% and 10% PEG, halo primed with 1% and 2% KCl and primed with 5 ppm and 10 ppm GA<sub>3</sub> for 12 and 24 hours. Seed moisture percentage, germination percentage, germination index, mean germination time, seedling length and dry weight, seed vigour index, field emergence, seed reserve utilization rate, seed reserve use efficiency, seed reserve depletion percentage, and biochemical parameters like electrical

\*Corresponding author: E-mail: [meghali.barua@aau.ac.in](mailto:meghali.barua@aau.ac.in);

conductivity, lipid peroxidation and  $\alpha$ -amylase were observed. Seed quality was gradually deteriorating due to ageing over the storage period. The rate of deterioration was faster for seedling vigour traits than germination parameters. There was a varietal difference in the rate of deterioration; it was slower in *Kon Joha*, an indigenous variety which also showed dormancy. The priming treatments were able to ameliorate the effect of seed ageing on seed germination, seedling growth and biochemical parameters in *Kon Joha*. Treatments with 1% KCl, 10 ppm GA<sub>3</sub> and 5% PEG enhanced the germination parameters and vigour indicators. KCl (1%) priming was the best priming agent; 24 hours of priming for all agents was better than 12 hours.

**Keywords:** Aromatic rice seed; biochemical parameters; physiological parameters; priming effect; storage.

## 1. INTRODUCTION

Seed, the primary and essential input for sustainable agriculture, has the genetic information for determining the yield potential of a crop. However, being a living entity, a seed is prone to deterioration due to ageing, especially in tropical regions with high temperatures and humidity Cortelazzo et al. [1]. Seed ageing, a natural change in seeds' physiological and biochemical properties, leads to vigour loss, declined and delayed germination, slower growth, stress susceptibility, and ultimately leading to seed death [2]. It also leads to DNA damage and membrane disruption, causing lipid peroxidation and protein synthesis disruption. Lipid peroxidation-mediated membrane damage is the most significant reason for seed deterioration during seed ageing [3]. The speed of the seed ageing process depends on the seed's ability to resist degradative changes and its protection mechanism [4], but with ageing, seeds become weak and unable to resist degradative changes efficiently.

The slow or poor germination problem of aged seeds can be ameliorated through many methods, and one of the best among them is seed priming. Seed priming is a pre-sowing controlled hydration treatment in which seeds are soaked in water or solution that allows them to complete the first two germination stages but does not permit radicle emergence. Priming seeds in aerated solution allows some of the metabolic processes for germination to occur without actual germination [5], thus improving the germination rate and uniformity Basra et al. [6]. Several priming treatments, such as hydropriming, halopriming, osmo-priming, hormonal priming, bio-priming, matrix priming, and thermo-priming, are used to invigorate and repair aged seeds. Priming has successfully enhanced seed performance for several crops, including cereals and vegetables Sarika et al. [7].

Rice (*Oryza sativa* L.), of the family Poaceae, is a staple food for more than 60% of the global population. India ranks second in rice production (129.66 mt) globally [8]. The world market has an excellent demand for aromatic fine-kernel rice varieties. *Keteki Joha* and *Kon Joha* are two popular aromatic varieties. *Keteki Joha*, with an average yield of 3.5-4 tons/ ha and duration of 150-160 days, is a very popular and an improved joha (scented) rice variety developed by crossing Sabitri and Badshabhog which was released by Assam Agricultural University (AAU). It has medium, slender grain, sweet aroma, good cooking quality and excellent palatability and taste. *Kon Joha* is also a popular indigenous short slender grain aromatic rice with excellent cooking qualities, 155-165 days of duration, mainly cultivated during the *Sali* season [9]. Hence, a study was conducted to investigate the effect of seed ageing on the physiological and biochemical parameters of small-grained aromatic rice seed and assess the ameliorating effect of priming treatments on the deteriorating effect of seed ageing.

## 2. MATERIALS AND METHODS

### 2.1 Materials

Breeder seeds of *Kon Joha* and *Keteki Joha* were collected from AAU, Jorhat. Seeds of both crops were divided into four parts to be stored for 0, 3, 6 and 9 months, respectively, in standard HDPE bags after treatment with malathion @ 2 g/ kg seed. After 9 months, the surface sterilized seeds of *Kon Joha* were primed with 5% and 10% PEG, 5 and 10 ppm GA<sub>3</sub>, 1% and 2% KCl and distilled water each for 12 and 24 hours. Priming solutions were stirred at regular intervals to facilitate aeration. After the soaking, the seeds were rinsed thoroughly with distilled water and dried to their original moisture content under shade at room temperature for 4-5 days. Unprimed dry seeds were taken as control.

## 2.2 Observations

After each storage period and after priming, seeds were evaluated for biochemical parameters like electrical conductivity (EC), lipid peroxidation (LP) and  $\alpha$ -amylase (AA), and physiological parameters like moisture percentage (oven dry method), germination percentage and index (GI), mean germination time (MGT), seedling length and dry weight, seed vigour index (SVI=germination percentage X average seedling length), field emergence (FE, conducted in tubs filled with field soils, FE percentage was calculated from the number of seedlings emerged at 30<sup>th</sup> day). Other physiological parameters were seed reserve utilization rate (SRUR in mg/seed), seed reserve use efficiency (SRUE in mg/mg), and seed reserve depletion percentage (SRDP). SRUR is the initial seed dry weight (before germination) – seed residue dry weight (only cotyledon after germination), SRUE is the seedling dry weight without cotyledon/ SRUR, whereas SRDP is obtained by SRUR/ initial seed dry weight Mohammadi et al. [10].

## 2.3 Estimation of Electrical Conductivity

3 samples of 25 seeds each were weighed to 2 decimal places and then placed in beakers containing 75 ml of deionized water at room temperature for 24 hours. All beakers were covered by aluminium foil to reduce contamination. After 24 hours, the electrical conductivity of the soaking solution was determined, and the results were expressed in  $\mu$ S.

## 2.4 Estimation of Lipid Peroxidation Value [11]

TBA-TCA reagent was prepared by dissolving 0.5 g TBA in 100 ml of 20% TCA solution. MDA (Malondialdehyde) was extracted, first by dissolving 1g dry powdered seed sample in 7ml TBA-TCA reagent, then incubating at 95°C for 30 minutes in capped reaction tubes; cooled in an ice bath and centrifuged at 5000 rpm for 10 minutes. The supernatant was collected, and absorbance was read at 535 nm and 600nm against check (without test sample, keeping all other reagents intact). MDA value was calculated as:

$$\text{MDA equivalent (nmol/ml)} = [(A_{535} - A_{600}) / 155000] \times 10^6$$

Where,

$A_{535}$  = 535 nm represents the maximum absorbance of the TBA-MDA complex

$A_{600}$  = 600 nm is the correction for non-specific turbidity

155000 = molar extinction coefficient for MDA.

## 2.5 Estimation of $\alpha$ -Amylase Activity [12]

0.2 M Sodium Acetate buffer was prepared at pH 4.7 by adding 46 ml of Glacial acetic acid (60.05 g/mole= 1.15 ml and volume made up to 100 ml) and 54 ml of Sodium acetate (82 g/mole=1.6 g in 100 ml water). 0.1 M was prepared by adding 100 ml distilled water to 0.2 M Sodium acetate buffer solution. 1% Starch solution was prepared by dissolving 1 g starch in 100 ml of 0.1 M Sodium acetate buffer. DNS reagent was prepared by adding 1 g 3,5-Dinitrosalicylic acid, 0.2 g crystalline phenol and 0.05 g sodium sulphate in 100 ml of 1% NaOH. 0.01g Maltose in 100ml distilled water gave Maltose solution. 1 g of crushed sample (germinated seed having radicle length more than 1 mm) was used to extract  $\alpha$ -amylase with 5 ml of ice-cold 10mM calcium chloride solutions (prepared by dissolving 0.15 g of calcium chloride in 1L of distilled water) overnight at 4°C or 3 hours at room temperature, and then the extract was centrifuged at 6,000 rpm at 4°C for 10minutes. The supernatant was used as an enzyme source. To prepare the enzyme assay, 1ml each of starch solution and properly diluted enzyme (test sample) was pipetted out in a test tube, which was incubated at 27°C for 15 minutes. The reaction was stopped by adding 2ml DNS reagent. The solution was then heated in a hot water bath for 5 minutes. While the solutions were warm, 1 ml of 40% Potassium Sodium Tartrate (PST) was added. Then, it was cooled in running tap water. The volume was made to 10 ml by adding 5 ml water. The absorbance was read at 560nm against check (without test sample, keeping all other reagents and procedure intact). A control sample was prepared by adding all the reagents and the test sample as described before and without incubation; hence, the reaction terminated at zero time in the control tubes. This is done to reduce the error. The absorbance reading recorded in the control should be subtracted from the absorbance reading recorded in samples. For standards, a stock solution was prepared by adding 0.01g maltose in 100 ml distilled water, i.e., 100  $\mu$ g/ 1 ml distilled water. A standard solution of 0, 0.1,

0.2, 0.4, 0.8 and 1ml maltose was prepared from the stock solution by adding 1, 0.9, 0.8, 0.6, 0.2- and 0-ml distilled water respectively. A standard graph with 0-100  $\mu\text{g}$  maltose against absorbance reading was prepared. The unknown concentration of the test sample (after subtracting the absorbance reading of the control) was taken using the standard maltose curve in  $\mu\text{g}/\text{ml}$ . The concentration was then converted into  $\mu\text{g}/5\text{ ml}$  (original extraction of 5 ml). Since 5 ml is extracted from 1 g, it can be represented as  $\mu\text{g}/\text{g}$ . This value was obtained for the incubation period of 15 minutes; hence, after dividing the value by 15, the concentration will be obtained in  $\mu\text{g}/\text{min}/\text{g}$ .

### 3. RESULTS AND DISCUSSION

#### 3.1 Effects of Seed Ageing on Physiological and Biochemical Parameters of *Kon Joha* and *Keteki Joha* Seed

Seed ageing leads to seed deterioration, which was determined by the seed's loss of viability and vigour. Significant seed quality deterioration takes place at the time of storage Khalequzzaman et al. [13]. The initial seed moisture content is a crucial factor for storage. The higher the seed moisture content, the higher the rate of deterioration. In the present study, moisture per cent and MGT for both the rice varieties (*Kon Joha* and *Keteki Joha*) increased with an increase in storage duration from 0 days to 9 months; this might be the reason why seeds deteriorated in the present study with the increase in storage duration (Table 1). To assess the seed quality, germination percentage, the rate, and the germination uniformity are equally important. GI shows the relationship between germination percentage and speed of germination, whereas MGT shows the rate and time spread of germination. The higher the GI and lower the MGT, the better will be the seed vigour. In this study, germination and GI decreased, but MGT increased with increased storage duration. The germination percentage in both the rice varieties was above seed certification standard, i.e., 80% till six months, but in the ninth month, it decreased drastically. This reduction may be due to the degradation of the mitochondrial membrane, leading to a reduction in the energy supply necessary for germination Gidrol et al. [14]. A decrease in GI and an increase in MGT with the increase in storage period indicated a reduction in the number of

seeds germinated and a slower rate of germination.

Seed vigour parameters (Table 2) like SVI, seedling length and dry weight, SRUR and SRDP decreased significantly with storage period in both *Kon Joha* and *Keteki Joha*, indicating loss of vigour of seeds with ageing, but SRUE was not affected by storage period. Though germination remained above seed certification till six months, seed vigour components decreased significantly throughout the storage period. Similar conclusions on seedling vigour as a result of ageing effects were also reported by Ellis et al. [15]; Jatoi et al. [16] in their study on rice and peas, respectively. SRUR, SRUE and SRDP are known as heterotrophic seedling characteristics because they are dependent on three components - initial seed weight, seedling dry weight and weight of mobilized seed reserve Mohammadi et al. [10]. Studying these traits was crucial because we can determine the fraction of mobilized seed reserve and seedling growth from these traits. SRUR represents the rate at which seed reserves are mobilized to the growing seedling tissues, whereas SRUE represents the conversion efficiency of mobilized reserves to seedling tissues. The reduction in SRUR and SRDP may be due to a lack of energy for converting mobilized seed reserve to seedling tissues because of ageing. This may indicate that SRUR and SRDP are sensitive components of seedling growth but not SRUE Nik et al. [17].

Biochemical traits are perfect indicators of seed quality. Here, EC and LP significantly increased with an increase in storage duration, whereas  $\alpha$ -amylase decreased for both rice varieties (Table 3). EC measures the electrolyte leakage from the seed. The higher the seed deterioration rate, the higher will be the electrolyte leakage, hence the higher EC. The increase in EC may be due to the change in cellular membrane permeability due to ageing Spano et al. [18]. LP is one of the significant causes of seed deterioration, which can be determined by estimating the seed's malondialdehyde (MDA) content [19]. The increase in MDA content may be due to free radicle damage and the deterioration of the membrane, which is probably involved in LP and associated free radical oxidative stresses, leading to membrane leakage Spano et al. [18]. The higher MDA and total peroxide in aged seeds might also result from ageing-induced changes in peroxide-scavenging enzyme activity [20].  $\alpha$ -amylase is an enzyme found in the

aleurone layer of seeds. It plays a vital role in hydrolyzing the seed reserve, which in turn provides the energy for the growth of the seedling tissues. Decreased  $\alpha$ -amylase may be because of protein denaturation and lack of ATP due to ageing. The pooled mean over both the rice varieties revealed a significant effect of the storage period on the germination and its related parameters. However, there was also a varietal difference in the rate of deterioration, but not so conspicuous as the storage period (Table 1, 2, 3). *Kon Joha* showed better germination (72%), longer and heavier seedlings (7.91 cm and 3.12 mg), and higher SRUR (5.96 mg) than *Keteki Joha* (66.67%, 7.25 cm and 2.78 mg, 5.45 mg respectively). EC and lipid peroxidation were less in *Kon Joha* (15.18  $\mu$ S and 0.125 n moles/ ml) than in *Keteki Joha* (16.07  $\mu$ S and 0.136 n moles/ ml), though  $\alpha$ -amylase activity (8.29  $\mu$ g/ min/ g) was found better than *Keteki Joha* (7.44  $\mu$ g/ min/ g). This indicates a slower rate of deterioration in the *Kon Joha* than in the *Keteki Joha*. As indigenous aromatic rice, *Kon Joha* might be harder than the high-yielding fragrant *Keteki Joha*, developed by crossing Sabitri and Badshabhog Das et al. [21]. As we know, indigenous varieties are harder and very well adapted to the local conditions. Hence, might have better chances of resisting the deteriorating effect of ageing. *Kon Joha* also showed dormancy, suggesting that there might be some relation between dormancy and tolerance to seed deterioration due to seed ageing. The varietal difference in rate of deterioration was also confirmed by Jatoi et al. [16] in pea.

### 3.2 Effect of Priming Treatments on Aged Seeds of Rice Variety *Kon Joha*

Seeds lose their vigour and ability to germinate during the ageing process and ultimately become less viable Bhattacharjee et al. [22]. Many researchers reported improved quality of aged seeds through priming due to increased enzyme activity, such as antioxidant enzymes and amylases Ansari et al. [23].

Seed priming, a simple and low-cost hydration technique, is the most effective method for achieving rapid and uniform emergence and enhancing high vigour, leading to better stand establishment and yield. Four priming treatments were considered for the present study to ameliorate the effect of deterioration due to ageing. They were hydro priming (HP), halo priming (KCl), Osmo-priming (polyethylene glycol-PEG) and hormonal priming (Gibberellic acid-GA<sub>3</sub>). Priming treatments were done after

nine months of storage. *Keteki Joha* showed no germination after nine months, indicating that the seeds were dead, which was confirmed by the tetrazolium test; hence, priming treatments were ineffective in enhancing the seeds' physiological and biochemical parameters.

Priming treatments (Table 4) effectively enhanced germination, seedling vigour and biochemical parameters over control. KCl (Table 4) was most effective for successfully improving germination percentage, GI, SVI, seedling length and dry weight, SRUR, SRDP and  $\alpha$ -amylase content. KCl priming was more effective for earlier and synchronized germination and emergence in rice Basra et al. [6]; Farooq et al. [24], improving seedling fresh and dry weight in rice seeds [25], increasing wheat yield under greenhouse conditions Khaing et al. [26]. This enhancement by KCl might be due to the enhancement of K<sup>+</sup> balance, thereby activating  $\alpha$ -amylase in converting starch to reducing sugar, which in turn provides energy for the growth of seedling tissues Farooq et al. [24]. PEG effectively enhanced seedling length and dry weight, SRUR and SRDP. Osmo-priming with PEG was more effective in strengthening seedling characteristics because osmo-priming can enhance DNA replication and DNA repair and promote mobilization of reserved materials, which contribute to initiating seed germination Aryal et al. [27] and as a result of which SRUR and SRDP enhanced effectively. GA<sub>3</sub> enhanced GI, seedling length and  $\alpha$ -amylase content. The results agreed with the observations of Zareh et al. [28], who concluded the positive effect of GA<sub>3</sub> on the seedling length of wheat. The possible reason for higher seedling growth traits with GA<sub>3</sub> treatment may be that GA<sub>3</sub> is a hormone that occurs naturally in plants, affects cell growth and elongation and significantly impacts germination, vigour and nutrient uptake Debbarma et al. [29]. HP was effective in enhancing SRUR, SRDP and alpha amylase content. The faster germination rate and better seedling characteristics were obtained by soaking seeds in water, probably due to more rapid water uptake and earlier initiation of the metabolism process, which determine radicle protrusion Ghassemi-Golezanik et al. [30]. However, any priming did not influence MGT, SRUE, EC, and lipid peroxidation.

Priming durations (Table 4) play an essential role in the success of priming since insufficient soaking duration will not complete the metabolic repair process Saini et al. [31], and long soaking

**Table 1. Germination parameters of rice varieties *Kon Joha* and *Keteki Joha* as influenced by storage period**

Storage periods	Moisture (%)			Germination (%)			GI			MGT		FE (%)			
	Kon	Keteki	Mean	Kon	Keteki	Mean	Kon	Keteki	Mean	Kon	Keteki	Mean	Kon	Keteki	Mean
0 days	9.65	9.59	9.62	94.67 (76.70)	94.00 (75.95)	94.33 (76.33)	14.33	15.56	14.95	3.42	3.07	3.24	92.67 (74.32)	90.67 (72.37)	91.50 (73.35)
3 months	10.80	11.58	11.19	91.33 (72.90)	89.33 (70.95)	90.33 (71.93)	12.80	14.22	13.51	3.55	3.33	3.44	88.67 (70.34)	86.00 (69.16)	87.33 (69.75)
6 months	12.53	12.65	12.59	89.33 (70.95)	83.33 (66.96)	86.33 (68.95)	11.40	9.13	10.27	4.89	5.21	5.05	79.33 (62.97)	76.00 (61.58)	77.67 (62.27)
9 months	13.95	13.99	13.97	12.67 (20.84)	0.00 (0.03)	6.33 (10.43)	1.05	0.00	0.53	6.36	0.00	3.18	0.00 (0.03)	0.00 (0.03)	0.00 (0.03)
Mean	11.73	11.95		72.00 (60.35)	66.67 (53.47)		9.90	9.73		4.55	2.90		65.17 (51.92)	63.17 (50.79)	
SE(M)	0.28	0.14	0.16	0.67	0.81	0.53	00.22	0.19	0.15	0.20	0.11	0.11	0.54	0.97	0.56
CD <sub>5%</sub>	0.91	0.47	0.47	2.17	2.66	1.60	0.73	0.61	0.44	0.66	0.35	0.34	1.75	3.17	1.66
CD <sub>5%</sub> (variety)	NS			1.13			NS			0.24			NS		

(Data in parenthesis are Arc sine transformed data), GI= Germination Index, MGT=Mean germination time, FE=Field emergence

**Table 2. Seedling parameters of rice varieties Kon Joha and Keteki Joha as influenced by storage period**

Storage periods	SVI		Mean	Seedling length (cm)		Mean	Seedling dry weight (mg)		Mean	SRUR (mg)		Mean	SRUE (mg/mg)		Mean	SRDP		Mean
	Kon	Keteki		Kon	Keteki		Kon	Keteki		Kon	Keteki		Kon	Keteki		Kon	Keteki	
0 days	1115.3 (3.05)	1160.3 (3.06)	1137.79 (3.06)	11.69	12.05	11.87	8.66	9.67	9.17	4.17	4.28	4.23	0.480	0.490	0.485	0.65	0.75	0.70
3 months	917.3 (2.96)	840.2 (2.92)	878.75 (2.94)	10.25	9.39	9.82	7.66	6.76	7.21	4.01	3.73	3.87	0.497	0.504	0.500	0.58	0.55	0.56
6 months	762.2 (2.88)	650.4 (2.81)	706.25 (2.85)	8.53	7.56	8.05	6.24	5.37	5.81	3.54	3.10	3.32	0.513	0.520	0.517	0.47	0.43	0.45
9 months	14.9 (1.17)	0.00 (0.00)	7.43 (0.59)	1.17	0.00	0.58	1.29	0.00	0.64	0.75	0.00	0.38	0.533	0.000	0.267	0.09	0.00	0.05
Mean	702.4 (2.52)	662.7 (2.20)		7.91	7.25		5.96	5.45		3.12	2.78		0.505	0.378		0.45	0.43	
SE(M)	0.02	0.02	0.015	0.13	0.16	0.10	0.07	0.19	0.10	0.14	0.07	0.08	0.014	0.012	0.009	0.01	0.02	0.01
CD <sub>5%</sub>	0.06	0.08	0.04	0.41	0.52	0.30	0.22	0.62	0.30	0.47	0.24	0.24	NS	0.040	0.027	0.03	0.06	0.03
CD <sub>5%</sub> (var.)	0.03			0.22			0.17			0.21			0.019			NS		

SVI=Seed vigour index, SRUR= seed reserve utilization rate, SRUE= seed reserve use efficiency, SRDP= seed reserve depletion percentage

**Table 3. Biochemical parameters of Kon Joha and Keteki Joha as influenced by storage period**

Storage periods	EC (µS)		Mean	AA (µg/ min/ g fw)		Mean	LP (n moles MDA/ ml)		Mean
	Kon	Keteki		Kon	Keteki		Kon	Keteki	
0 days	12.01	13.54	12.77	11.53	12.56	12.05	0.082	0.090	0.085
3 months	14.45	15.61	15.03	9.51	8.64	9.08	0.090	0.105	0.098
6 months	15.36	16.50	15.93	8.35	6.84	7.60	0.122	0.146	0.134
9 months	18.90	18.62	18.76	3.79	1.75	2.77	0.207	0.204	0.205
Mean	15.18	16.07		8.29	7.44		0.125	0.136	
SE(M)	0.15	0.13	0.10	0.098	0.15	0.09	0.006	0.006	0.005
CD <sub>5%</sub>	0.50	0.42	0.30	0.32	0.49	0.27	0.018	0.022	0.014
CD <sub>5%</sub> (variety)	0.21			0.19			0.010		

EC= electric conductivity, AA= α-amylase, LP= lipid peroxidation

**Table 4. Germination, seedling, heterotrophic and biochemical parameters of *Kon Joha* as influenced by different priming treatments**

Treatments	Germination (%)	GI	MGT (days)	SVI	Seedling length (cm)	Seedling dry weight (mg)	SRUR (mg/ seed)	SRUE (mg/ mg)	SRDP (%)	EC ( $\mu$ s)	$\alpha$ -amylase ( $\mu$ g/ min/ g fw)	LP (nano moles MDA/ ml)
Control	12.67	1.05	6.36	14.87	1.17	0.75	1.29	0.533	0.093	18.90	3.79	0.207
Treatment	24.48	2.42	5.74	154.64	6.18	2.07	4.04	0.497	0.295	18.27	4.55	0.181
Mean												
HP	22.67	2.26	5.84	131.98	5.82	1.91	4.05	0.468	0.305	18.29	4.50	0.182
KCl	27.00	2.68	5.54	181.08	6.50	2.21	4.24	0.510	0.313	18.23	4.92	0.177
PEG	23.50	2.14	5.76	147.08	6.21	2.14	4.13	0.492	0.303	18.29	4.29	0.182
GA <sub>3</sub>	23.83	2.53	5.86	147.10	6.02	1.96	3.75	0.503	0.265	18.28	4.46	0.183
HP 12 hours	24.00	2.37	6.15	138.57	5.75	1.79	3.91	0.460	0.287	18.17	4.67	0.178
HP 24 hours	21.33	2.15	5.53	125.38	5.88	2.03	4.20	0.477	0.323	18.40	4.33	0.187
KCl 12 hours	26.67	2.50	5.64	173.88	6.35	2.19	4.21	0.508	0.310	18.25	4.89	0.178
KCl 24 hours	27.33	2.85	5.44	188.27	6.65	2.22	4.27	0.512	0.317	18.21	4.95	0.176
PEG 12 hours	22.33	2.09	5.94	134.63	6.02	2.10	4.05	0.497	0.312	18.36	4.26	0.184
PEG 24 hours	24.67	2.19	5.59	159.54	6.40	2.18	4.20	0.487	0.293	18.21	4.33	0.180
GA <sub>3</sub> 12 hours	18.67	2.03	6.14	102.40	5.47	1.47	3.03	0.492	0.192	18.56	4.16	0.192
GA <sub>3</sub> 24 hours	29.00	3.02	5.58	191.79	6.57	2.45	4.46	0.515	0.338	18.01	4.76	0.175
KCl 1%	33.00	3.52	5.54	243.78	7.36	2.87	5.05	0.558	0.373	18.02	5.50	0.164
KCl 2%	21.00	1.84	5.54	118.37	5.64	1.54	3.42	0.462	0.253	18.43	4.33	0.190
PEG 5%	24.67	2.31	5.86	162.60	6.51	2.44	4.29	0.523	0.335	18.08	4.33	0.179
PEG 10%	22.33	1.97	5.67	131.57	5.90	1.83	3.97	0.460	0.270	18.49	4.26	0.184
GA <sub>3</sub> 5 ppm	21.00	2.06	6.10	121.74	5.70	1.77	3.40	0.495	0.250	18.39	4.25	0.189
GA <sub>3</sub> 10 ppm	26.67	2.99	5.62	172.45	6.34	2.15	4.09	0.512	0.280	18.18	4.67	0.178
SE(M)	1.03	0.10	0.16	6.06	0.21	0.07	0.07	0.022	0.012	0.15	0.17	0.006
CD <sub>5%</sub>	2.98	0.30	0.47	17.50	0.62	0.23	0.21	0.065	0.037	0.45	0.48	0.018

GI= Germination Index, MGT=Mean germination time, SVI=Seed vigour index, SRUR= seed reserve utilization rate, SRUE= seed reserve use efficiency, SRDP= seed reserve depletion percentage, EC= electric conductivity, LP= lipid peroxidation



**Table 5. Pearson’s correlation coefficients among the physiological and biochemical parameters of *Kon Joha* during storage**

	MP	GP	SVI	GI	MGT	SL	SW	SRUR	SRUE	SRDP	FE	EC	AA
GP	-0.81												
SVI	-0.93	0.97**											
GI	-0.89	0.99**	0.99**										
MGT	0.97**	-0.90	-0.96**	-0.94									
SL	-0.92	0.97**	1.00**	1.00**	-0.96**								
SDW	-0.87	0.99**	0.99**	1.00**	-0.94	0.99**							
SRUR	-0.93	0.97**	1.00**	0.99**	-0.97**	1.00**	0.99**						
SRUE	1.00**	-0.83	-0.94	-0.91	0.96**	-0.94	-0.89	-0.94					
SRDP	-0.93	0.97**	1.00**	0.99**	-0.97**	1.00**	0.99**	1.00**	-0.94				
FE	-0.85	1.00**	0.98**	1.00**	-0.93	0.99**	1.00**	0.98**	-0.87	0.98**			
EC	0.97**	-0.89	-0.97**	-0.95*	0.94	-0.97**	-0.93	-0.97**	0.98**	-0.97**	-0.92		
AA	-0.96**	0.94	0.99**	0.98**	-0.96**	0.99**	0.96**	0.99**	-0.97**	0.99**	0.96**	-0.99**	
LPV	0.93	-0.97**	-0.99**	-0.99**	0.98**	-1.00**	-0.99**	-1.00**	0.93	-1.00**	-0.98**	0.95*	-0.98**

\*\*/\* = significant at 1% and 5% respectively

durations may lead to radicle emergence. *Kon Joha* seeds were primed with all the priming agents for 12 hours and 24 hours. For KCl and PEG, both the duration showed similar effects except for GI in KCl and SVI in PEG, where 24 hours of priming showed better results than 12 hours. 24 hours of GA<sub>3</sub> priming was significantly better than 12 hours in improving all the parameters and ameliorating the ageing effect except in SRUE and lipid peroxidation. The effect of hydropriming duration was observed in MGT, seedling dry weight and SRUR, where longer time was found to be better than shorter duration, indicating 24 hours as the optimum duration for priming. Chivasa et al. [32], in their study on the determination of optimum seed priming time for maize and sorghum, reported that the rate of emergence significantly increased with an increase in the priming period and 24-hour priming of maize was better in every aspect of crop emergence and early growth.

Priming concentration is also an essential factor for the success of priming, which had variable effects on the traits (Table 4). In KCl priming, 1% was better than 2% for enhancing germination per cent, GI, SVI, seedling length and dry weight, SRUR, SRDP, alpha-amylase content and reduction of lipid peroxidation, whereas in GA<sub>3</sub> higher concentration, i.e., 10 ppm was better than 5 ppm for enhancing germination, GI, SVI, seedling length and dry weight, SRUR and reduced MGT. 1% KCl also enhanced anaerobic germination and its related traits in rice Doley et al. [33], enhancing seed yield and quality in green gram Devi et al. [34]. Khan et al. [35] in sorghum concluded that soaking seeds with low concentrations of priming agents like KCl and KNO<sub>3</sub> enhances germination and rapid seed emergence. In their study on rice, Ruttanaruangboworn et al. [36] suggested that priming rice seeds with 1% KNO<sub>3</sub> is better than priming with 2% KNO<sub>3</sub> for maintaining plant vigour. For PEG priming, 5% was better than 10% for enhancing GI, SVI, seedling dry weight, and SRUR. Better performance in lower concentrations might be due to a faster imbibition rate than in higher concentrations; the seeds could complete the two phases of germination faster at lower concentrations Ruttanaruangboworn et al. [36].

#### 4. CORRELATION

GP was positively correlated with SVI, GI, seedling length and dry weight, SRUR, SRDP, field emergence and AA activity but negatively

correlated with moisture content, EC and LP; hence, these parameters are good indicators to estimate the quality of seed and their performance in the field (Table 5). High seed moisture led to delayed and poor germination (high MGT and SRUE), and a high rate of deterioration (High LP and EC) showed a positive correlation. Hence, there was a gradual deterioration of seed quality due to ageing during storage.

#### 5. CONCLUSION

It can be concluded that at the initial stage of the storage, the rate of deterioration was slower but faster at a later period, especially after 6 months and was faster in seedling vigour traits than germination parameters. Reduction in heterotrophic seedling characteristics, viz., SRUR and SRDP, led to a reduction in seedling length, but seed ageing did not affect the mobilized reserve's conversion efficiency. There was varietal difference in the rate of deterioration. The rate of deterioration in *Kon Joha*, an indigenous variety, was slower than in *Keteki Joha*. The priming treatments were able to ameliorate the effect of seed ageing on seed germination, seedling growth and biochemical parameters in *Kon Joha*. Treatments with 1% KCl, 10 ppm GA<sub>3</sub> and 5% PEG enhanced the germination parameters and vigour indicators. KCl (1%) priming was the best priming agent; 24 hours of priming for all agents was better than 12 hours. The seed germination, seedling growth and biochemical parameters varied their performance concerning different priming agents, concentration and duration. The treatments could enhance the vigour, but once the seeds are dead, treatments cannot ameliorate the effect of deterioration.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Cortelazzo AL, Coutinho J, Granjeiro PA. Storage and ageing of french beans (*Phaseolus vulgaris* L.): Effect on seed viability and vigor. Brazilian Journal of Morphological Sciences. 2005;22(2):121-128.
2. Walters C. Understanding the mechanisms and kinetics of seed aging. Seed Science Research. 1998;8(2):223-244.

3. Koostra PT, Harrington JF. Biochemical Effect of age on membranal lipids of *Cucumis sativus* L. Seed. Proceedings of the International Seed Testing Association. 1969;34:329-340.
4. Gupta A, Aneja KR. Seed deterioration in soybean varieties during storage-physiological attributes. Seed Research. 2004;32(1):26-32.
5. Bradford KJ. Manipulation of seed water relations *via* osmotic priming to improve germination under stress conditions. Horticultural Science. 1986;59(2):672-676.
6. Basra SMA, Farooq M, Wahid A, Khan MB. Rice seed invigoration by hormonal and vitamin priming. Seed Science and Technology. 2006;34(3):753-758.
7. Sarika G, Basavaraju GV, Bhanuprakash K, Chaanakeshava BC, Paramesh R, Radha BN. Investigations on seed viability and vigour of aged seeds by priming in french bean (*Phaseolus vulgaris* L.). Vegetable Science. 2013;40(2): 169-174.
8. Anonymous; 2022. Available:<https://newsonair.gov.in/Main-News-Details.aspx?id=441328>
9. Anonymous; 2019. Available:<https://www.rkbassam.in/bd/premium-quality-rice-varieties#>
10. Mohammadi H, Soltani A, Sadeghipour HR, Zeinali E. Effects of seed aging on subsequent seed reserve utilization and seedling growth in soybean. International Journal of Plant Production. 2011;5(1): 65-70.
11. Heath RL, Packer L. Photo-peroxidation in Isolated Chloroplasts: I. Kinetics and Stoichiometry of Fatty Acid Peroxidation. Archives of Biochemistry and Biophysics. 1968;125(1):180-198.
12. Khan MA, Faust MA. Effect of growth retardants on  $\alpha$ -amylase production in germinating barley seeds. Physiologia Plantarum. 1967;20(3):673-681.
13. Khalequzzaman KM, Rashid MM, Hasan MA, Reza MA. Effect of storage containers and storage periods on the seed quality of french bean (*Phaseolus vulgaris*). Bangladesh Journal of Agricultural Research. 2012;37(2):195-205.
14. Gidrol X, Noubhani A, Mocquot B, Fournier A, Pradet A. Effect of accelerated aging on protein synthesis in two legume seeds. Plant Physiology and Biochemistry. 1998;26(3):281-288.
15. Ellis RH, Hong TD, Roberts EH. Handbook of seed technology for gene banks. Principles and methodology. 1985;1.
16. Jatoi SA, Afzal M, Nasim S, Anwar R. Seed deterioration study in pea, using accelerated ageing techniques. Pakistan Journal of Biological Sciences. 2001; 4(12):1490-1494.
17. Nik SMM, Tilebeni HG, Zeinali E, Tavassoli A. Effects of seed ageing on heterotrophic seedling growth in cotton. American-Eurasian Journal of Agricultural & Environmental Sciences. 2011;10(4):653-657.
18. Spano C, Buselli R, Castiglione MR, Bottega S, Grilli I. RNases and nucleases in embryos and endosperms from naturally aged wheat seeds stored in different conditions. Journal of Plant Physiology. 2007;164(4):487-495.
19. Asakawa T, Matsushita S. Colouring condition of thiobarbituric acid test for detecting lipid hydroperoxide. Lipid. 1980;15:137-140.
20. Sung JM, Jeng TL. Lipid peroxidation and peroxide-scavenging enzymes associated with accelerated ageing of peanut seed. Physiologia Plantarum. 1994;91(1):51-55.
21. Das A, Kesari V, Rangan L. Aromatic joha rice of assam-A review. Agricultural Reviews. 2010;31(1):1-10.
22. Bhattacharjee A, Maity S, Banerjee G, Roy M, Pal C, Pal B, Chakrabarti D. Chemical induced prolongation of seed viability and stress tolerance capacity of mung bean seedlings. Seed Science and Technology. 2000;28(1):155-162.
23. Ansari O, Sharif Zadeh F. Osmo and hydro priming improvement germination characteristics and enzyme activity of mountain rye (*Secale Montanum*) seeds under drought stress. Journal of Stress Physiology and Biochemistry. 2012; 8(4):253-261.
24. Farooq M, Barsa SM, Wahid A. Priming of field-sown rice seed enhances germination, seedling establishment, allometry and yield. Plant Growth Regulation. 2006;49(2-3):285-294.
25. Zheng HC. Effect of seed priming with mixed- salt solution on germination and physiological characteristics of seedling in rice (*Oryza sativa* L.) under stress conditions. Journal of Zhejiang University. (Agric Life Sci). 2002;28:175-178.
26. Khaing M, Ultra VU, Chul Lee S. Seed priming influence on growth, yield, and

- grain biochemical composition of two wheat cultivars. *Journal of Agricultural Science and Technology*. 2020;22(3):875-888.
27. Aryal K, Shrestha A, Subedi R. Effect of various seed priming methods on germination characteristics of black gram. *J. of Protein Research and Bioinformatics*; 2020.  
DOI: 10.24966PRB-1545100009
  28. Zareh M, Mehrabioladi A, Sharafzade S. Evolution effects gibberellic acid (GA<sub>3</sub>) and IAI on germination and seedling growth of wheat under salinity. *Journal of Agricultural Science*. 2006;4:855-865.
  29. Debbarma A, Devi J, Barua M. Seed priming durations and concentrations influence on germination and seedling growth of bitter gourd. *Vegetable Science*. 2018;45(1):137-139.
  30. Ghassemi-Golezani K, Aliloo AA, Valizadeh M, Moghaddam M. Effects of hydro and osmo-priming on seed germination and field emergence of lentil (*Lens culinaris*). *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*. 2008;36(1): 29-33.
  31. Saini R, Rai PK, Bara BM, Sahu P, Anjer T, Kumar R. Effect of different seed priming treatments and its duration on seedling characters of bitter gourd (*Momordica charantia* L.). *Journal of Pharmacognosy and Phytochemistry*. 2017;6(5):848-850.
  32. Chivasa W, Harris D, Chiduza C, Mashingaidze AB, Nyamudeza P. Determination of optimum on-farm seed priming time for maize (*Zea mays* L.) and sorghum (*Sorghum bicolor* L.) for Use to improve stand establishment in semi-arid agriculture. *Tanzania Journal of Agricultural Sciences*. 2000;3(2):103-112.
  33. Doley D, Barua M, Sarma D, Barua PK. Screening and enhancement of anaerobic germination of rice genotypes by pre-sowing seed treatments. *Current Science*. 2018;115(6):1185.
  34. Devi K, Barua PK, Barua M. Integrated effect of pre-sowing seed treatment, sowing windows and seasons on seed yield and quality of greengram. *Legume Research*; 2019.  
DOI: 10.18805/LR-4174
  35. Khan K, Gul Ullah Z, Afsar A, Uddin I, Ullah H. Effect of different osmopriming sources and levels on germination and root length of sorghum. *Weekly Science Research Journal*. 2014;2(4): 5255-5264.
  36. Ruttanaruangboworn A, Chanprasert W, Tobunluepop P, Onwimol D. Effect of seed priming with different concentrations of potassium nitrate on the pattern of seed imbibition and germination of rice (*Oryza sativa* L.). *Journal of Integrative Agriculture*. 2017;16(3):605-613.

© 2023 Bora et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/107251>