



# Influence of Organic Manure and Inorganic Fertilizers on Growth and Yield of Radish (*Raphanus sativus* L.) in Gird Region of Madhya Pradesh, India

Sumit Kumar <sup>a</sup>, Pushpendra Kumar <sup>a\*</sup>, Vinod Jatav <sup>a</sup>,  
Pinkey Dukpa <sup>a</sup> and Shekhar Suman <sup>a</sup>

<sup>a</sup> Department of Horticulture, School of Agriculture, ITM University, Gwalior, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i203804

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106251>

Original Research Article

Received: 10/07/2023  
Accepted: 13/09/2023  
Published: 19/09/2023

## ABSTRACT

A field experiment was conducted to study the effect of organic manure and inorganic fertilizers on growth and yield of Radish (*Raphanus sativus* L.) in Grid region of Madhya Pradesh as factorial based on the randomized block design(RBD) with three replication and Ten treatments viz., T<sub>1</sub>- (100% NPK through inorganic sources), T<sub>2</sub>-(100% NPK through Farm Yard manure (16t/ha)), T<sub>3</sub>- (100% NPK through Vermicompost (12t/ha)), T<sub>4</sub>-(100% NPK through Poultry Manure (9.48t/ha)), T<sub>5</sub>-(75% NPK+25%N through FYM), T<sub>6</sub>-(75% NPK+25% N through Vermicompost), T<sub>7</sub>-(75% NPK+25% N through Poultry manure), T<sub>8</sub>-(50% NPK+50% N through FYM), T<sub>9</sub>-(50% NPK+50% N through Vermicompost), T<sub>10</sub>-(50% NPK+50% N through Poultry manure) were made up of

\*Corresponding author: E-mail: [pkgoyal4699@gmail.com](mailto:pkgoyal4699@gmail.com);

singlevariety Ivory white the experiment was planted in the CRC-II, School of agriculture, ITM University Gwalior. At all of the periodic crop growth phases except at 15 DAS, T<sub>6</sub> showed significant plant height and leaf count increases across crop phases. T<sub>5</sub>, T<sub>6</sub> had maximum heights; T<sub>3</sub>, T<sub>10</sub> had minimum. T<sub>5</sub>, T<sub>6</sub>, T<sub>8</sub> displayed the highest leaf counts; T<sub>1</sub> the least. Leaf lengths followed a similar trend, with T<sub>8</sub>, T<sub>6</sub> having the longest; T<sub>1</sub> the shortest. Overall, T<sub>6</sub> exhibited prominent growth in both height and leaf attributes, while T<sub>1</sub> consistently showed the lowest values throughout all stages and measurements. The conventional practices T<sub>6</sub> had significant positive influence on yield and yield attributing parameters, viz; root length (35.05cm), weight of whole plant (418.91g), initial fresh weight of roots (355.54g) but initial fresh weight of leaves (72.52g) in recorded in treatment T<sub>2</sub>, root diameter (50.61mm), girth of root (12.3cm), dry weight of leaves (13.49g) measured in T<sub>2</sub>, dry weight of root (102.19g), root yield per plot (21.33kg), root yield per hectare (319.98qha<sup>-1</sup>). Conclusion:- Economically the application of 75% NPK+25% N through vermicompost gave maximum net return (Rs. 262553.80) and BC ratio (3.94).

**Keywords:** Economics; FYM; NPK; poultry manure; vermicompost.

## 1. INTRODUCTION

Vegetables are the most important crops which are grown and consumed worldwide. India is second largest producer of vegetable after china; the share of the total vegetable production is almost 13.82 percent (APEDA). Radish (*Raphanus sativus* L.) 2n=18 is an important root vegetable crop belongs to the family Brassicaceae and originated from Europe. It is called 'Mooli'.

Radish is grown in almost all the states of India with an area of 206 thousand hectare and production 3304 thousand MT (2<sup>nd</sup> Advance estimates for 2021-22 as per PIB data base, 2022). However, as per the final estimated data for 2020-21, the area was 207 thousand hectares with production of 3263 thousand MT (NHB data base, 2022). Haryana is leading producer of Radish. The production of Radish is 550.07 thousand tonnes in Haryana (Agriexchange Apeda, 2021-22). Large scale production of radish in the field is a prevalent practice in several states across India, including Haryana, West Bengal, Punjab, Bihar, Assam and Madhya Pradesh. In Madhya Pradesh specifically, radish cultivation is noteworthy, with approximately 11381.25 hectares of land dedicated to its cultivation, resulting in a substantial production output of 174821.14 tonnes (MPH data base, 2019-20).

Radish holds significant medicinal importance, offering various therapeutic uses. It can be employed to alleviate neurological headaches, address sleeplessness issues, and manage chronic diarrhea. The roots of the radish are particularly beneficial for urinary complaints and piles. Moreover, radish leaves are valuable

resources for large scale protein extraction, while radish seeds have the potential to yield non-drying fatty oil suitable for soap production, illumination and edible purposes as documented study in [1]. Radish is predominantly a cool season vegetable crop, it sown during winter from September to January in northern plains of India; Asiatic types can tolerate higher temperatures than European varieties, so in the mild climate of peninsular India, radish can be grown almost all the year round expect for few months of summer. It is an annual or biennial crop depending upon the type for the purpose it is grown (PCARRD, 2009).

Organic manure like farm yard manure (FYM), poultry manure and Vermicompost should also be used as they also make the soil fertile and give nutrition to plant. FYM helps to improving the physical, chemical and biological properties of soil [2]. In comparison to chemical fertilizers, vermicompost stands out as a superior option for improving soil quality and productivity. Similarly, poultry manure, another organic fertilizer, contains elevated levels of essential nutrients like nitrogen, phosphorus, and potassium when compared to manure from other animals. Moreover, poultry manure contributes to enhanced soil fertility, improved soil aeration, and increased water-holding capacity. These findings emphasize the advantages of using organic alternatives like vermicompost and poultry manure over chemical fertilizers for sustainable and fruitful agriculture as demonstrated by Khatri et al., [3]. Chemical fertilizer deteriorates the quality of produce and are expensive too, leading to reduction in the net profit and returns to the farmers. The integrated nutrient management system approach utilizes a judicious combination of inorganic fertilizer and organic manure in

building soil fertility and to the increase the production potential of the crop (Kumar *et al.*) [4].

## 2. MATERIALS AND METHODS

The experimental site is located in Gwalior's district of Madhya Pradesh which is situated in subtropics at an elevation of 196 m above sea level with coordinates at 26° 21'N latitude and 78° 17' E longitude which represents Indo-gangetic plains region.

Gwalior experiences a distinct pattern of rainfall and temperature throughout the year. The majority of its annual rainfall, approximately 80-90%, occurs during the southwest monsoon season, which typically spans from July to September. Occasional showers from cyclonic rains can also be observed during the winter or late spring months. This means that rainfall is concentrated mainly between July and September, with limited precipitation during the winter and spring seasons. The climate in Gwalior exhibits significant temperature fluctuations during both summer and winter. Summers are characterized by scorching heat, with mean maximum temperatures reaching as high as 48°C. Hot and desiccating winds are common during this period. In contrast, winters are quite cold, with mean minimum temperatures dropping to 0°C or even lower. Frost is a regular occurrence during the winter months.

The experiment was carried out using a randomised block design (RBD Factorial). Ten treatments viz., T<sub>1</sub>-(100 % NPK through inorganic sources), T<sub>2</sub>-(100% NPK through Farm Yard manure (16t/ha)), T<sub>3</sub>-(100% NPK through Vermicompost (12t/ha)), T<sub>4</sub>-(100% NPK through Poultry Manure (9.48t/ha)), T<sub>5</sub>-(75% NPK+25%N through FYM), T<sub>6</sub>-(75% NPK+25% N through Vermicompost), T<sub>7</sub>-(75% NPK+25% N through Poultry manure), T<sub>8</sub>-(50% NPK+50% N through FYM), T<sub>9</sub>-(50% NPK+50% N through Vermicompost), T<sub>10</sub>-(50% NPK+50% N through Poultry manure) with three (3) replications, were made up of single variety Ivory white to determine the influence of organic manure and inorganic fertilizers on growth and yield attributes for increased output of radish. On November 28, 2022, the experiment was planted in the CRC-II. In the experiment, NPK (nitrogen, phosphorous, and potash) was applied using urea, DAP, and MOP in accordance with specific treatment protocols. The basal dressing involved the application of the entire phosphorous and potash

quantity, along with one-third of the total nitrogen, as per the respective treatment specifications. The remaining nitrogen was administered as top dressing at two different stages: 15 days after sowing (DAS) and 30 DAS. Each treatment was allocated a specific plot area, and within each plot, five randomly selected plants were identified and tagged. These tagged plants were used for the purpose of collecting various measurements and data throughout the course of the experiment. The data was analyzed using the statistically method outlined by Panse and Sukhatme, (1989).

## 3. RESULTS AND DISCUSSION

### 3.1 Growth Components

The results summarized in Table 1 reveal treatment T<sub>6</sub>-(75% NPK+25% N through vermicompost) as the consistent leader in promoting plant height growth across all stages, showcasing its efficacy with a remarkable 28.6 cm height at harvest. However, the growth journey is far from monotonous, with treatments like T<sub>4</sub>-(100% NPK through Poultry Manure (9.48t/ha)) and T<sub>5</sub>-(75% NPK +25%N through FYM) demonstrating competitive potential at specific stages. The variability in growth patterns underlines the diverse interactions between treatments and growth factors. The close findings are supported by Kumar *et al.*, [4] in radish, Basir kutawa *et al.*, (2020) in okra, Kushwah *et al.*, [5] in carrot and Aswathi P *et al.*, [6]. Application of different sources of nutrients resulted in significant variation for number of leaves per plant at all growth stages. Highest number of leaves was recorded with T<sub>6</sub>-(75% NPK+25% N through vermicompost) at 30 and 45 days while T<sub>8</sub> at harvest followed by T<sub>3</sub>-(100 % NPK through Vermicompost (12t/ha)), T<sub>8</sub>-(50% NPK+50% N through FYM), T<sub>6</sub>-(75% NPK+25% N through Vermicompost at 30, 45 DAS and at harvest) while lowest number of leaves was noted with T<sub>1</sub>-(100% NPK through inorganic sources). Treatment T<sub>6</sub>-(75% NPK+25% N through vermicompost) excels with the highest leaf count due to gradual nutrient release from vermicompost. This enriches soil nutrients, yielding more leaves per plant. The study highlights vermicompost's role in sustained nutrient supply, enhancing leaf growth under T<sub>6</sub>-(75% NPK+25% N through Vermicompost) at 30 and 45 DAS while T<sub>8</sub>-(50% NPK+50% N through FYM) at 15 DAS and at harvest. Similar findings have been reported by Bhattarai and Maharjan, [7] Kumar *et al.*, [4] and Khalid *et al.*, [8] in

**Table 1. Effect of organic manure and inorganic fertilizers on growth parameters**

Treatment	Plant Height(cm)				Number of leaves per plant				Length of Leaves per plant(cm)			
	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest
T1	10.87	18.54	20.83	23.21	3.07	5.07	10.47	15.00	7.45	12.46	17.33	20.79
T2	11.58	17.83	22.61	23.07	4.00	6.87	15.60	19.33	11.18	17.42	21.53	22.66
T3	9.94	18.97	23.23	23.84	4.67	7.73	16.33	18.67	10.21	18.57	22.15	23.44
T4	12.09	19.39	23.91	25.01	4.53	6.93	14.97	15.67	10.35	18.98	22.49	23.61
T5	13.59	19.51	24.23	27.39	5.20	7.53	15.20	17.73	10.51	19.11	22.83	23.98
T6	11.10	20.57	27.13	28.67	4.87	7.87	17.60	19.73	11.23	20.17	24.06	26.27
T7	11.45	18.24	22.82	23.26	4.20	7.47	15.73	18.53	11.35	17.84	21.55	22.86
T8	11.43	20.03	22.43	24.58	4.97	6.80	16.73	20.47	11.75	19.63	23.70	26.93
T9	10.82	19.67	24.17	24.63	4.40	7.47	14.80	15.80	10.55	20.61	23.10	24.89
T10	11.11	16.71	22.39	22.26	3.60	6.33	15.60	17.80	10.70	16.29	20.65	21.86
SE(m)±	0.55	1.44	0.77	0.93	0.42	0.48	1.21	1.15	0.60	1.34	1.22	1.17
C.D. at 5%	1.66	NS	2.29	2.78	1.25	1.43	3.61	3.45	1.81	4.00	3.65	3.49

radish. Among treatments, highest length of leaves was recorded with T<sub>8</sub>-(50% NPK+50% N through FYM) 15 DAS and at harvest followed by T<sub>6</sub>-(75% NPK+25% N through Vermicompost) at 45 DAS and at harvest, while minimum length of leaves was recorded with T<sub>1</sub>-(100 % NPK through inorganic sources). Similar findings have been reported by Kumar et al., (2014) in radish [4].

### 3.2 Yield and Yield Components

The data presented clearly in Table 3. Maximum length of root (35.05 cm) was recorded under the treatment T<sub>6</sub>-(75% NPK+25% N through Vermicompost) and it was followed by (33.55 cm) T<sub>9</sub>-(50% NPK+50% N through Vermicompost) and minimum length of root (28.61 cm) recorded in T<sub>7</sub>-(75% NPK+25% N through Poultry manure). Treatments recorded significant effect on diameter of root. Maximum diameter of root (50.61 mm) recorded under the treatment T<sub>6</sub>-(75% NPK+25% N through Vermicompost), it was (46.05 mm) followed by T<sub>9</sub>-(50% NPK+50%

N through Vermicompost) and minimum (35.47mm) in T<sub>10</sub>-(50% NPK+50% N through Poultry manure). These findings are in agreement with reported by Uddain et al., (2010), [9] Kumar et al. (2014) [16] in radish and Kumar et al. (2014) [4] in carrot.

The Table 2 presented that treatments indicated significant effect on fresh weight of root. Maximum Initial fresh weight of root (402.21 g) was observed with treatment T<sub>6</sub>-(75% NPK+25% N through Vermicompost) followed by (285.78g) in T<sub>8</sub>-(50% NPK+50% N through FYM) and minimum root weight (180.36g) observed in T<sub>10</sub>-(50% NPK+50% N through Poultry manure). The treatments recorded significant effect on Initial fresh weight of leaves (73.52g) in T<sub>2</sub> followed by (63.37g) in T<sub>6</sub> (75% NPK+25% N through Vermicompost) and minimum fresh weight of leaves (44.52g) was recorded in T<sub>10</sub> (50% NPK + 50% N through Poultry manure). Similar findings related to Kushwah et al. (2016) in radish, Kumar et al. [4] in radish. The data summarized in Table 3.

**Table 2. Effect of organic manure and inorganic fertilizers on fresh weight of plant of Radish**

Treatment	Weight of whole plant(g)	Initial fresh weight of roots(g)	Initial fresh weight of leaves(g)
T1	254.81	204.54	50.27
T2	348.43	274.91	73.52
T3	285.86	234.98	50.88
T4	321.95	259.81	62.13
T5	291.83	241.72	50.11
T6	418.91	355.54	63.37
T7	256.95	210.27	46.68
T8	346.51	285.78	60.73
T9	317.43	264.67	52.76
T10	224.88	180.36	44.52
SE(m)±	13.27	10.24	3.23
C.D. at5%	39.72	30.67	9.66

**Table 3. Effect of organic manure and inorganic fertilizers on root of Radish**

Treatment	Root length(cm)	Root diameter(mm)	Girth of Root(cm)
T1	32.53	41.907	10.47
T2	31.94	42.74	10.27
T3	31.03	41.14	10.07
T4	33.41	43.587	10.25
T5	29.92	42.52	9.9
T6	35.05	50.607	12.3
T7	28.61	39.833	9.56
T8	32.63	40.833	11.38
T9	33.55	46.047	11.69
T10	30.29	35.467	8.33
SE(m)±	1.06	1.62	0.41
C.D. at 5%	3.19	4.87	1.24

the treatment T<sub>6</sub>-(75% Vermicompost (50.61 mm)) exhibit the widest root diameter among all treatments. Treatments T<sub>9</sub>-(46.05 mm), T<sub>4</sub>-(43.59 mm), and T<sub>2</sub>-(42.74 mm) also show relatively wider root diameters while (35.47 mm) recorded lowest in T<sub>10</sub>-(50% NPK+50% N through Poultry manure). The variation in root diameters across treatments emphasizes the role of various factors in shaping the root development patterns of radish crops. Maximum Root girth (12.3 cm) was observed in T<sub>6</sub>-(75% NPK+25% N through Vermicompost) followed by T<sub>9</sub>-(11.69 cm) and minimum Girth of root (8.33 cm) was found in T<sub>10</sub>-(50% NPK+50% N through Poultry manure). Similar findings are agreement with Kumar *et al.* (2009), Uddain *et al.*, [9] Kumar *et al.*, [4] and Kushwah *et al.*, (2016) in radish.

The treatment indicated significant effect on dry weight of root and leaves. The data in Table 4 represents that the treatment T<sub>6</sub>-(75% NPK+25% N through Vermicompost) 102.19g exhibits the highest dry weight of roots among all treatments. Treatments T<sub>8</sub>-(99.24g), T<sub>2</sub>-(96.91g), and (95.99g) in T<sub>9</sub>-(50% NPK+50% N through Vermicompost) also show relatively higher root dry weight while treatment T<sub>10</sub>-(47.29 g) has the lowest and Treatments T<sub>6</sub>-(12.25g), T<sub>8</sub>-(12.04g), and T<sub>7</sub>-(11.29g) also show relatively higher leaf dry weights while treatments T<sub>8</sub>-(7.00 g) and T<sub>5</sub>-(7.87g) have relatively lower dry weight. The close findings are supported by Kumar *et al.*, (2009), Uddain *et al.*, [9] and Kumar *et al.*, [4]. The outcome of the investigation presented in (Table 5). Root yield per plot and per hectare was significantly increased due to use of different nutrient management in radish crop [10-12]. The highest root yield of radish per plot (21.33kg) and per hectare (319.98q/ha) was obtained in T<sub>6</sub>-

(75% NPK+25% N through Vermicompost), while lowest root yield (10.82 kg, 162.3q/ha) in T<sub>10</sub>-(50% NPK+50% N through Poultry manure). The close findings are supported by Kushwah *et al.*, [5] (2016) in radish, Kumar *et al.*, [15] (2014) in radish, Kumar *et al.*, [4] (2014) in carrot, Narayan *et al.* in potato [16].

### 3.3 Economics

The suitability of any agricultural practice for cultivating a crop is determined by its ability to maximize economic returns [13,14]. In a cost analysis, the most economically favorable treatment, T<sub>6</sub>, which involved 75% NPK fertilizer and 25% nitrogen through vermicompost, yielded the highest net profit of Rs. 262,553.8 per hectare. Additionally, this treatment achieved the maximum benefit-to-cost ratio. B: C of 3.9 (Table 6) [17].

**Table 4. Effect of organic manure and inorganic fertilizers on dry weight of root and leaves of Radish**

Treatment	Dry Weight of Root(g)	Dry weight of leaf(g)
T1	90.89	9.91
T2	96.91	13.49
T3	79.93	7.00
T4	93.43	9.85
T5	83.67	7.87
T6	102.19	12.25
T7	76.73	11.29
T8	99.24	12.04
T9	95.99	9.19
T10	47.29	9.26
SE(m)±	3.25	0.46
C.D. at 5%	9.73	1.38

**Table 5. Effect of organic manure and inorganic fertilizers on yield of Radish**

Treatment	Root Yield per plot(kg)	Root yield (q/ha)
T1	12.273	184.087
T2	16.497	247.417
T3	14.1	211.483
T4	15.59	233.83
T5	14.5	217.547
T6	21.333	319.983
T7	12.62	189.24
T8	17.147	257.203
T9	15.88	238.2
T10	10.82	162.327
SE(m)±	0.615	9.218
C.D. at 5%	1.841	27.601

**Table 6. Effect of organic manure and inorganic fertilizers on economics of Radish**

<b>Treatment</b>	<b>Treatment Details</b>	<b>Radish yield q ha-1</b>	<b>Gross return (a) Rs. ha-1</b>	<b>Cost of cultivation (b) Rs. ha-1</b>	<b>Net return (a- b) Rs. ha-1</b>	<b>Benefit: cost ratio [(a/b)]</b>
T1	T1-100 % NPK through inorganic sources	184.09	202495.70	54690.00	147805.70	3.70
T2	T2-100 % NPK through Farm Yard manure (16t/ha)	247.42	272158.70	97640.00	174518.70	2.79
T3	T3-100 % NPK through Vermicompost (12t/ha)	211.48	232631.30	97640.00	134991.30	2.38
T4	T4-100 % NPK through Poultry Manure (9.48t/ha)	233.83	257213.00	79640.00	177573.00	3.23
T5	T5-75% NPK + 25%N through FYM	217.55	239301.70	65427.50	173874.20	3.66
T6	T6-75% NPK +25% N through Vermicompost	319.98	351981.30	89427.50	262553.80	3.94
T7	T7-75% NPK + 25% N through Poultry manure	189.24	208164.00	75927.50	132236.50	2.74
T8	T8-50% NPK + 50% N through FYM	257.20	282923.30	76165.00	206758.30	3.71
T9	T9-50% NPK + 50% N through Vermicompost	238.20	262020.00	76090.00	185930.00	3.44
T10	T10-50% NPK + 50% N through Poultry manure	162.33	178559.70	67090.00	111469.70	2.66

#### 4. CONCLUSION

Based on the current research findings, it can be concluded that treatment T6, which consisted of 75% NPK fertilizer and 25% nitrogen supplied through vermicompost, emerged as the most effective treatment combination for enhancing plant growth, root yield, and achieving the highest benefit-to-cost ratio (B: C) for Radish (*Raphanus sativus* L.) of the Ivory White variety grown in the Gird region of Madhya Pradesh.

#### ACKNOWLEDGEMENT

The author expresses gratitude to the Dean, School of Agriculture at ITM University Gwalior (Madhya Pradesh) for generously providing all the necessary facilities essential for the successful completion of the research.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Politud ERR. Growth and Yield Performance of Radish (*Raphanus sativus* L.) cv 'SNOWWHITE' in Response to Varying Levels of Vermicast Applications. International Journal of Scientific and Research Publications. 2016;6(5):53-57.
2. Mengistu M, Abebe Y, Mekonnen Y, Tolessa T. In vivo and in vitro hypotensive effect of aqueous extract of *Moringa stenopetala*. African Health Sciences. 2012;12(4):545-551.
3. Khatri KB, Ojha RB, Pande KR, Khanal BR. The effects of different sources of organic manures in growth and yield of radish (*Raphanus sativus* L.). International Journal of Applied Sciences and Biotechnology. 2019; 7(1):39-42.
4. Kumar P, Meghwal PR, Painuli DK. Effect of organic and inorganic nutrient sources on soil health and quality of carrot; 2014.
5. Kushwah L, Sharma RK, Kushwah SS, Singh OP. Influence of organic manures and inorganic fertilizers on growth, yield and profitability of radish (*Raphanus sativus* L.). Annals of Plant and Soil Research. 2020;22(1): 14-18.
6. Aswathi P, Prasad VM, Deepanshu VB. Integrated nutrient management in radish (*Raphanus sativus* L.) cv. scarlet red globe; 2021.
7. Bhattarai BP, Maharjan A. Effect of organic nutrient management on the growth and yield of carrot (*Daucus carota* L.) and soil fertility status. Nepalese Journal of Agricultural Sciences. 2013;16.
8. Khalid M, Yadav BK, Yadav MP. Studies on the effect of integrated nutrient management on growth and yield attributes of radish (*Raphanus sativus* L.). Annals of horticulture. 2015; 8(1):81-83.
9. Uddain J, Chowdhury S, Rahman MJ. Efficacy of different organic manures on growth and productivity of radish (*Raphanus sativus* L.). International Journal of Agriculture, Environment and Biotechnology. 2010;3(2):189-193.
10. Horticulture statistic at a glance. apeda.gov.in.
11. Horticulture statistic at a glance 1<sup>st</sup> advance Estimate. 2021-22. Available: <https://agriexchange.apeda.gov.in/>
12. Anonymous Indian Horticulture Database, National Horticulture Board, Gurgaon. 2016;172
13. Anonymous (2021-22) (Horticulture statistic at a glance 2<sup>nd</sup> Advance estimates for 2021-22 as per PIB data base, 2022). [www.nhb.gov.in](http://www.nhb.gov.in)
14. Horticulture statistic at a glance (mp government horticulture data base 2019-20), [www.mphorticulture.gov.in/](http://www.mphorticulture.gov.in/)
15. Sandeep K, Sutanu M, Sanjay K, Singh HD. Efficacy of organic manures on growth and yield of radish (*Raphanus sativus* L.) cv. Japanese White. International Journal of Plant Sciences (Muzaffarnagar). 2014;9(1):57-60.
16. Narayan S, Kanth RH, Narayan R, Khan FA, Saxena A, Hussain T. Effect of planting dates and integrated nutrient management on productivity and profitability of potato (*Solanum tuberosum*) in Kashmir valley. Indian Journal of Agronomy. 2014;59(1):145-150.
17. Muhammad M, Kutawa AB, Adamu M. Influence of NPK fertilizer and poultry



manure on the growth of okra  
(*Abelmoschus esculentus* L. Moench) in  
Northern Sudan Savanna region of

Nigeria. International Journal of  
Horticulture, Agriculture and Food  
Science (IJHAF). 2020;4(6).

---

© 2023 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

*<https://www.sdiarticle5.com/review-history/106251>*