



# Commonly Practiced Induced Breeding Techniques of Carp Species in Three Important Fish Hatcheries of Kurigram District, Bangladesh

Md. Hedayetul islam <sup>a</sup>, Sadia Afrin Kamal <sup>a</sup>,  
Kazi Md. Navid Islam <sup>b</sup>, Kazi Towsif Ahmed <sup>c</sup>  
and Ireen Parvin <sup>a\*</sup>

<sup>a</sup> Department of Fisheries Biology & Genetics, Sher-e-Bangla Agricultural University, Dhaka-1207, Bangladesh.

<sup>b</sup> ACI Animal Health, ACI Limited, Dhaka-1208, Bangladesh.

<sup>c</sup> Department of Fisheries Biology & Genetics, Bangladesh Agricultural University, Mymensingh- 2202, Bangladesh.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJFAR/2023/v25i5706

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109724>

Original Research Article

Received: 15/09/2023

Accepted: 22/11/2023

Published: 29/11/2023

## ABSTRACT

The study was carried out to investigate the induced breeding practices and variations in effective hormone dosage used in induced breeding procedures in three different fish hatcheries of Kurigram District of Bangladesh. The hatcheries mainly produced carp fish seeds, including five native fish species such as Rui (*Labeo rohita*), Catla (*Catla catla*), Mrigal (*Cirrhinus mrigala*), Bata

\*Corresponding author: Email: ireen@sau.edu.bd;

(*Labeo bata*), Punti (*Puntius sarana*) and four exotic fish species namely Silver carp (*Hypophthalmichthys molitrix*), Bighead carp (*Aristichthys nobilis*), Grass carp (*Ctenopharyngodon idella*) and Common carp (*Cyprinus carpio*). Exotic carps were the highest produced fish seeds (38%) in the Kurigram district, followed by major carps (35%), Bata (16%), Punti (6%) and others (5%). Three main types of inducing agents including PG (pituitary gland), HCG (human chorionic gonadotropin) and Ovupin (GnRHa injection) were used for induced spawning. The rate of 1st dose of injection of PG, HCG, and GnRHa varied from 0.5 to 2 mg/kg body weight, 100 to 200 IU/kg body weight, and GnRHa (only one dose) 0.2 to 0.4 ml/kg body weight, respectively. The rate of 2nd dose of injection of PG varied from 2 to 12 mg/kg body weight. The PG is usually primarily used for first and second doses; the ovulation period of different fish species varied between 12 to 72 hours, whereas the hatching rate ranged from 70% to 95%. Due to the variations in inducing agents and doses, different ovulation, fertilization, and hatching rates were observed.

**Keywords:** Artificial spawning; hatchery; brood; carp; hormone; Bangladesh; PG; HCG.

## 1. INTRODUCTION

In the fiscal year of 2019-20 [1] Bangladesh emerged as one of the leading fish-producing nations globally, with a total production of 45.03 lakh MT. The country is gifted with vast aquatic assets with a wide variety of aquatic species. The country's fisheries sector contributes 3.52% to the national GDP and 26.37% to the agricultural GDP, making it capable of being self-sufficient in fish production [1]. Bangladesh ranked third in the world in farmed fish production by the Food and Agriculture Organization in 2022, which is considered an exceptional accomplishment of the fisheries sector. Previously aquaculture practices in Bangladesh started with natural seeds, but due to various factors such as the devastation of habitats and breeding grounds, degradation of ecological balance, overfishing, the building of drainage structures, industrial pollutants, and agrochemicals it has now mostly been replaced by hatchery seed production.

The induced breeding or hypophysation technique has significantly transformed quality fish seed production and opened up vast opportunities for advancing the inland fisheries and aquaculture sector. Since 1975, Bangladesh has successfully embraced induced breeding techniques and cost-effective hatchery designs. Bangladesh has 103 government-run hatcheries and 963 private hatcheries, that highlighting the growing adoption of these techniques [2]. The country has accomplished self-reliance in carp seed production, although quality seeds remain limited in scale [3]. The principle of hypophysation technique involves stimulating ripe broods by pituitary or any other synthetic

hormones to breed in captive conditions. It promotes the timely release of sperm and eggs, enabling seed incubation, hatching, and rearing under secure and weather-independent conditions.

The understanding of artificial breeding holds great significance as it enables the intensive production of certain species under controlled conditions, ensuring a reliable and consistent supply of high-quality fish seeds and pure spawn [4]. Pituitary gland (PG) extract and Human Chorionic Gonadotropic (HCG) hormone are commonly used as inducing agents. However, some other agents like Ovupin, Ovaprim etc. are also used for both Indian major carp and other species.

For successful induced breeding, proper hatchery and broodstock management is essential. But many problems exist in fish hatcheries, such as improper hatchery management, poor broodstock management, adverse selection, undesirable sizes of brood fish, inbreeding depression, lack of technical knowledge, lack of good quality fish seeds, inadequate brood fish, collection of broods from the same stock, water quality problem and inadequate supply of nutritive food for brood fish are the most common. However, day by day, the demand for fry and fingerlings is increasing as more farmers show interest in fish culture. So, there is a huge chance to develop this sector by developing a management system and reliable induced breeding technique. Our study was conducted to learn about the induced breeding technique in Kurigram, the basic strategy of brood fish and hatchery management, and the results according to different management techniques.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

The study was carried out between March and April of 2021 in three different fish hatcheries (Ma

Matsya, M. Hossain, and Ananya Matsya) of kurigram town under Phulbari upazila (Fig. 1). Ma Matsya, M. Hossain and Ananya Matsya hatcheries are located in Chandrokhna, Shimulbari and Barabhita areas, respectively of Kurigram district.



Fig. 1. Observed fish hatcheries and their locations in Kurigram district

### 2.2 Study Design




Fig. 2. Conducted flow chart for the study

## 2.3 Data Collection

Primary data was collected through pre-arranged interviews of the operators and hatchery managers, observing the procedures, and surveying each study area. Additional pertinent secondary data and information were gathered from the District and Upazilla Fisheries Offices of the Department of Fisheries (DoF).

## 2.4 Data Processing and Analysis

The collected data were verified for trustworthiness and precision with the help of hatchery personnel. All the accumulated data were cautiously summarized and examined, and then, for further analysis, several pertinent tables were prepared.

## 3. RESULTS

### 3.1 Brood Stock Collection

Broods were primarily collected from natural sources such as different lakes, rivers, haors, baors, beels, jheels etc. The optimal time for brood collection usually falls between November to March. In this study, the hatcheries' broodstock sources are different (Table 1).

### 3.2 Brood Fish Selection for Induced Breeding

The success of induced breeding depends on adequately selecting the broods based on their peripheral characteristics. The mature adult males can be distinguished from the females by the presence of denticulation on the dorsal surface of the pectoral fin, which feels rough to the touch. On the other hand, the ripe females exhibit a soft and bulging stomach with a swollen pinkish genital opening.

The hatchery managers mainly selected the broods based on their weight and age (Table 2), as well as their physical characteristics including size, coloration and growth rate.

### 3.3 Inducing Agents and Hormonal Dose Selection

In the examined hatcheries, mainly three types of inducing agents: pituitary gland (PG), Human Chorionic Gonadotropin (HCG), and Gonadotropin Releasing Hormone analogues

(GnRHa) were used for induced breeding purposes, which were marketed as Ovupin (100 mg Domperidone + 0.2 mg S-GnRHa). Notably, each hatchery has a different selection of dosages and hormone choices depending on species and specific inducing agents utilized (Fig. 3).

Ma Matsya Hatchery used 0.1 ml/kg as first dose of given solution for all carp species. Second dose is administered after 6 hours of first dose and it varies in Indian Carps (Rui, Mrigal, Puti, Bata, Common Carp) and Exotic Carps (Bighead, Grass carp, Common carp). Indian carps are given solution at either 0.6 ml/kg for female or 0.15 ml/kg for males. Whereas exotic carps are given solution at either 1.2 ml/kg or 0.15 ml/kg for males. Second dose of pituitary gland (PG) is given at 1 mg/kg for females and 1.5 mg/kg for males for every species. The dilution rate is ten individual PG, which is 30 mg, mixed with 3 ml of water (Table 3).

As for M. Hossain hatchery, three types of inducing agents are used: pituitary gland (PG), human chorionic gonadotropin (HCG), and Ovupin. For first dose in females, 0.5 to 2 mg/kg PG or 200 IU/kg HCG is used for both types of carps. Additionally, for second dose in females, 6 to 10 mg/kg PG is administered in Indian carp and 6 to 12 mg/kg in exotic carp. Second dose of HCG level for females is 1200 IU/kg for Indian and exotic carp. As for males, 2 mg/kg PG, 200 IU/kg HCG, or 0.3 l/kg Ovupin is given as second dose (Table 4).

At the Ananya Matsya hatchery, only one dose is given to male carp species, whereas two are given to females. For Indian carp species, 1 to 2 mg/kg PG is applied for males and 1 to 2 mg/kg as first dose, followed by 6 to 8 mg/kg PG as second dose is applied for females. No PG is given to male Indian carp. Furthermore, 150 to 200 IU/kg HCG is used for male exotic carp. As for females, 100 to 150 IU/kg HCG and 8 to 12 mg/kg PG are given as first and second doses, respectively (Table 5).

### 3.4 Hormone administration

In all the observed hatcheries, two places were used to inject hormones into a fish to instigate hypophysation. The first method involved an intraperitoneal injection, which was administrated through the ventral bottom part behind either the

**Table 1. Brood fish source of the hatcheries**

SL No.	Hatchery name	Brood fish source
1.	Ma Matsya Hatchery	Collected from local pond.
2.	M. Hossain Hatchery	Carp- Jamuna and Halda river origin fish. Puntii- China origin.
3.	Ananya Matsya Hatchery	Collected from local pond & river.

**Table 2. Age and weight of selected brood fish**

Type	Common Name	Scientific name	Weight	Age
Native carp	Rui	<i>Labeo rohita</i>	1-4 kg	2+
	Catla	<i>Catla catla</i>	2-6 kg	2+
	Mrigal	<i>Cirrhinus mrigala</i>	1-4 kg	2+
	Bata	<i>Labeo bata</i>	0.2-0.8 kg	1+
	Sarputi	<i>Puntius sarana</i>	0.15-0.5 kg	1+
Exotic carp	Silver carp	<i>Hypophthalmichthys molitrix</i>	2-4 kg	2+
	Grass carp	<i>Ctenopharyngodon idella</i>	3-5 kg	2+
	Common Carp	<i>Cyprinus carpio</i>	2-5 kg	1+
	Big head carp	<i>Aristichthys nobilis</i>	2-4 kg	2+



**Fig. 3. Hormone preparation of PG and Ovupin in M. Hossain hatchery**

**Table 3. Inducing Agents & Hormonal Dose Selection of Ma Matsya Hatchery.**

Species	Ma Matsya Hatchery					
	1 <sup>st</sup> Dose	2 <sup>nd</sup> Dose (6 hour after 1 <sup>st</sup> Dose)				
	Female Given Solution (ml/kg)	Female PG (mg/kg)	Given Solution (ml/kg)	PG (mg/kg)	Male Given Solution (ml/kg)	PG (mg/kg)
Rui	0.1	1	0.6	6	0.15	1.5
Mrigal	0.1	1	0.6	6	0.15	1.5
Puti	0.1	1	0.6	6	0.15	1.5
Bata	0.1	1	0.6	6	0.15	1.5
Common Carp	0.1	1	0.6	6	0.15	1.5
Bighead	0.1	1	1.2	12	0.15	1.5
Grass Carp	0.1	1	1.2	12	0.15	1.5
Silver Carp	0.1	1	1.2	12	0.15	1.5

**Dilution rate: 10 PG (30 mg) mixed with 3 ml water**

**Table 4. Inducing Agents & Hormonal Dose Selection of M. Hossain Hatchery**

M. Hossain Hatchery								
Species	1 <sup>st</sup> Dose		2 <sup>nd</sup> Dose ( 6 hour after 1 <sup>st</sup> Dose)					
	Female		Female			Male		
	PG (mg/kg)	HCG (IU/kg)	PG (mg/kg)	HCG (IU/kg)	OVUPIN (ml/kg)	PG (mg/kg)	HCG (IU/kg)	OVUPIN (ml/kg)
Bata	0.5-2.0	200	6-10	1200	0.6	2	200	0.3
Mrigal	0.5-2.0	200	6-10	1200	0.6	2	200	0.3
Sar Puti	0.5-2.0	200	6-10	1200	0.6	2	200	0.3
Rui	0.5-2.0	200	6-10	1200	0.6	2	200	0.3
Common Carp	0.5-2.0	200	6-10	1200	0.6	2	200	0.3
Silver Carp	1.0-2.0	200	6-12	1200	0.6	2	200	0.3
Bighead	1.0-2.0	200	6-12	1200	0.6	2	200	0.3
PG	PG is used on both 1 <sup>st</sup> and 2 <sup>nd</sup> Dosage.							
HCG	HCG is mainly used for 1 <sup>st</sup> Dose and rarely used as 2 <sup>nd</sup> Dose.							
OVUPIN	OVUPIN is used as 2 <sup>nd</sup> Dosage or one single Dose.							

**Table 5. Inducing Agents & Hormonal Dose Selection of Ananya Matsya Hatchery**

Ananya Matsya Hatchery					
Species	Male		Female		
	One Dose		1 <sup>st</sup> Dose		2 <sup>nd</sup> Dose
	PG (mg/kg)	HCG(IU/kg)	PG (mg/kg)	HCG(IU/kg)	PG (mg/kg)
Rui	1-2		1-2		6-8
Catla	1-2		1-2		6-8
Mrigal	1-2		1-2		6-8
Bata	1-2		1-2		6-8
Sarputi	1-2		1-2		6-8
Common Carp	1-2		1-2		6-8
Silver Carp		150-200		100-150	8-12
Grass carp		150-200		100-150	8-12
Bighead		150-200		100-150	8-12

pelvic fin (Fig. 4) or pectoral fin, and intramuscular injections, typically performed on the dorsal part of the fish above the lateral line. During hypophysation, two dosage levels were commonly utilized: a preparatory dose and a decisive or final dose, usually with a time gap of 6 to 12 hours between the two injections. Generally, the preparatory dose constitutes 10-25% of the total. However, no preliminary doses were necessary for males.

### 3.5 Stripping and Fertilization

At first eggs were collected in a dry plastic bowl by employing gentle pressure near the genital opening of the fish. After the eggs were collected, the milt was squeezed out from the male fish into the same bucket similarly as the eggs (Fig. 5A and 5B). Then a little amount of water was added to harden the eggs and remove

excess sperm. Both eggs and milt were mixed by shaking the bowl. In this study, dry fertilization method was used to fertilized the sperm and eggs of fish. Fertilized eggs were then placed for five minutes for better contact with the sperm.

### 3.6 Incubation and Hatching

For incubation funnel incubators were commonly used in all three hatcheries (Fig. 6). Incubation period depends on species and incubation temperature. According to Aktar et al. [5], incubation period of various warm water fish species varies, ranging from a minimum of 12 to a maximum of 72 hours. Valeta et al. [6] also discovered that the incubation period tends to decrease with higher incubation temperatures. Aeration is essential for the embryonic development of eggs just before the hatching of larvae; they need more oxygen because of

wriggling activities to come out from the shell (Fig. 7). During incubation, egg density in the incubator influences both the hatching rate and embryonic development.

### 3.7 Ovulation Rate

The calculation of ovulation rates was performed using the following formula:

$$\text{Ovulation rate (\%)} = (\text{No. of fish ovulated} / \text{Total number of fish injected}) \times 100 \text{ [7]}$$

The ovulation rate of the hatcheries was 95% for Ma Matsya Hatchery and 90% for both M. Hossain and Ananya Matsya Hatchery, respectively (Fig. 8). Ma Matsya Hatchery used PG at the rate of 7mg/kg body weight for Indian carps and 13mg/kg body weight for exotic carps. On the other hand, M. Hossain hatchery used PG (6.5-12/kg body weight for Indian carps, 7-14mg/kg body weight for exotic carps), HCG (200 IU/kg body weight for 1st dose), or Ovupin (0.6 ml/kg body weight for one single dose). Ananya Matsya Hatchery used PG (7-10 mg/kg body weight) and HCG (100-200 IU/kg body weight for 1<sup>st</sup> dose).

### 3.8 Fertilization Rate

The fertilization rates were calculated as follows:

$$\text{Fertilization rate (\%)} = (\text{No. of fertilized eggs} / \text{Total no. of eggs}) \times 100 \text{ [7]}$$

The highest fertilization rate was observed in M. Hossain Hatchery (90-95%), followed by Ananya Matsysya Hatchery at 85-95%, and the lowest fertilization rate was observed in Ma Matsya Hatchery which is 80-90% (Fig. 9).

### 3.9 Hatching Rate

The formula for the hatching rate is:

$$\text{Hatching rate (\%)} = (\text{No. of eggs hatched} / \text{Total no. of fertilized eggs}) \times 100 \text{ [7]}$$

During the study, the maximum hatching rate was observed in M. Hossain Hatchery, which was 85%-95%, following the hatching rates of Ananya Matsysya Hatchery and Ma Matsya Hatchery, which was 80%-90% and 70%-90%, respectively (Fig. 10).



Fig. 4. Intra peritoneal hormone administration in M. Hossain hatchery.




Fig. 5. Stripping of male and female brood fish in Ma hatchery




**Fig. 6. Funnel incubators in Anannya hatchery**



**Fig. 7. Hatchlings produced in M. Hossain hatchery**




**Fig. 8. Ovulation rate of the hatcheries**



**Fig. 9. Fertilization rate of the hatcheries**





**Fig. 10. Hatching rate of the hatcheries**

#### 4. DISCUSSION

We observed the uppermost ovulation rate at Ma Matsya hatchery when they used 7mg PG/kg body weight for Indian carp and 13mg PG/kg for exotic carp species. Notably, deviations from these dosages of PG levels resulted in a decreased ovulation rate. Chaudhuri [8] found that induced breeding experiments yielded better results when the giver and receiver fishes belonged to the same species. Bhuiyan et al. [9] also discovered that the ovulation rate is highest at 7 mg PG/kg body weight, while higher (9mg PG/kg) or lower doses (5 mg PG/kg) led to a decreased rate. Fontenel [10], Atz and Pickford [11], Das and Khan [12], Chaudhuri [13], Mirza et al. [14], Alam and Bhuiyan [15] also implemented similar practices for induced breeding in various fishes.

On the other hand, Ovupin (0.6 ml/kg body weight) showed the best result in the best fertilization rates. Previously Antara et. al [16], and Chakrabarti [17] also found higher fertilization and hatching rate using ovaprim as inducing agent. Two different hormones for two doses (HCG 100-150 IU/kg body weight +PG 6-12 mg/kg body weight) showed better results than only using PG for two doses (1 mg+ 6-12mg/kg body weight).

Moreover, using ovupin gives the highest hatching rate than others. HCG+PG also gives better results than PG, but not as much as

Ovupin. These findings align with the research of Yeasmin et al. [18] on common carp, where Ovaprim (similar to Ovupin) exhibited better fertilization and hatching rates compared to HCG and PG. A similar result was reported by Indira et al. [19] who observed better ovulation, fertilization, and hatching rates in Indian major carp treated with Ovaprim than those treated with PG. Highest ovulation rate observed in Ma Matsya Hatchery (95%) and in terms of fertilization and hatching rate it was observed in M. Hossain Hatchery (90-95% and 85-95%).

In summary, the study highlights the ovulation, fertilization, and hatching rate of larvae primarily influenced by the hormone doses, brood stock mismanagement, quality of the brood, maturity of eggs, seasonal variation, incubation process, water flow during incubation, quality of hatchery water and handling procedure of the broods. It was also detected that the fish bred successfully under artificial conditions when the appropriate hormonal injections were applied.

#### 5. CONCLUSION

This study was performed to observe the induce breeding technique and the differences in production rate due to the use of different hormones at a different dosage of three important fish hatcheries in Phulbari upazilla of Kurigra district. In present study, we observed ovupin hormone gives the highest hatching rate than the PG and HCG hormone. The differences

observed in ovulation rate, fertilization and hatching rate between three hatcheries during this study due to variation in hormones, their dosages and administration timing. These rates can also vary due to the variations in broodstock management, handling, water source, brood fish quality etc. Fish hatcheries significantly impact inland fish production and aquaculture development. Induced breeding has proven to be a revolutionary technique in the fisheries sector worldwide. To ensure high-quality and abundant fish production, providing financial and material support, facilitating information sharing, and organizing workshops and seminars conducted by DoF and Universities is crucial. Additionally, there is a need to focus on developing hatchery facilities and further enhancing induced breeding techniques to foster the growth and progress of aquaculture in Bangladesh.

## ACKNOWLEDGMENTS

The authors would like to express their most profound respect and gratitude to everyone who helped in successfully carrying out this study.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. DoF. Yearbook of Fisheries Statistics of Bangladesh, 2019-20. Fisheries Resources Survey System (FRSS), Department of Fisheries. Bangladesh: Ministry of Fisheries and Livestock, 2020; 37:141.
2. DoF. Yearbook of Fisheries Statistics of Bangladesh, 2020. Fisheries Resources Survey System (FRSS), Department of Fisheries. Bangladesh: Ministry of Fisheries and Livestock, 2021;37:127.
3. Roy NC. Fish Hatchery Management: Broodstock, Fish Breeding & Genetics. Sylhet Agricultural University, Sylhet-3100, Bangladesh. 2016;241.
4. Mollah MFA, Amin MR, Sarowar MN, Muhammadullah. Induced breeding of the riverine catfish *Rita rita*. J. Bangladesh Agril. Univ. 2016;6(2):361– 366.
5. Aktar N, Islam NM. Induced Breeding Practices of the Fish Hatcheries in the North-Western Region of Bangladesh. J Aquac Mar Biol 2015;2(6):00048. DOI: 10.15406/jamb.2015.02.00048.
6. Valeta JS, Likongwe JS, Kassam D, Maluwa AO. Temperaturedependent egg development rates, hatchability and fry survival rate of Lake Malawi Tilapia (*Chambo*), *Oreochromis karongae* (Pisces: Cichlidae). Int. J. Fish. Aqua. 2013;5(4):55-59.
7. Mosaddequr MR, Hossain MY, Md Istiaque Hossain MI, Provhat SJ, Islam MS, Hossain MB. Induced Breeding of the Stinging Catfish, *Heteropneustes fossilis*: Comparison among Different Inducing Agents. Turkish Journal of Fisheries and Aquatic Sciences. 2013;13: 523-527.
8. Chaudhury H. Experiments on induced spawning of Indian carps with pituitary injections. Indian J. Fish., 1960;7(1):20-48.
9. Bhuiyan AS, Jesmin Akhter, Syeda Mushahida- Al-Noor. Efficacy of Two Inducing Agents, PG and DOM+SGNRH on the Induced Breeding of the Major Carp, *Kalibaus (Labeo calbasu)*. Our Nature, 2013;11(1):17-24.
10. Fontenel O. Injecting pituitary (*hypophyseal*) hormones into fish to induce spawning. Prog. Fish. Cult. 1955;17(2):71-75.
11. Atz JW and Pickford GE. The use of pituitary hormones in fish culture. Endeavour. 1959;18(71):125-129.
12. Das SM and Khan HA. The pituitary and pisciculture in India with an account of the pituitary of some Indian fishes and review of techniques and literature on the subject. Ichthyobiologica 1962;1: 43-58.
13. Chaudhuri H. Uses of hormones in induced spawning of carps. J. Fish. Res. Board. Can. 1976;33:940-947.
14. Mirza ZS, Naik IU and Bhatti MZ. Induced spawning of Indian major carps using carp pituitary homogenate and human chorionic gonadotropin in the Punjab (*Pakistan*). Proc. Pak. Cong. Zool. 1993;12:423-428.
15. Alam MM and Bhuiyan AS. Determination of the optimum PG dose for induced spawning of *Labeo rohita (Hamilton, 1822)*. Univ. j. zool. Rajshahi Univ. 1999;18:103-108.
16. Antara G, Rahman MA, Paul P, Ferdous A, Hemal S, Ali MA. Comparative efficacy of PG and Ovaprim in induced breeding of *Kalibaus (Labeo calbasu)* for conservation and aquaculture. World Applied Science Journal. 2020;38(4):309-316.
17. Chakrabarti Saumen. A concise review report on induced breeding of indian major carps through pituitary extract and

- synthetic hormone analogues. International Journal of Recent Scientific Research. 2020;11(A):40011-40016.
18. Yeasmin SM, Rahman MA, Haq M, Hossain MMM, Rahman MH. Effects on hormone on ovulation, fecundity, fertilization and hatching of common carp (*Cyprinus carpio*). Int. J. anim. fish. Sci (online), 2013;1(1):01-07.
19. Indira T, Damodaran R, Priyadarshini R. Comparative study of synthetic hormones Ovaprim and carp pituitary extract used in induced breeding of Indian major carp. Advanced Biotech. 2013;12(7):49-52.

---

© 2023 Islam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/109724>