

Asian Journal of Biology

Volume 20, Issue 1, Page 32-42, 2024; Article no.AJOB.112447 ISSN: 2456-7124

Perceptions of Cat Rearing, Feeding, and Health Management by Cat Owners in Rajshahi

Md. Niyamat Ullah ^a, Md. Abul Kalam Azad ^b, Md. Royhan Gofur ^{a*} and Md. Jalal Uddin Sarder ^a

Department of Veterinary and Animal Sciences, University of Rajshahi, Bangladesh.
 Institute of Environmental Science, University of Rajshahi, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJOB/2024/v20i1384

Open Peer Review History:

Original Research Article

Received: 05/01/2024 Accepted: 25/01/2024 Published: 07/02/2024

ABSTRACT

Background and Objectives: Perceptions of cat owners may affect the lives of pet animals. The study was conducted to learn about of cat owners as well as their contentment with and motivations for raising cats in Rajshahi districts of Bangladesh.

Materials and Methods: Data were collected by a face-to-face interview with 150 pet owners who came to the recognized pet clinics in Rajshahi city from January to December 2023 and analyzes using chi-square test.

Results: This study showed that cat owners made up 70% by female of the total, while 30% made up by male. The majority of cat owners were young people, between the ages of 21 and 30 (49.3%). Urban residents tend to rear cat (62.0%) and have better educational levels (41.5%). Among the cat owners, 87.7% raised their cats indoors, and 11.3% nurtured them outdoors. Residence and educational attainment of cat owners have a substantial association (p<0.05). The breed of cats and their origins were likewise of great worth (p<0.05). The feeding habits of cats

*Corresponding author: E-mail: royhangm@gmail.com;

Asian J. Biol., vol. 20, no. 1, pp. 32-42, 2024

varied from owner to owner; however, the majority (20.7%) fed their cats cooked meat, cooked fish and pellets. Different factors influenced cat-raising with the majority (30.0%) being hobbies. Impact studies showed that biting had a significant negative impact and that unconditional love had a significant good impact. During the investigation, various cat health issues were noted; alopecia was the most common issue (24.7%). The vaccination and deworming rates for their cat were 54.0% and 52.7%, respectively.

Conclusion: It was concluded that different owners raised cats for different reasons, with a different preference in choice of cat breed, feeding practice, medication approach, and through a fuller understanding of these features, such research will aid in the development of a cat-rearing approach that improves the well-being of both pets and their owners.

Keywords: Cat owner; rearing; feeding; health; Rajshahi District.

1. INTRODUCTION

Pet animals can significantly impact human lives by bringing companionship, support, amusement. There was proof that having a pet as company might enhance psychological wellbeing in people by fostering close emotional connections [1,2]. Every year, large quantities of money were spent, often ineffectively, on therapeutic treatment and preventative medicine for people with a variety of physical and mental health issues. Several studies reported that cats can improve human health status and wellbeing. Pet ownership has been linked to lower levels of depression in owners [3,4]. A companion animal may also serve as a form of social support for its owner [5,6] meaning that having the animal around will likely make the owner's life better overall and may even make it easier for them to handle difficult situations. Some owners have reportedly found emotional support in their cats [7], particularly when they used them as a kind of nonjudgmental confidante [8,9]. Despite the fact that the owner is frequently referred to as the career, companion animals may also be able to fulfill the reciprocal duty of the source of security and protection in an attachment connection [10]. They may also bring happiness and comfort to their owner [11], they may be a source of joy and comfort [12,13,14], and during their absence, they may make their owner feel missed [14].

In Bangladesh, the urban population has grown quickly during the past 40 years [15]. People have a wide variety of opportunities to work in different fields as a result of these rapid urbanizations, which helps them enhance their income [16]. Due to the prevalence of nuclear families in urban life and the activities of the majority of family members, people are becoming more and more interested in raising pets [17]. They commonly raised pets, especially dogs and cats, to combat loneliness [18,19, 20,21].

Control methods like quarantine, lockdown, and social seclusion during the COVID-19 pandemic may impact not only people (like impact on emotion, health and lifestyle) but also their companion animals [22,23]. The chances of COVID-19 transmission from pets to people were extremely unlikely, according to research published shortly after the COVID-19 pandemic [22,24]. However, it's possible that the media caused a cultural reaction that led to thousands of abandoned pets and the euthanasia of many more [23]. However, qualitative insights revealed that cat owners considered their experience of the lockdown as being made easier by having a pet. Oliva and Johnston [25] reported that cat ownership was not a significant predictor of loneliness scores during lockdown. Along with combating loneliness, pets, especially cats, rear for many reasons. includina companionship, loneliness and depression and hobby, animal welfare, depression, anxiety, marital complexity, children's demands, and others. According to reports, pet owners are happier and more satisfied with their lives than non-owners [26]. Additional responsibilities related to proper pet care include housing, feeding, deworming, immunization, grooming, and exercise [17]. The well-known zoonotic disease of rabies can be prevented in cats by proper anti-rabies immunization. Other vaccines those help to prevent illness and strengthen a cat's immune system included a combination of antiviral vaccines against feline parvovirus, feline herpes, and feline infectious rhinotracheitis. A crucial factor in keeping cats worm-free is the regular use of anthelmintic medications. Regular grooming improves a cat's attractiveness and helps to prevent ectoparasites. The same holds true for nutritionally balanced diets, which support protection against several chronic diseases and improve health [27]. There were many commercial cat foods accessible in Bangladesh, but the majority was imported

because there was no such cat food production facility in Bangladesh. Some cat owners solely fed their pet's homemade food or leftovers, while others fed them processed foods, but most owners were unaware of their balanced diet. The purpose of the current study was to ascertain cat owners' socio-economic conditions, their pet rearing technique and perceptions in Rajshahi district in Bangladesh.

2. MATERIALS AND METHODS

2.1 Study Area

The study was conducted in the divisional city of Rajshahi in Bangladesh on 150 cat owners.

2.2 Data Collection

A comprehensive, coordinated questionnaire was developed to gather information from cat owners at three different pet clinics in Rajshahi. The questionnaire includes questions about the pet owner's age, gender, place of residence, level of education, as well as the pet animals' breed, food preference, feeding habits, deworming, vaccinations, and overall health.

Pet owners were questioned face-to-face utilizing a questionnaire on the hospital grounds. Pet owners were surveyed from January 2023 to December 2023, and information was gathered from three pet clinics namely: Kittycat care and cure; Birds Pet Animal Clinic and Dr. Shazid's Pet clinic. The sample size was 150 pet owners who brought their cats to the hospital for treatment or routine health checks were interviewed during this study.

2.3 Data Analysis

Following data collection, data were compiled and analyzed using SPSS version 2021. Significant differences among the categorical variables were calculated using a chi-square test with a significance level set at alpha <0.05.

3. RESULTS

3.1 Social Characteristics of Cat Owners

3.1.1 Demographic status of pet owners

The majority of cat owners (70%) were women, with 30% being men. The age of the respondents ranged from 10 to more than 40 years. The age distribution was as follows: less than 10 years (3.3%), 10 to 15 years (6.7%), 16 to 20 years (15.3%), 21 to 30 years (49.3%), 31 to 40 years (13.3%), and more than 40 years (12.0%). Urban residents make up 62.0% of cat owners, followed by rural residents (16.7%) and suburban residents (21.3%). According to the educational level, there were four levels of education: primary (18.0%), secondary (28.0%), higher (44.7%), and none (9.3%). Indoor and outdoor cat rearing systems make up, respectively, 88.7% and 11.3% of the population (Table 1).

Table 1. Demographic status of pet owners

Characteristics		Frequency (%)	
Sex	Male	45 (30.0%)	
	Female	105 (70.0%)	
	<10	05 (3.3%)	
	10-15	10 (6.7%)	
	16-20	23 (15.3%)	
Age (year)	21-30	74 (49.3%)	
	31-40	20 (13.3%)	
	>40	18 (12.0%)	
	Urban	93 (62.0%)	
Residence	Rural	25 (16.7%)	
	Sub-urban	32 (21.3%)	
	Primary	27 (18.0%)	
	Secondary	42 (28.0%)	
Educational Status	Higher	67 (44.7%)	
	None	14 (9.3%)	
Rearing system of cat	Indoor	133 (88.7%)	
	Outdoor	17 (11.3%)	

3.1.2 Educational status and residence of cat owners

The correlation between educational level and residence of cat owners was greatly significant (p<0.05). Out of 150 cat owners, 93 lived in urban areas where the majority had higher levels of education (Table 2). In contrast, there were 25 cat owners in rural areas who have lower education levels than urban residents. There were 32 respondents in suburban areas where owners' educational levels are higher than those in rural areas (Table 2).

3.2 Technical Characteristics of Pet Cat Owners

3.2.1 Available cat breeds in Bangladesh with their sources

In Bangladesh right now, there are many cat breeds accessible. Due to the high value of foreign cats, raising foreign breeds was done by financially stable individuals. The analysis revealed several cat-producing sources. Cat breeds came from a variety of sources, including gifts, street collections, purchases, and adoptions. The study's findings indicate that native breeds were preferred over imported ones in the majority of cases. The comparison between cat breed and sources was significant at p<0.05 (Table 3).

3.2.2 Feeding management and practices

Cat owners offer different types of food to their pets. They have supplied both homemade foods and some commercial foods (pellets, cans, and pouches). Most cat owners (20.7%) offered their pets cooked meat, cooked fish, and pellets. Usually, three to four times a day were set aside for feeding (Table 4).

3.2.3 Vaccination, deworming and common health issues of cat

Vaccination was a significant concern for cats. The rabies vaccine aids in the prevention of zoonotic illnesses. The combined antiviral vaccine which includes calcivirus, feline rhinotracheitis virus, and feline panleukopenia virus was the most significant vaccination. The administration of a combination antiviral vaccines was increased the cat's immunity. According to the present research, 54.0% of cat owners

vaccinated their cats (Table 5). The cat was kept free of worms by routine deworming. Our research reported that 52.7% of cat owners dewormed their cats (Table 5). Many health issues were monitored while raising cats. Among the common health problems were diarrhea, alopecia, inappetance, vomiting, and indigestion. The most frequent health problem was vomiting 22.70% (Fig. 1).

3.3 Motivation for Pet Cat Rearing

3.3.1 Experience and reasons of cat rearing

Different owners had different experiences with raising cats. A maximum experience of two years (35.30%) of cat-rearing was obtained (Fig. 2). There were some fascinating justifications for raising cats. The common causes included pressures from children, marital difficulty, animal welfare, loneliness, depression, anxiety, and other factors. The most common reason among them was their hobbies (Fig. 3).

3.3.2 Impact analysis of pet management

This study examined the effect of cat ownership on cat owners' mental health. Both positive and negative effects have been observed. Unconditional love dominated the favorable effects (34.7%), whereas biting (21.3%) had the most adverse effects (Table 6).

4. DISCUSSION

A pet is defined as a cat, dog, or companion animal that the subject can interact with [28]. Interacting with animals improves a person's attitudes, behaviors, sense of personality and social support [29]. This study showed that women (70%) are more involved in cat rearing than men (30%).

A study conducted in the United Kingdom and Ireland suggested that women were more likely than men to keep cats and that this link should also be applied to children's ownership of cats and dogs [30,31,32]. This may be because, in Bangladesh, the majority of women are responsible for household duties, so they feel spending their free time with pets. In addition, women may be more interested companionship with pets during pregnancy. In this study, females owned significantly more pets.

Table 2. Relation between educational status and residence of pet owners

Characteristics		Educational status				Chi-Square Tests			
			Primary	Secondary	Higher	None	Total	Value	AsympSig.(2 sided)
	Urban	Count	11	27	49	6	93		
Residence	Ulbali	Expected	16.7	26.0	41.5	8.7	93.0		
Rur Sub	Dimal	Count	5	6	10	4	25	13.563	0.035**
	Rurai	Expected	4.5	7.0	11.2	2.3	25.0		
	Cultural and	Count	11	9	8	4	32		
	Sub-urban	Expected	5.8	9.0	14.3	3.0	32.0		
Total		Count	27	42	67	14	150		
		Expected	27.0	42.0	67.0	14.0	150.0		

*p<0.01

Table 3. Various cat breed and their sources

		Sources of cat					Pearson Chi-Square Test	
Cat breed	Adoption	Buying	Collection from street	Gift	Others	Total	Value	Asymp. Sig. (2 sided)
Deshi	33	1	24	7	5	70		
Deshi+Persian	2	6	0	0	0	8		
Mixed+Persian	0	7	0	0	0	7		
Deshi+ Mixed	6	0	0	1	0	7		
Deshi+Mixed+Persian	4	2	0	1	1	8		
Mixed	5	8	1	0	0	14	400 400	0.000***
Persian	0	15	0	2	0	17	129.498	0.000***
Turkish angora	0	5	0	0	0	5		
British short hair	1	4	0	0	0	5		
Maine coon	0	2	0	1	0	3		
Others	3	0	0	2	1	6		
Total	54	50	25	14	7	150		

***p<0.001

Table 4. Feeding practices of cat

Food items	Frequency (%)	
Pellet (commercial food)	14 (9.3 %)	
Others	01 (0.7%)	
Pellet+ Cooked fish+ Cooked meat	31 (20.7%)*	
Cooked fish+ Cooked Meat	21 (14.0%)	
Rice+ Milk	14 (9.3%)	
Rice+ Fish bone	05 (3.3%)	
Cooked fish+ Cooked meat+ Rice+ Vegetable	08 (5.3%)	
Cooked fish+ Cooked meat+ Rice	05 (3.3%)	
Pellet+ Canned	04 (2.7%)	
Commercial canned food	07 (4.7%)	
Bone (fish)+ Rice	02 (1.3%)	
Bone (fish)+ Biscuit+ Vegetable	06 (4.0%)	
Cooked fish	18 (12.0%)	
Cooked meat	05 (3.3%)	
Rice	04 (2.7%)	
Bone (fish)	05 (3.3%)	
Total	150 (100.0%)	

Table 5. Vaccination and deworming status of cat rearing

Characteristics	Frequency (%)	
Vaccination		
Yes	81 (54.0%)	
No	69 (46.0%)	
Total	150 (100.0%)	
Deworming		
Yes	71 (52.7%)	_
No	17 (47.3%)	
Total	150 (100.0%)	

Table 6. Perceived effects of cat rearing

Characteristics	Frequency (%)	
Positive impact		
Unconditional love	51 (34.7%)	
Home safety	07 (4.7%)	
Nurture	19 (12.7%)	
Health benefit	21 (14.0%)	
Playing with cat	27 (18.0%)	
Spent time	25 (16.7%)	
Total	150 (100.0%)	
Negative impact		
Biting	32 (21.3%)	
Others	08 (5.3%)	
Stool/ Urination	23 (15.3%)	
Living expenses	11 (7.3%)	
Allergies	25 (16.7%)	
Respiratory problem	07 (4.3%)	
Damage house/ property	20 (13.3%)	
Disturbances of sleep	08 (5.3%)	
Loss of food or curry	05 (3.3%)	
Bad odor	11 (7.3%)	
Total	150 (100.0%)	




Fig. 1. Common health issues of cat




Fig. 2. Experience of cat rearing




Fig. 3. Reasons of cat rearing

The present research showed that owners of cats were able to lower their depression levels by receiving the unconditional love of their pets. According to Garrity et al. [33], having a pet has a favorable effect on depression, emotional wellbeing, and physical well-being. Pet ownership has been shown by McConnell et al. [34] to improve well-being and reduce depression in those who have unmet social needs. [29] conducted a research and found that interacting with animals improves a person's attitude, behavior, sense of responsibility, and social support.

Cats are mostly found in indoor environments (80.7%) and outdoors (11.3%). Although it has been demonstrated that domestic cats' home ranges were wider in rural than urban areas, the quantity of outdoor activity relies on the location of the home and the neighborhood [35,36]. Contrarily, research showed that indoor-only pet cats interact with their owners more than those kept outdoors as pet cats [37]. According to those studies, the majority of pet owners were well-educated and financially stable. We also found the same.

According to our results, the majority of cat owners (49.3%) were young people between the ages of 21-30 years old. A previous research by Purewal et al. [38] reported that the younger age group's lower involvement in pet rearing activities may be attributed to a lack of free time and education-related activities.

Contrarily, older people may raise cats as a way to pass the time and show them unconditional love. Most cat owners in our study were rear deshi cats, and their primary source of cats was adoption. Fewer individuals keep exotic breeds (such as Persian, Maine Coon, Russian Blue, and Turkish A ngoora), and their primary source was purchasing them because of their financial value. Persian breeds dominated among foreign breeds due to their availability in Rajshahi, Bangladesh. In Rajshahi, several people offer Persian breeds commercially [39].

In order to provide nutritious, high-quality food for their pets, pet owners should be aware of quality meals and consult with vets. The majority of dogs and cats were being fed commercial pet food, but many poor participants also said they provided homemade recipes to their pets [17]. In addition, pet owners may use homemade feeding regimens as a means of providing affection or strengthening their bond with their animals. They

may have found these feeds to be more palatable or believed that they were healthier than commercially prepared diets [40]. However, this was at odds with other experiments conducted by Baldwin et al. [40] and Freeman et al. [41], which proposed that the food available on the market is a nutrient-dense, well-balanced, easily prepared diet. In our study, the majority of cat owners offer their cat's homemade food in addition to a few commercial foods (cooked fish, cooked meat, and pellets). But if homemade food is not prepared properly, that food will cause several health problems for cats. Some owners directly give raw fish and meat to their cats, which cause dyspepsia and a higher risk of worm infestation. Cats who consume spicy food may develop a variety of skin disorders, such as dermatitis and alopecia were reported by the pet owners during the interviewed in present study.

A number of health problems were claimed by the pet owners during the interview, and the most common which was alopecia Ωf (24.70%) and vomiting (22.70%) in the present study. Similar results also found in the UK, Japan, and Sweden, dermatological illnesses were among the most common disease categories [42,43,44]. Numerous health effects were discovered in the study, with the most notable good impact being unconditional love (37.7%) and the most notable negative effect being the cat's propensity for biting (21.3%). According to the survey, 52.7% of cats were dewormed, and 54.0% of cats received vaccinations. There was a greater risk to owners if the cat was not vaccinated against rabies. A combined antiviral vaccine is also required for cat immunity building. Deworming was essential for maintaining worm-free conditions and preventing zoonotic worm infestations. Public education was crucial for reducing the risk of parasitic zoonosis in both humans and companion animals [45,46,47].

5. CONCLUSIONS AND RECOMMANDA-TIONS

A big number of cats were being raised indoors on a daily basis. Cat owners perceived both positive and negative impacts. However, urban people with higher education were more inclined to raise the cat. Moreover, the majority of owners were unaware of zoonotic diseases. Therefore, regularly immunizing and deworming their cats and counseling cat owners about health issues and zoonotic diseases of cats were crucial. It is recommended that Rajshahi district residents

have access to contemporary medical facilities for the treatment of their pets.

ACKNOWLEDGEMENTS

Authors would like to thank the Institute of Environmental Science, University of Rajshahi, Bangladesh for financial support to conduct the research. Authors also thank to the owners of pet clinics (Kittycat Care and Cure, Birds and Pet Animal Clinic, Dr. Shazids Pet Clinic), and pet owners for helping in data collection.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Ståhl A, Salonen M, Hakanen E, Mikkola S, Sulkama S, Lahti J, Lohi H. Pet and owner personality and mental wellbeing associate with attachment to cats and dogs. IScience. 2023;26(12):108423. DOI: 10.1016/j.isci.2023.108423
- 2. Wells D LJ. The effects of animals on human health and well-being. Journal of Social Issues. 2009;65(3):523–543.
- 3. Clark Cline KM. Psychological effects of dog ownership: Role strain, role enhancement, and depression. The Journal of Social Psychology 2010;150:117–131.
- 4. Chakma SK, Islam TT, Shahjalal M, Mitra DK. Depression among pet owners and non-pet owners: a comparative cross-sectional study in Dhaka, Bangladesh. F1000Research. 2022;10: 574.
- 5. Staats S, Wallace H, Anderson T. Reasons for companion animal guardianship (pet ownership) from two populations. Society and Animal. 2008;16:279–291.
- 6. Beetz A, Julius H, Turner D, Kotrschal K. Effects of social support by a dog on stress modulation in male children with insecure attachment. Frontiers Psychology. 2012;3:352.
- 7. Stammbach KB, Turner DC. Understanding the human-cat relationship: Human social support or attachment. Anthrozoos. 1999; 12:162–168.
- 8. Hafen M, Rush BR, Reisbig AM, McDaniel KZ. The role of family therapists in veterinary medicine: Opportunities for

- clinical services, education, and research. Journal of Marital Family Therapy. 2007:33:165–176.
- Pachana NA, Massavelli BM, Robleda-Gomez S. A developmental psychological perspective on the human–animal bond. In the psychology of the Human-Animal Bond; Blazina, C, Boyraz G., Shen-Miller, D., Eds; Springer: New York, NY, USA. 2011;151–165.
- 10. Rockett B, Carr S. Animals and Attachment Theory. Society and Animal 2014;22:415–433.
- 11. Kurdek LA. Pet dogs as attachment figures for adult owners. Journal of Family. Psychology. 2009;23:439–446.
- 12. Prato-Previde E, Fallani G, Valsecchi P. Gender differences in owners interacting with pet dogs. An Observational Study Ethology. 2006;112:64–73.
- 13. Kurdek LA. Pet dogs as attachment figures. Journal of Social and Personal Relationship. 2008;25: 247–266.
- 14. Walsh F. The Relational significance of companion animals. Human-animal bonds. Family Process. 2009;48:462–480.
- Helal M, Hossain MA. Four decades of economic development of Bangladesh: An assessment. Journal of the Asiatic Society of Bangladesh (Hum.). 2013;58:335–362.
- 16. Rana MMP. Urbanization and sustainability: challenges and strategies for sustainable urban development in Bangladesh. Environment, Development and Sustainability. 2011;13:237–256.
- Bhowmik P, Mimi H K, Datta A, Adhikary K, Akter N, Barua K, Hossain A K. Food, nutrition and health status of the pet animals in Dhaka and Chattogram city of Bangladesh. Bangladesh Journal of Veterinary and Animal Sciences. 2020; 8:171-179.
- Kuzniar AA. Melancholia's dog: Reflections on our animal kinship. University of Chicago Press, USA;2006.
- 19. Beck A. Between Pets and People: The Importance of Animal Companionship. Purdue University Press, India;2011.
- Irvine L. Animals as life changers and lifesavers: Pets in the redemption narratives of homeless people. Journal of Contemporary Ethnography. 2013;42:3– 30.
- 21. Bradshaw J. The animals among us: The new science of anthrozoology. Penguin UK:2017.

- 22. Bowen J, Garcia E, Darder P, Argűelles J, Fatjo J. The effects of the Spanish COVID-19 lockdown on people their pets and the human-animal bond. Journal of Vet Behavior. 2020;40:75–91.

 Available:https://doi.org/10.1016/j.jveb.2020.05.013
 - PMID: 32837452.
- 23. Parry NMA. COVID-19 and pets. When pandemic meets panic. Forensic Science International Reports. 2020;2:100090.
- Goumenou M, Spandidos DA, Tsatsakis A. Possibility of transmission through dogs being a contributing factor to the extreme Covid-19 outbreak in North Italy. Molecular Medicine Reports. 2020;21:2293–2295. Available:https://doi.org/10.3892/mmr.2020.11037
 - PMID: 32236594
- Oliva JL, Johnston KL. Puppy love in the time of Corona: Dog ownership protects against loneliness for those living alone during the COVID-19 lockdown. International Journal of Social Psychiatry. 2021;67(3):232-242.
 - DOI: 10.1177/0020764020944195
- 26. Bao KJ, Schreer G. Pets and happiness: Examining the association between pet ownership and wellbeing. Anthrozoos. 2016;29:283–296.
- 27. Case LP, Carey DP, Hirakawa DA. Canine and Feline Nutrition: A resource for companion animal professionals. mosby inc; 2nd edition; 1995.
- 28. Parslow RA, Jorm AF, Christensen H, Rodgers B, Jacomb P. Pet ownership and health in older adults: findings from a survey of 2.551 community-based australians aged 60-64. Gerontology. 2004;51:40-47.
- 29. Serpell J. Guest editor's introduction: Animals in children's lives. Society and Animals 1999;7(2): 87–94.
- 30. Downes M, Canty MJ, More SJ. Demography of the pet dog and cat population on the island of Ireland and human factors influencing pet ownership. Preventive Veterinary Medicine. 2009;92: 140–149.
- Murray JK, Browne WJ, Roberts MA, Whitmarsh A, Gruffydd-Jones TJ. Number and ownership profiles of cats and dogs in the UK. Veterinary Record. 2010;166:163– 168.
- 32. Westgarth C, Pinchbeck GL, Bradshaw JWS, Dawson S, Gaskell RM, Christley RM. Factors associated with cat ownership

- in a community in the UK. Veterinary Record. 2010:166:354–357.
- 33. Garrity TF, Stallones LF, Marx M B. Pet ownership and attachment as supportive factors in the health of the elderly. Anthrozoös. 1989;3(1):35–44. DOI:10.2752/089279390787057829
- McConnell AR, Brown CM, Shoda TM, Stayton LE, Martin CE. Friends with benefits: on the positive consequences of pet ownership. Journal of Personality and Social Psychology. 2011;101(6):1239-52. DOI: 10.1037/a0024506
- 35. Metsers EM, Seddon PJ, Van Heezik YM. Cat-Exclusion zones in rural and urban-fringe landscapes: How large would they have to be? Journal of Wildlife Research. 2010;37:47–56. DOI: 10.1071/WR09070
- 36. Hall CM, Bryant KA, Haskard K, Major T, Bruce S, Calver MC. Factors Determining the Home Ranges of Pet Cats: A Meta-Analysis. Biological Conservation 2016;203:313–320. DOI: 10.1016/j.biocon.2016.09.029
- 37. Hubrecht RC, Turner DC. Companion animal welfare in private and institutional settings. In: Wilson CC, Turner DC. editors. Companion animals in human health. Sage publications inc. Thousand Oaks. CA. USA: pp. 1998:267–289.
- 38. Purewal R, Christley R, Kordas K, Joinson C, Meints K, Gee N, and Westgarth C. Companion animals and child/adolescent development: A systematic review of the evidence. International Journal of Environmental Research and Public Health. 2017:14: 234.
- 39. Remillard RL. Homemade diets: attributes, pitfalls, and a call for action. Topics in Companion Animal Medicine. 2008;23:137–142.
- Baldwin K, Bartges J, Buffington T, Freeman L M, Grabow M, Legred J, and Ostwald Jr D. AAHA nutritional assessment guidelines for dogs and cats. Journal of the American Animal Hospital Association. 2010;46:285–296.
- 41. Freeman L, Becvarova I, Cave N, MacKay C, Nguyen P, Rama B, Takashima G, Tiffin R, van Beukelen P, Yathiraj S. WSAVA nutritional assessment guidelines. Journal of Feline Medicine and Surgery. 2011; 13:516–525.
- 42. Neill DG, Church D, Mc Greevy P, Thomson P, Brodbelt DC. Prevalence of disorders recorded in cats attending

- primary-care veterinary practices in England. Veterinary Journal. 2014;202: 286–91.
- DOI: 10.1016/j.tvjl.2014.08.004
- Inoue M, Hasegava A, Sugiura K. Morbidity pattern by age, sex, and breed in insured cats in Japan (2008-2013). Journal of Feline Medicine and Surgery. 2016; 18(12):1013-1022.
 DOI: 10.1177/1098612X15616433
- 44. Egenvall A, Bonnett BN, Häggström J, Ström HB, Möller L, Nødtvedt A. Morbidity of insured Swedish cats during 1999-2006 by age, breed, sex, and diagnosis. Journal of Feline Medicine and Surgery. 2010;12:948–59.

- DOI:10.1016/j.jfms.2010.08.008
- 45. Baneth G, Thamsborg S M, Otranto D, Guillot J, Blaga R, Deplazes P, Solano-Gallego L. Major parasitic zoonoses associated with dogs and cats in Europe. Journal of Comparative Pathology. 2016; 155(1 Suppl 1):S54-74.
 - DOI: 10.1016/j.jcpa.2015.10.179
- 46. Traversa D. Pet roundworms and hookworms: a continuing need for global warming. Parasites and Vectors. 2012; 5:91.
- 47. Uribe Corrales N. The significance of education in the preparedness for zoonotic diseases. IntechOpen;2023.

 DOI: 10.5772/intechopen.1002409

© 2024 Ullah et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
https://www.sdiarticle5.com/review-history/112447