



Under Environmental Stress Increase CO₂ Level and the Photosynthetic Response of Plant

Ayesha Arif ^a, Sadia Batool ^a, Alveena Rafiq ^{a*}, Iaraib barira ^a, Zubaira BiBi ^a, Zabeehullah Burhan ^a and Urooj Bashir ^a

^a *Department of Botany, University of Agriculture, Faisalabad, Pakistan.*

Authors' contributions

This work was carried out in collaboration among all authors. Author AR was responsible for creating the study and writing the protocol. Authors AR and AA handled the preparation of the materials, data collection, and analysis. Author AR wrote the first draft of the manuscript, and author SB provided feedback on earlier iterations. Authors Ib, ZB and Zb the literature searches and contributed a lot to Strategies Portion. The final part of the manuscript is hinder hunger written by authors UB and ZB. Author AR was in charge of managing the references and citations. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115199>

Review Article

Received: 23/01/2024

Accepted: 27/03/2024

Published: 01/04/2024

ABSTRACT

Although the importance of respiration and photosynthesis to plants is well-established, the antioxidant system's response to abiotic stresses remains an area of intense interest in the study of physiological stress. While reports and reviews have been conducted on a single important metabolic process and its reaction to climate change, there has been little coverage of an integrated study that would include several biological processes at different scales. Along with other important abiotic stresses like drought, heat, nitrogen limitation, and ozone pollution, this review will

*Corresponding author: E-mail: alveenarafiq2001@gmail.com;

provide a synthesis of the mechanisms to elevated CO₂ and its responses at various scales, including cellular, molecular, physiological, and biochemical, and individual aspects. While it contains what has been well-established in earlier reviews, the current comprehensive evaluation may contribute considerable and pertinent information about the issue in recent research. An introduction to the essential biological processes and a synopsis of their functions in controlling the environment follows. The second part of the article discusses the current state of study on the many subtopics, such as how plants adjust their antioxidant system, respiration, and photosynthetic capacity to either CO₂ enrichment or other forms of climate change. In the end, we go over some of the possible uses for plant responses to different degrees of climate change. aided by this review, which is currently of paramount concern on a global scale.

Keywords: Antioxidant; abiotic stresses; bio-chemical processes; comprehensive evaluation; of physiological stress.

1. INTRODUCTION

Increases in atmospheric CO₂, global warming, changes in precipitation patterns, and the feedback loops between these and other environmental variables are the main drivers of climate change [1]. Due to the significant rise in CO₂ emissions to the atmosphere this century, its level has been steadily rising at a rate of around 1.0 μmol mol⁻¹ per year [1,2].

Our attention here is directed to the essential plant biological processes involved in climate change, such as respiration, photosynthesis, metabolic activities, and the antioxidant system. Both respiration and photosynthesis are essential to plant life. Photosynthesis fixes carbon dioxide and releases oxygen into the air, while respiration produces energy and works on substrate metabolisms like those that supply the carbon skeleton.

According to many studies Singh & Agrawal, [3] Long et al., 2004; Peñuelas et al., 2013; Matros et al., [4] the respiration, photosynthesis, and antioxidant systems, among other important secondary metabolisms in plants, have been impacted by the increasing levels of CO₂. These basic biological reactions may be the basis for all additional impacts of increased CO₂ on specific plants and ecosystems. Increased CO₂ levels may affect genetic variations linked to biological process traits. For the reasons already mentioned, studying how plants adapt to a changing environment has been a major focus of botanical studies all sizes in the last few decades. Photosynthesis, stomatal behavior, and other biological reactions to elevated CO₂ levels and how it interacts with changes in the environment have been extensively studied in several papers (Long et al., 2004).

Plant development, N and carbon (C) allocations, and gas exchange reactions to increased CO₂ in the context of drought and high temperatures were the subjects of our previous review by Xu et al. (2013a). Whether we're talking about the molecular, cellular, biochemical, physiological, organ, person, or ecosystem level, this review did touch on water use efficiency (WUE), how higher plants' growth, photosynthesis, and changed when contact with CO₂ upgrading with abiotic variables (Ainsworth and Long, 2005).

However, it did not go into detail on the numerous underlying mechanisms by which these critical biological processes were affected, modulated, or controlled. To be more specific, there has been no well evaluated systematic evaluation of these materials. Consequently, we aimed to methodically summarize the substantial study found that when exposed to increased CO₂ levels together with other environmental factors, the respiratory, photosynthesisal, and antioxidant systems reacted differently. Our review was in light of the need to update and consolidate all relevant studies on plant biology and climate change (Ainsworth Rogers, 2007).

In particular, we examined the processes at work, the paths for a response, and the linkages between these. Lastly, we provide and discuss briefly the future prospects of our work in relation to the potential consequences. As a result, this review may be relevant to the present day due to its multidisciplinary and systematic synthesis, which includes detailed information on significant past and present experimental findings, theoretical analysis, underlying processes, and possible submissions that could spur additional research.

2. RESPONSES THE PROCESSES OF CRITICAL BIOLOGICAL TO ELEVATED CO₂

2.1 Stomatal Conductance Response to Elevated CO₂ Concentration

Guard cells are innately able to sense CO₂, and it is believed that they react more to CO₂ in the intercellular space (ci) than to CO₂ at the surface of the leaf. We will just provide a brief overview of guard cell signaling and metabolism. The amounts of organic and ionic solutes facilitate the turgor pressure inside the guard cells, which controls the stomatal opening. The potential of guard cell membrane must be depolarized in order for the stomata to close. Based on electrophysiological studies, it was found that when CO₂ levels are high, certain potassium channels are more active than others. On the other hand, inward rectifying potassium channels are less active, S type anion channels are more active, guard cell Cl⁻ release is stimulated, and guard cell Ca²⁺ concentration is increased [5]. The stomata close as a result of the depolarization of guard cell membrane potential brought forth by these alterations. A smaller stomatal aperture is the consequence of more depolarization with higher CO₂ [6].

To review, when CO₂ levels are high, stomatal aperture typically decreases in the near term. Dysregulation of stomatal aperture, stomatal index, or density of stomata may lead to a decline in gs over time. According to Gray et al. [7] the HIC gene produces a potential *3-keto acyl coenzyme A synthase*, a protein that negatively regulates stomatal development. In contrast to the majority of plants, mutant hic plants respond to high CO₂ levels by increasing their stomatal density by as much as 42%. This is likely because high CO₂ levels impair the signal transduction system that normally controls stomatal patterning [7]. Additionally, research by Lake et al. [8] suggests that mature leaves limit stomatal development in response to CO₂. Immature leaves get signals from mature leaves about levels of CO₂ and adjust their stomatal growth according on those signals [8]

Abscisic acid, ethylene, and jasmonic acid may all have a role in long-distance signaling, according to new research using Arabidopsis mutants; moreover, there are separate controls for adaxial and abaxial stomatal responses [9]. Many Arabidopsis thaliana ecotypes and a broad range of other species have their stomatal

density reduced during growth in environments with increased CO₂ levels (Hetherington & Woodward 2003; Woodward et al., [10]. In FACE studies, the drop in gs with increased CO₂ was not due to a substantial change in stomatal density Reid et al., [11] Marchi et al., [12] Tricker et al., [13].

3. RESPONSE OF PHOTOSYNTHESIS TO INCREASED CO₂ CONCENTRATIONS

3.1 Quantity of Response

Ainsworth and Rogers (2007), Nowak et al. [14] Ainsworth and Long (2005), Long et al. (2004) and are just a few of the several publications that have evaluated the reactions of photosynthesis to increasing CO₂ concentrations (Ainsworth and Rogers, 2007). Ainsworth and Long (2005) and Ainsworth and Rogers (2007) found that the stimulation magnitude varied across the range of plant functional types (PFTs) is as follows: trees and C₃ grasses exhibit the highest PFTs, shrubs, C₃ and C₄ crops, and legumes have moderate PFTs, and C₄ grass has the lowest PFTs (despite a negative response). Since plant species, PFTs, and environmental factors such as nutrition and water resource availability all play a role, the stimulation by high CO₂ levels may vary greatly [15].

Also, C₄ plants may enhance CO₂ fixation while using light more efficiently by avoiding photorespiration (Long et al., 2006; von Caemmerer and Furbank, [16]. High internal and stomatal resistances, increased levels of starch, diluted concentration of chlorophyll, and nitrogen restriction are other potential outcomes of photosynthetic adaptation to elevated CO₂ levels [17]. Carbohydrate accumulations, including starch size and chloroplast number, may be augmented under increasing CO₂, in part because the carbon substrate increases [18]. Nevertheless, a decrease in photosynthetic capability may result from physical damage to chloroplasts or feedback inhibition caused by an overabundance of carbohydrates [19].

how plants adapt their photosynthetic processes to higher concentrations of carbon dioxide (CO₂), especially when there is a nitrogen availability deficit and the plants need to store more CO₂ for the long terms.

3.2 Molecular Mechanisms

Function of Rubisco Duarte et al. (2014), Ainsworth Rogers (2007), and Long et al. (2004)

all found that short-term increased CO₂ stimulated photosynthesis in C₃ species. This finding was confirmed under almost all experimental settings, including FACE. Acclimation of photosynthesis capability may occur with prolonged exposure to high CO₂ or other constraints; the specifics of this process depend on the species, stage of development, and environmental factors. According to research by Eichelmann et al. [20] the enzyme Rubisco regulates the rate of carbon fixation. Here, we provide a brief overview of the five main hypotheses that might account for Rubisco's reaction to increasing CO₂ levels.

An instantaneous rise in the Rubisco carboxylation velocity may be caused by growing CO₂ levels, since there is an increase in the availability of carbon substrate. (2) Carboxylation and oxygenation are two inherent side characteristics of Rubisco catalytic activity. Ainsworth and Rogers (2007) found that at 25°C, the oxygenation process may consume about one-third of the ribulose 1,5-bisphosphate (RuBP), as the carboxylation rate is approximately 2.2 times higher. Therefore, increased CO₂ levels can promote RuBP carboxylation by up-regulating Rubisco's affinity for CO₂, and inhibit RuBP oxygenation (light-dependent photorespiration) by down-regulating Rubisco's affinity for O₂.

4. RESPIRATORY RESPONSE TO INCREASED CO₂

4.1 Photorespiration

Decreased photosynthesis may result from enhanced photo peroxidation in chloroplasts, which can destroy chlorophyll and cause membrane disintegration. The photosynthetic machinery may be protected from oxidative stress if photorespiration is limited by increased CO₂, which may also lower H₂O₂ products Zinta et al., 2014; Watanabe et al., [21].

Since photorespiration prevents photo-oxidation, it may work by increasing glutamine synthetase (GS2) activity, which recycles ammonia and reduces photo-oxidation and photo-inhibition (Zinta et al., 2014). Another problem arises because, as CO₂ levels rise, photorespiration decreases, which might counteract the protective effect and cause more photo-oxidation than the increased rate of carboxylation can sustain. Research into climate change adaptation strategies, such as modified photorespiration

bioprocess manipulation, is necessary to address this challenge [22].

5. INCREASED CO₂ INTERACTIONS WITH A VARIETY OF ABIOTIC STRESSORS

Global warming is a growing problem that will soon affect many parts of the biosphere, including humans, animals, and plants. Extreme weather events are a known risk to global food supplies because they might disrupt agricultural output, which is highly sensitive to weather patterns. Significant fluctuations in worldwide temperatures and precipitation patterns have the potential to render agricultural plants susceptible to both biotic and abiotic stresses, including pathogens, vegetation, ozone, cold, intense light, nutrients, and drought.

In addition, shifting weather patterns may need new farming techniques, which in turn affect the dynamics between plants and their parasites [23]. For example, soil-borne illnesses are more common in areas where practices such as stubble retention and limited tillage are utilized to maintain soil moisture [24].

One of the main causes of global warming is atmospheric CO₂. Carbon dioxide levels in the atmosphere increased from 280 ppm to 400 ppm throughout the last century, coinciding with industrialization and deforestation. By the end of this century, experts anticipate that these levels will reach 730 to 1000 ppm [1,25]. More and more evidence points to the fact that eCO₂, in addition to its involvement in climate change, may impact agricultural productivity in a number of ways, both directly and indirectly. While it is known that eCO₂ can have positive effects on crop production, especially in C₃ plants, these effects are highly dependent on nutrient and water availability and were mainly studied when stresses weren't present (AbdElgawad et al., 2016; Becklin et al., [26] Ghini et al., [27] Gray and Brady, [28].

Depending on the sort of interactions between plants and parasites, eCO₂ may have negative, neutral, or even favorable effects on plant health. As shown in numerous studies (Jwa and Walling, 2001; Górial et al., [29] Mcelrone et al., [30] Zavala et al., [31,32] Kobayashi et al., [33] Sharma et al., 2016; Xie et al., [34] Váry et al., 2015;), eCO₂ can directly and/or indirectly affect many aspects of host and parasite biology during plant-biotic interactions.

6. CONCLUSION

1. In general, when CO₂ levels are high, the Anet goes up. However, the specific beneficial effects on different plant functional groups and species are quite species and functional group dependent. For example, although C₃ species are predicted to be stimulated by increasing CO₂, C₄ plants are the only ones that would experience this effect under water scarcity circumstances (caused by the CCM). The CO₂ enrichment of Rubisco improves its ability to fix carbon, which is a dual feature. Nevertheless, in the presence of carbon sink restrictions and/or a reduced ATP:ADP ratio, as well as diluted N and excessive photosynthate buildup in response to continuously increasing CO₂, the photosynthetic capability may be downregulated.
2. While the amounts of structural carbohydrates, lignin, and lipids stay relatively constant. On the other hand, CO₂ enrichment improves the concentration of some secondary metabolites such phenolic acids, tannins, and phenylpropanoids. Increased CO₂ levels may reduce isoprene emissions because photosynthetic metabolism requires ATP and NADPH for equilibrium.
3. Under various environmental conditions, elevated CO₂ may reduce the negative impacts of abiotic stresses by promoting antioxidant defense metabolism, decreasing photorespiration, increasing resource use efficiency, and improving photosynthesis.

7. POSSIBLE USES IN LIGHT OF FUTURE CLIMATE CHANGE

Key metabolic processes in living organisms and the feedback loops between them may be affected by future climate change. To combat future climate change, we may also use biotechnological instruments like the protective function against ROS.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. IPCC Climate change synthesis report RK. Pachaur, RK, Meyer LA. (Eds.), Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, IPCC, Geneva, Switzerland. 2014;151.
2. NASA GCC. Global climate change: Vital signs of the planet. NASA Climate Factsheet; 2014.
3. Singh A, Agrawal M. Effects of ambient and elevated CO₂ on growth, chlorophyll fluorescence, photosynthetic pigments, antioxidants, and secondary metabolites of *Catharanthus roseus* (L.) G Don. grown under three different soil N levels. Environmental Science and Pollution Research. 2015;22:3936-3946.
4. Matros A, Amme S, Kettig B, BUCK-SORLIN GH, Sonnewald UWE, MOCK HP. Growth at elevated CO₂ concentrations leads to modified profiles of secondary metabolites in tobacco cv. SamsunNN and to increased resistance against infection with potato virus Y. Plant, Cell & Environment. 2006;29(1):126-137.
5. Hanstein SM, Felle HH. CO₂-triggered chloride release from guard cells in intact fava bean leaves. Kinetics of the onset of stomatal closure. Plant Physiology. 2002; 130(2):940-950.
6. Raschke K, Shabahang M, Wolf R. The slow and the quick anion conductance in whole guard cells: their voltage-dependent alternation, and the modulation of their activities by abscisic acid and CO₂. Planta. 2003;217:639-650.
7. Gray JE, Holroyd GH, Van Der Lee FM, Bahrami AR, Sijmons PC, Woodward FI, Hetherington, AM. The HIC signalling pathway links CO₂ perception to stomatal development. Nature. 2000; 408(6813): 713-716.
8. Lake JA, Quick WP, Beerling DJ, Woodward FI. Signals from mature to new leaves. Nature. 2001;411(6834):154-154.
9. Lake JA, Woodward FI, Quick WP. Long-distance CO₂ signalling in plants. Journal of Experimental Botany. 2002;53(367):183-193.
10. Woodward FI, Lake JA, Quick WP. Stomatal development and CO₂: ecological consequences. New Phytologist. 2002;477-484.

11. Reid CD, Maherali H, Johnson HB, Smith SD, Wullschlegel SD, Jackson RB. On the relationship between stomatal characters and atmospheric CO₂. *Geophysical research letters*. 2003;30(19).
12. Marchi S, Tognetti R, Vaccari FP, Lanini M, Kaligarič M, Miglietta F, Raschi A. Physiological and morphological responses of grassland species to elevated atmospheric CO₂ concentrations in FACE-systems and natural CO₂ springs. *Functional plant biology*. 2004; 31(2):181-194.
13. Tricker J., Trewin H, Kull O, Clarkson GJ, Eensalu E, Tallis MJ, Taylor G. Stomatal conductance and not stomatal density determines the long-term reduction in leaf transpiration of poplar in elevated CO₂. *Oecologia*. 2005;143:652-660.
14. Nowak RS, Ellsworth DS, Smith SD. Functional responses of plants to elevated atmospheric CO₂—do photosynthetic and productivity data from FACE experiments support early predictions?. *New phytologist*. 2004;162(2):253-280.
15. Markelz RC, Strellner RS, Leakey AD. Impairment of C₄ photosynthesis by drought is exacerbated by limiting nitrogen and ameliorated by elevated [CO₂] in maize. *Journal of experimental botany*. 2011;62(9):3235-3246.
16. von Caemmerer S, Furbank RT. The C₄ pathway: an efficient CO₂ pump. *Photosynthesis research*. 2003; 77:191-207.
17. Teng N, Jin B, Wang Q, Hao H, Ceulemans R, Kuang T, Lin J. No detectable maternal effects of elevated CO₂ on *Arabidopsis thaliana* over 15 generations. *PLoS One*. 2009;4(6):e6035.
18. Teng N, Wang J, Chen T, Wu X, Wang Y, Lin J. Elevated CO₂ induces physiological, biochemical and structural changes in leaves of *Arabidopsis thaliana*. *The New Phytologist*. 2006;172(1):92-103.
19. Aranjuelo I, Cabrera-Bosquet L, Morcuende R, Avice JC, Nogues S, Araus JL, Perez P. Does ear C sink strength contribute to overcoming photosynthetic acclimation of wheat plants exposed to elevated CO₂? *Journal of Experimental Botany*. 2011;62(11):3957-3969.
20. Eichelmann H, Talts E, Oja V, Padu E, Laisk A. Rubisco in planta k cat is regulated in balance with photosynthetic electron transport. *Journal of experimental botany*. 2009;60(14):4077-4088.
21. Watanabe CK, Sato S, Yanagisawa S, Uesono Y, Terashima I, Noguchi K. Effects of elevated CO₂ on levels of primary metabolites and transcripts of genes encoding respiratory enzymes and their diurnal patterns in *Arabidopsis thaliana*: possible relationships with respiratory rates. *Plant and Cell Physiology*. 2014; 55(2):341-357.
22. Moroney JV, Jungnick N, DiMario RJ, Longstreth DJ. Photorespiration and carbon concentrating mechanisms: two adaptations to high O₂, low CO₂ conditions. *Photosynthesis research*. 2013; 117:121-131.
23. Furlong MJ, Zalucki MP. Climate change and biological control: the consequences of increasing temperatures on host-parasitoid interactions. *Current opinion in insect science*. 2017;20:39-44.
24. Kazan K, Gardiner DM. *Fusarium crown rot caused by Fusarium pseudograminearum in cereal crops: recent progress and future prospects*. *Molecular plant pathology*. 2018;19(7):1547-1562.
25. Meehl GA, Covey C, Delworth T, Latif M, McAvaney B, Mitchell JF, Taylor KE. The WCRP CMIP3 multimodel dataset: A new era in climate change research. *Bulletin of the American meteorological society*. 2007;88(9):1383-1394.
26. Becklin KM, Walker SM, Way DA, Ward JK. CO₂ studies remain key to understanding a future world. *New Phytologist*. 2017;214(1):34-40.
27. Ghini R, MacLeod RE, Santos MS, Silva CE. Elevated atmospheric carbon dioxide concentration increases eucalyptus plantlets growth and reduces diseases severity. *Procedia Environmental Sciences*. 2015;29:206-207.
28. Gray SB, Brady SM. Plant developmental responses to climate change. *Developmental biology*. 2016;419(1):64-77.
29. Górial MM, Ghini R, Bettio W. Elevated atmospheric CO₂ concentration increases rice blast severity. *Plant Pathol*. 2013; 38:253-257
30. Mcelrone AJ, Reid CD, Hoye KA, Hart E, Jackson RB. Elevated CO₂ reduces disease incidence and severity of a red maple fungal pathogen via changes in host physiology and leaf chemistry. *Global Change Biology*. 2005;11(10):1828-1836.

31. Zavala JA, Casteel CL, DeLucia EH, Berenbaum MR. Anthropogenic increase in carbon dioxide compromises plant defense against invasive insects. Proceedings of the national academy of sciences. 2008; 105(13):5129-5133.
32. Zavala JA, Nability PD, DeLucia EH. An emerging understanding of mechanisms governing insect herbivory under elevated CO₂. Annual review of entomology. 2013; 58:79-97.
33. Kobayashi T, Ishiguro K, Nakajima T, Kim HY, Okada M, Kobayashi K. Effects of elevated atmospheric CO₂ concentration on the infection of rice blast and sheath blight. Phytopathology. 2006;96(4):425-431.
34. Xie H, Zhao L, Yang Q, Wang Z, He K. Direct effects of elevated CO₂ levels on the fitness performance of Asian corn borer (Lepidoptera: Crambidae) for multigenerations. Environmental Entomology. 2015;44(4):1250-1257.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/115199>