



Roving Survey of Root Rot Disease of Pea Caused by *Fusarium solani* f. sp. *pisi* in Rajasthan, India

**Yogita Nain ^{a*}, Nitin Chawla ^a, Priyanka Kumari Meena ^a,
R. P. Ghasolia ^b, S.K. Goyel ^b and Pooja Nain ^c**

^a Department of Plant Pathology, Rajasthan Agricultural Research Institute, Durgapura, Jaipur, 302018, India.

^b Department of Plant Pathology, SKNAU, Jobner, Jaipur, 302018, India.

^c Department of Soil Science, GBPUA and T, Pantnagar, Uttarakhand, 263145, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i62544>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116386>

Original Research Article

Received: 16/03/2024

Accepted: 20/05/2024

Published: 30/05/2024

ABSTRACT

Pea, a vegetable crop grown in cool regions of the world. It has a great economic importance in India. The survey conducted in the major pea-growing regions of Rajasthan during the *Rabi* season of 2021-22 highlighted root rot as a significant issue caused by *Fusarium solani* f. sp. *pisi*. The study covered areas in Jaipur, Sikar, Nagaur, Ajmer, and Alwar regions. To assess disease incidence, researchers randomly selected 25 square meter areas at four spots in each field, totaling 100 square meters per field. In each village, fields were chosen, and plants displaying typical root rot symptoms were collected for further analysis.

*Corresponding author: E-mail: nainyogita9@gmail.com;

Cite as: Nain, Y., Chawla, N., Meena, P. K., Ghasolia, R. P., Goyel, S., & Nain, P. (2024). Roving Survey of Root Rot Disease of Pea Caused by *Fusarium solani* f. sp. *pisi* in Rajasthan, India. *Journal of Experimental Agriculture International*, 46(6), 917–923. <https://doi.org/10.9734/jeai/2024/v46i62544>

Upon examination, an average disease incidence of 23.68 percent was recorded across the surveyed districts. The highest incidence was observed in Jaipur at 30.85 percent, while the lowest was in Nagaur at 16.91 percent. To further understand the pathogen responsible for root rot, samples were collected and brought to the laboratory for isolation. This comprehensive survey provides valuable insights into the prevalence of root rot in pea crops across different regions of Rajasthan, aiding in the development of effective management strategies.

Keywords: *Fusarium solani* f. sp. *pisi*; survey; pea; root rot.

1. INTRODUCTION

Pea is a top vegetable crop grown in cool regions of the world. It is used as a vegetable (green pea) and pulse (dried pea). Duke [1] reported that pea is four types i.e. Garden pea (*P. Sativum* sp. *Hortense* (Asch. And Graebn.)), dry pea (*P. Sativum* sp. *Arvense* (L.) Poir.), edible podded pea (*P. Sativum* sp. *Macrocarpon*), and early dwarf pea (*P. Sativum* var. *Humile*). It is a diploid species of the legume family. Pea is a crop in tropical and subtropical regions. The protein content of pea ranges from 15.5-39.7 per cent, fresh 100g green pea contain 44 calories, 75.6 per cent water, 6.2 g protein, 0.4 g fat, 16.9 g carbohydrate, 2.4 g crude fibre, 0.9 g ash, 32 mg Ca, 102 mg P, 1.2 mg Fe, 6 mg Na, 350 mg K, 405 µg-β carotene equivalent, 0.28 mg thiamine, 0.11 mg riboflavin, 2.8 mg niacin and 27 mg ascorbic acid; while dried pea contain 10.9 per cent water, 22.9 per cent protein, 1.4 per cent fat, 60.7 per cent carbohydrate, 1.4 per cent crude fiber and 2.7 per cent ash [1-3].

Peas are cultivated both as a vegetable and as a pulse crop, often intercropped with cereals to enhance soil fertility through nitrogen fixation in their root nodules. This crop holds significant economic importance in India, where it ranks fourth in terms of area (10.53%) and fifth in production (6.96%) globally [4].

Root rot disease poses a significant threat to pea cultivation, particularly in the third A zone of Rajasthan, causing substantial losses during the initial plant establishment. The root rot disease incited by *Fusarium solani* (Mart.) Appel and Wollweber f. sp. *pisi* (E. R. John) Snyder and Hansen is a serious disease of pea (*Pisum sativum* L) throughout the world. *Fusarium solani* f. sp. *pisi* produces three types of microscopic spores. Microconidia are small, one-celled and elliptical. Macro conidia are much larger, septate, and slightly curved. Chlamydospores are thick-walled and rounded. The chlamydospores are resistant to extreme environmental conditions and can persist in soil for 5 years or more in the

absence of pea. Root decay generally begins on the finer feeder roots and progresses gradually to the main tap root of the plant. In some cases, all roots are destroyed, leaving only remnants below the attachment of the seed. Root rot in peas is characterized by cortical decay and a brilliant red discoloration of vascular tissues in the root [5]. Pea plants infected with *Fusarium solani* f. sp. *pisi* show poor growth, yellowing, and drying of foliage with the partially or fully damaged root system [6]. Looking to this, the survey was made to know the occurrence of this important disease in Rajasthan region.

2. MATERIALS AND METHODS

Based on the survey conducted in Rajasthan during *Rabi* season 2021 to assess the distribution and prevalence of root rot of peas, the survey covered several districts including Jaipur, Sikar, Ajmer, Nagaur, and Alwar. In each district, two tehsils were selected. Within each tehsil, four villages were chosen. From each village, two fields were selected for the study. Disease incidence was recorded by counting both diseased and healthy plants in randomly selected 25 square meter areas at four random spots within each field, totaling 100 square meters per field. In each village, plants displaying typical symptoms of root rot were collected from the surveyed areas. These samples were carefully labeled and placed in paper bags for transportation to the Rajasthan Agricultural Research Institute, Durgapura, Jaipur laboratory. In the laboratory, efforts were made to isolate the pathogen responsible for root rot from the collected plant samples. This step is crucial for identifying the specific causative agent of the disease.

Based on observation, the disease incidence was calculated by following formula (Horsfall and Cowling, 1978).

$$\text{Per cent disease incidence} = \frac{\text{Number of diseased plants}}{\text{Total number of plants observed}} * 100$$

3. RESULTS AND DISCUSSION

A survey was conducted in major pea growing region of Rajasthan during *Rabi* season of 2021-22. It was concluded that root rot caused by *Fusarium solani* f. sp. *lisi* is fundamental obsessive issue for the major part of area of Jaipur, Sikar, Nagaur, Ajmer and Alwar regions. An average disease incidence of 23.68 per cent was recorded in these surveyed districts. The higher disease frequency of 30.85 % was assessed in Jaipur while least in Nagaur location (16.91 %).

The disease made its appearance at 21-90 days old crop. Initially, the leaves of infected pea plants turn light pale in colour starts drying and finally died. The severity of the disease was highest in Jaipur region which ranged from 30.85 to 29.64 per cent with an average of 30.25 per cent followed by Ajmer, Alwar, Sikar and Nagaur district.

Jaipur: Jobner and Mozmabad tehsils in the Jaipur district were selected for the survey. In Jobner tehsil four village *i.e.* Kalkh, khejrawas, Bobas and Hingoniya were chosen, while in Mozmabad tehsil four village *i.e.* Boraj, Jhag, Sewa, Gangati village were chosen. Four fields were chosen at random in each community for a roving survey. The average disease incidence in Jaipur district was 30.25 per cent with the highest average disease incidence in Jobner tehsil (30.85%) followed by Mozmabad tehsil (29.64%).

Sikar: Dhod and Sikar tehsils in the Sikar district were selected for the survey. In Dhod tehsil four villages *i.e.* Mandota, Bidoli, Arjunpura, and Banura were chosen, while in Sikar tehsil four villages *i.e.* Gokulpura, Dujod, Palwas, and Kasli village were chosen. Four fields were chosen at random in each community for a roving survey. The average disease incidence in Sikar district was 21.48 per cent with the highest average disease incidence in Dhod tehsil (22.59%) followed by Sikar tehsil (20.37%).

Nagaur: Nawa and Merta tehsils in the Nagaur district were selected for the survey. In Nawa tehsil four villages *i.e.* Lohrana, Maroth, Rajliya and Piprali were chosen, while in Merta tehsil four villages *i.e.* Prithvipura, Satlas, Sogawas, and Bhuriyasani were chosen. Four fields were chosen at random in each community for a roving survey. The average disease incidence in Nagaur district was 16.91 per cent with the

highest average disease incidence in Nawa tehsil (17.93%) followed by Merta tehsil (15.90%).

Ajmer: Ajmer and Kishangarh tehsils in the Ajmer district were selected for the survey. In Ajmer tehsil four villages *i.e.* Nand, Kayampura, Gegal, and Picholiya were chosen, while in Kishangarh tehsil four villages *i.e.* Moondoti, Salemabad, Raghunathpura, and Silora village were chosen. Four fields were chosen at random in each community for a roving survey. The average disease incidence in Ajmer district was 26.77 per cent with the highest average disease incidence in Ajmer tehsil (27.77%) followed by Kishangarh tehsils (25.76%).

Alwar: Rajgarh and Bansur tehsils in the Alwar district were selected for the survey. In Rajgarh tehsil four villages *i.e.* Kaleshgarh, Goth, Berli, and Bhangarh were chosen, while in Bansur tehsil four villages *i.e.* Chula, Harsora, Bhoopsera, and Balawas village were chosen. Four fields were chosen at random in each community for a roving survey. The average disease incidence in Alwar district was 22.99 per cent with the highest average disease incidence in Rajgarh tehsil (24.51%) followed by Bansur tehsils (21.46%).

The root rot of pea caused by *Fusarium solani* is an exigent pathological concern especially in Jaipur, Sikar, Nagaur, Ajmer and Alwar. According to the data in Table 1, Fig. 1 the average disease incidence of 23.68 per cent was recorded in five districts of Rajasthan. Jaipur (30.25%) had the highest disease incidence followed by Ajmer (26.77%), Alwar (22.99), Sikar (21.48%) and Nagaur (16.91%). No area or location in the surveyed region was completely devoid of the root rot of pea.

Similar results were also reported by Swaroop *et al.* [7] that conducted a survey on root rot of pea disease and they recorded incidence observed from 18.22% to 32.43% in the Jobner and nearby areas. Thakur *et al.* [8] also reported the incidence of root rot caused by *Fusarium solani* complex pea growing area in Zone IV of Himachal Pradesh. The disease noticed severe form with incidence of 54.7 per cent at HAREC, Kukumseri whereas, in Zone II and III, the status of the disease remained moderate with an incidence of 17.7 to 35.3 per cent at Bajaura and Palampur, respectively.

Table 1. Per cent disease incidence of root rot of pea incited by *Fusarium solani* f. sp. *lisi* in different regions of Rajasthan

S.N.	District	Tehsils	Name of village surveyed	PDI in surveyed 2 fields of each village	Soil type	Avg. PDI (%) of tehsil	
1	Jaipur	Jobner	1. Kalkh	28.19 (1)	28.88 (2)	Sandy loam	30.85
			2. Khejrawas	36.37 (3)	26.48 (4)		
			3. Bobas	25.81 (5)	31.32 (6)		
			4. Hingoniya	32.38 (7)	37.36 (8)		
	Jaipur	Mozmabad	1. Boraj	26.67 (9)	35.00 (10)	Sandy loam	29.64
			2. Jhag	33.34 (11)	24.14 (12)		
			3. Sewa	31.34 (13)	20.19 (14)		
			4. Gangati	32.37 (15)	26.00 (16)		
Mean						30.25	
2	Sikar	Dhod	1. Mandota	19.45 (17)	20.75 (18)	Sandy loam	22.59
			2. Bidoli	16.71 (19)	25.28 (20)		
			3. Arjunpura	23.00 (21)	26.04 (22)		
			4. Banura	27.47 (23)	21.00 (24)		
	Sikar	Sikar	1. Gokulpura	17.65 (25)	23.34 (26)	Sandy loam	20.37
			2. Dujod	24.25 (27)	19.15 (28)		
			3. Palwas	13.89 (29)	28.58 (30)		
			4. Kasli	17.65 (31)	18.52 (32)		
Mean						21.48	
3	Nagaur	Nawa	1. Lohrana	21.31 (33)	24.14 (34)	Sandy loam	17.93
			2. Maroth	16.15 (35)	10.72 (36)		
			3. Rajliya	24.13 (37)	17.82 (38)		
			4. Piprali	18.11 (39)	13.04 (40)		
	Nagaur	Merta	1. Prithvipura	16.19 (41)	10.46 (42)	Sandy loam	15.90
			2. Satlas	19.43 (43)	12.50 (44)		
			3. Sogawas	15.62 (45)	15.71 (46)		
			4. Bhuriyasani	23.34 (47)	15.90 (48)		
Mean						16.91	
4	Ajmer	Ajmer	1. Nand	24.88 (49)	25.81 (50)	Loamy soil	27.77
			2. Kayampura	32.15 (51)	33.49 (52)		
			3. Gegal	27.93 (53)	24.24 (54)		

S.N.	District	Tehsils	Name of village surveyed	PDI in surveyed 2 fields of each village		Soil type	Avg. PDI (%) of tehsil
	Ajmer	Kishangarh	4. Picholiya	30.29 (55)	23.34 (56)	Loamy soil	25.76
			1. Moondoti	24.28 (57)	23.67 (58)		
			2. Salemabad	21.52 (59)	22.69 (60)		
			3. Raghunathpura	20.06 (61)	26.67 (62)		
			4. Silora	34.26 (63)	22.88 (64)		
						Mean	26.77
5	Alwar	Rajgarh	1. Kaleshgarh	20.45 (65)	18.75 (66)	Clay loam	24.51
			2. Goth	20.00 (67)	26.49 (68)		
			3. Berli	32.36 (69)	27.25 (70)		
			4. Bhangarh	22.67 (71)	28.04 (72)		
	Alwar	Bansur	1. Chula	20.12 (73)	24.00 (74)	Clay loam	21.46
			2. Harsora	22.65 (75)	25.43 (76)		
			3. Bhoopsera	19.56 (77)	26.04 (78)		
			4. Balawas	15.30 (79)	18.52 (80)		
						Mean	22.99
						Overall mean	23.68




Fig. 1. Per cent disease incidence of root rot of pea incited by *Fusarium solani* f. sp. *pisi* in different regions of Rajasthan

Williamson-Benavides *et al.* [9] concluded that root rot is a devastating disease of pea which causes significant yield losses (15–60%) across the world. Similarly, Yan and Nelson (2022) reported that soil type significantly affected disease development, with higher severity in the lighter soils of sandy loam and silt loam compared with clay. Soil type also interacted with *Fusarium* species, in which the maximum severity was observed in sandy loam for *F. solani*.

4. CONCLUSION

Root rot disease is a potentially serious threat to pea production area. Data clearly showed that the root rot of pea caused by *Fusarium solani* is an exigent pathological concern especially in Jaipur, Sikar, Nagaur, Ajmer and Alwar. The average disease incidence of 23.68 per cent was recorded in five districts of Rajasthan. Jaipur (30.25%) had the highest disease incidence followed by Ajmer (26.77%), Alwar (22.99), Sikar (21.48%) and Nagaur (16.91%). No area or location in the surveyed region was completely devoid of the root rot of pea. Since *Fusarium solani* are soil inhabitants and survive long time in soil. The practical implications of research findings on root rot disease in pea crops for farmers, agronomists, are significant. Integrated disease management Implementing IPM strategies can help in controlling root rot disease while minimizing reliance on chemicals. This may involve practices such as biological control using antagonistic microorganisms, crop sanitation, and judicious use of fungicides only when necessary and as part of an overall management plan.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Duke JA. Handbook of legumes of world economic importance. Plenum Press, New York. 1981;199-265.
2. Hulse JH. Nature, composition and utilization of food legumes. In: Muehlbauer F.J. and Kaiser W.J. (eds.), Expanding the production and use of cool Season Food Legumes. Kluwer Academic Publishers. Dordrecht, the Netherland. 1994;77-97.
3. Kadam AM, Chavan SS, Dhutraj DN, Rewale KA. Survey of dry root rot of chickpea incidence in Marathwada region. Journal of Pharmacognosy and Phytochemistry. 2018;7(1S):3004-8.
4. FAO. FAOSTAT. Food and Agriculture Organization of the United Nations, Rome, Italy; 2021.
5. Lin YS, Sun W, Wong PH. Fusarium root rot and wilt of garden peas in Taiwan. Journal of agricultural research of China. 1984;33(4), 395-405.
6. Hamid A, Bhat NA, Sofi TA, Bhat K. A., and Asif, M. 2012. Management of root rot of pea (*Pisum sativum* L.) through bio-agents. African Journal of Microbiology Research. 2012;6(44):7156-7161.
7. Swaroop KR, Ahir RR, Kumar JN. Survey of root rot disease in Jobner vicinity induced by *Fusarium solani* f. sp. *pisi*. Trends in Biosciences. 2012;7(20):3311-3314.
8. Thakur BR, Kumari N, Singh A.. Occurrence of pea root rot/wilt complex disease in Himachal Pradesh. Himachal Journal of Agricultural Research. 2016; 42(2):187-191.
9. Williamson-Benavides BA, Sharpe RM, Nelson G, Bodah ET, Porter LD, Dhingra, A. Identification of *Fusarium solani* f. sp. *pisi* (Fsp) responsive genes in *Pisum sativum*. Frontiers in Genetics. 2020;11: 950.
10. Yan H, Nelson Jr B. Effects of soil type, temperature and moisture on development of *Fusarium* root rot of soybean by *Fusarium solani* (FSSC 11) and *usarium tricinatum*. Plant Disease, (ja). Plant Disease. 2022;106(11), 2974-2983.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/116386>